

Library copy

MARY GUNN LIBRARY
NATIONAL BOTANICAL INSTITUTE
PRIVATE BAG X101
PRETORIA 0001
REPUBLIC OF SOUTH AFRICA

LIBRARY
DIVISION OF BOTANY AND
PLANT PATHOLOGY
Catalogue No.
2 - 3 = 1944
BIBLIOTHEEK
AFDELING PLANTKUNDE EN
PLANTSIEKTEKUNDE

DEPT. VAN LANDBOU EN VISSERVE
NAVORSINGSINSTITUUT VIR PLANTKUNDE
BOTANICAL RESEARCH INSTITUTE
DEPT. OF AGRICULTURE AND FISHERIES

MARY GUNN LIBRARY




0000017866

South African National
Biodiversity Institute

Bothalia.

A RECORD OF
CONTRIBUTIONS
FROM THE
NATIONAL HERBARIUM
UNION OF SOUTH AFRICA
PRETORIA




EDITED BY

I. B. POLE EVANS, C.M.G., M.A. (Cantab.), D.Sc. (Wales), F.L.S.

CHIEF, DIVISION OF BOTANY AND PLANT PATHOLOGY, DEPARTMENT OF AGRICULTURE,
PRETORIA; AND

DIRECTOR OF THE BOTANICAL SURVEY OF THE UNION OF SOUTH AFRICA



PRETORIA

THE GOVERNMENT PRINTING AND STATIONERY OFFICE

1921

PREFACE.

WITH the advent of Union and the ever-increasing demands made by the public for information regarding the vegetable resources of the country, its plant poisons, and plant pests, considerable stimulus has been given to botanical research in South Africa.

One of the results of the progress made has been the establishment of the National Herbarium at Pretoria under the Department of Agriculture.

The National Herbarium now embraces all the more important private collections in the country, including those of Mr. E. E. Galpin, the late Mr. H. G. Flanagan, Madame A. Dieterlen's Basutoland herbarium, Miss A. Pegler's Kentani plants, Madame J. Borle's collections from Portuguese East Africa, and Mr. E. G. Bryant's plants from the Prieska and Hay Districts.

It also possesses the largest collection of South African fungi in existence, amongst which may be mentioned a large number of specimens of the superficial ascomycetes affecting our forest trees, and the original collections of Dr. Medley Wood, including a number of type specimens, while to it have been bequeathed the valuable cryptogamic collections of Dr. T. R. Sim; Tyson's collections of marine algae, including his private herbarium, have also been acquired by purchase.

A vast amount of material has thus been accumulated, and the special work which has been done on it in the past few years makes it very necessary and desirable that it should be published and made available to workers in other parts of the country. Consequently it has been decided to issue from time to time as occasion demands a publication which will consist of contributions from the National Herbarium.

In view of the great agricultural development which has taken place in South Africa since Union, and bearing in mind that much of this was due to the policy and influence of the first Union Premier and Minister for Agriculture—the late General Botha—no more fitting name for the official organ of the National Herbarium could be found than that of

“BOTHALIA.”


Bothalia is intended primarily as the medium for the publication of papers and monographs based on the material in the National Herbarium, presenting new facts and items of general interest.

It will include descriptions of new or little known plants, cryptogamic and phanerogamic, and it is hoped will mark a further step in the progress of botanical science in South Africa.

I. B. POLE EVANS.

Pretoria,

1st January, 1921.



Digitized by the Internet Archive
in 2016 with funding from
South African National Biodiversity Institute Libraries

<https://archive.org/details/bothaliavolume101unse>

C O N T E N T S.

Part I: 1921

South African Ascomycetes.....	5
The Genus "Bersama".....	33
A Revision of the African Species of "Sesbania".....	40
The Natal Species of the "Sapindaceae".....	57

Part II: 1922

South African Ascomycetes.....	65
The Thorn Pears.....	83
The Genus Ochna.....	87
The Genus Olinia.....	97
The Genus Cyclopia, Vent.....	105

Part III: 1922

Revision of South African Species of Rhynchosia.....	113
The Genera Aloe and Mesembryanthemum.....	139
South African Gramineae.....	170
A Fungus of Economic Importance on the Avocado.....	189
Species of Elephantorrhiza in the South African Herbaria.....	187

Part IV: 1924

South African Ascomycetes in the National Herbarium..	195
South African Gramineae.....	222
Index to Volume I.....	305

MARY GUNN LIBRARY



0000017867

South African National
Biodiversity Institute

CONTENTS.



	PAGE
I. SOUTH AFRICAN ASCOMYCETES.....	5
II. THE GENUS "BERSAMA".....	33
III. A REVISION OF THE AFRICAN SPECIES OF "SESBANIA".....	40
IV. THE NATAL SPECIES OF THE "SAPINDACEAE".....	57

SOUTH AFRICAN ASCOMYCETES

IN THE

NATIONAL HERBARIUM.

By ETHEL M. DOIDGE, M.A., D.Sc., F.L.S., Assistant Chief, Division of
Botany and Plant Pathology.

PART I.

Most of the South African fungi described in the early days were collected by MacOwan, Medley Wood, and others, who made a practice of sending a part of their material to Europe for determination and retaining the duplicates; many of these duplicates are now incorporated in the National Herbarium. These specimens are not strictly co-types, as they were not examined by the describers of the species, but since the specimens were all numbered, one rarely finds that they are not identical with the type specimens bearing the same numbers. The same applies to specimens sent to Sydow in more recent years, descriptions of which were published in the "Annales Mycologici."

Many alterations in nomenclature and classification have been introduced of late years, chiefly by V. Höhnelt, Theissen, and Sydow, who have published revised descriptions from the type specimens. In some cases it is possible to add a few details to these descriptions from more abundant and more recent collections, and to record the fungus on other hosts and from new localities.

There are also a number of ascomycetes in the Herbarium which have been previously recorded from other parts of the world, or which are apparently undescribed.

In every case the numbers quoted in square brackets are those of the mycological section of the National Herbarium.

I am indebted to Mr. W. E. Schilz for the preparation of a fine series of sections of the fungi under consideration and for considerable assistance in the translation of some of the descriptions from the original German.

1. *Schneepeia radiata* Doidge.

Syn. *Lembosia radiata* Doidge (South African Microthyriaceae, Trans. Roy. Soc. South Africa, VIII, Part 4, 1920).

On leaves of undetermined shrub (Leguminosae) Rikatli, Portuguese East Africa, Junod [11729].

This fungus was originally described as a *Lembosia*, but by studying sections through the leaf of the host an intramatrical hypostroma may be detected. This consists of a colourless hyphal mass lying between the cuticle and the epidermis, and sometimes penetrating downwards into the epidermal cells. In certain of the epidermal cells compact balls of dark coloured hyphae are formed, and at these points the fungus breaks through the cuticle to form the ascostroma.

The ascostroma is attached at several points and is almost 40 μ high; the hypothecium is thin and colourless.

The intramatrix hypostroma with superficial ascomata is typical of the *Polystomellaceae*; the radial arrangement of the loculi which are attached at several points, the paraphysate asci, and two-celled brown spores are characteristic of the genus *Schneepeia*, to which this fungus must therefore be referred.

2. *Cycloschizon brachylaenae* (Rehm), P. Henn.

Engl. Bot. Jahrb. XXXIII, p. 39 (1902); Syll. Fung. XVII, p. 896.

Syn. *Schneepeia brachylaenae* Rehm, Hedwigia 1901, p. 173.

On living leaves *Brachylaena nerifolia*, Bains Kloof, near Wellington, C.P., 12.11.10, Doidge [1981].

On *Brachylaena dentata*, Port Elizabeth, West [761]; Van Stadens Pass, 9.3.16, Bottomley [9562]; Van Stadens Pass, 17.5.18, Pole Evans [1445].

On *Brachylaena discolor*, Lemana, Zoutpansberg District, 14.8.11, Doidge [1789]; Duikerfontein, Natal, 10.10.11, Moon [1887]; Port Shepstone, Natal, 15.10.12, Pole Evans [5605]; Umgeni, near Durban, 15.11.16, V. d. Bijl [10088].

On *Brachylaena elliptica*, East London, 24.11.17, Doidge [10912].

On *Brachylaena* sp., Rikatli, Portuguese East Africa, September, 1918, Junod [11724].

Sydow [Ann. Myc. XIII (1915), p. 207], described this fungus in detail from the type specimen (Exsic. Rabh. P., F. Eur. et extra-europ., 4264), which agrees in every respect with the material in the National Herbarium. The original is evidently not quite mature, and in this connection Theissen and Sydow writes as follows:—"Die Art unterscheidet sich generisch nur durch die farblosen Sporen von *Dielsiella*: da die Asken des Originals noch nicht ganz ausgereift sind, wäre es nach diesem Exemplar nicht ausgeschlossen, dass die Sporen sich später bräunten; dan fiel die Gattung *Cycloschizon* mit *Dielsiella* (1903) zusammen und letztere müsste gestrichen werden. Zu demselben Ergebnis kam V. Höhnelt (Fragm. n. 634 und 635). . . . Das oben erwähnte Exemplar auf *B. dentata* scheint jedoch völlig ausgereift zu sein. Wir fanden hier schon ausserhalb der Schläuche zahlreiche, stets völlig hyaline Sporen, Wahrscheinlich stellt *Cycloschizon* daher eine gute Gattung dar."

After examining the abundant material detailed above, it is evident that the spores of *Cycloschizon brachylaenae* become fuscous, and later brown, at maturity. The genus *Dielsiella* is therefore identical with *Cycloschizon* and the species *Dielsiella pritzelii* P. Henn., and *D. alyxiae* (Mass) Th. and Syd. become *Cycloschizon pritzelii* (P. Henn.) and *C. alyxia* (Mass), respectively.

An apparently undescribed species on *Catha edulis* also belongs to the genus *Cycloschizon*, and is described below.

3. *Cycloschizon fimbriatum* n. sp.

On living leaves of *Catha edulis*, Letaba Drift, Zoutpansberg District, 6.8.11, Doidge [1798] and [1833]; Rikatli, Portuguese East Africa, 31.5.18, Junod [11682]; Louis Trichardt, 8.4.19, Putterill [11835].

Stromata amphigenous, scattered, circular, 1.5 to 2 mm. diam., surface dull black, crustaceous, with an irregular annular ridge marking the position of the loculi; attached to the leaf by a massive central foot, which is black and pseudo-cellular, arising from a dense hypostroma. The hypostroma consists of closely interwoven and packed hyphae, which completely destroy the palisade cells or the mesophyll cells in the immediate vicinity of the foot and penetrate right through the thickness of the leaf in the intercellular spaces. The increasing pressure of the hypostroma ruptures the epidermis, and at this point the foot is formed.

Except for the central foot, the stroma lies free on the leaf surface. The outer membrane is radial in structure, rather loosely interwoven, and composed of hyphae 3.5-4 μ thick. At the periphery it spreads out into a short fringe of brown, radiating hyphae about 3.5 μ thick; these are undulating, branched, and sometimes more or less

coherent. The central portion of the stroma is sterile, the loculi being arranged in a circle round the centre; loculi linear, about $270\ \mu$ wide and $100-115\ \mu$ high, dehiscing by a longitudinal fissure. Hypothecium pseudo-cellular, rather pale.

Asci paraphysate; ovate when immature, thick walled above, $60\ \mu \times 30-33\ \mu$, and with irregularly tristichous or conglobate spores. Mature asci clavate or ellipsoid, $75-90\ \mu \times 23-26\ \mu$ with distichous spores. Asci stain brick red with iodine; there is no blue reaction.

Paraphyses numerous, hyaline, linear, about $2.5\ \mu$ thick. Spores brown, opaque at maturity, two-celled, slightly constricted, ellipsoid, cells equal or sub-equal, $28-30\ \mu \times 12-13.5\ \mu$.

Cycloschizon fimbriatum Doidge, n. sp.

Stromata amphigena, sparsa, $1.5-2\ \text{mm. diam.}$, superficialia, centro affixa; centro sterili magno $400-500\ \mu$ diam.; irregulariter radiatum ex hyphis septatis $3.5-4\ \mu$ crassis brunneis composita, ambitu hyphis longiusculis radiantibus, $3.5\ \mu$ crassis, flexuosis, ramosis cincta; loculo annulari integro v. interrupto, $270\ \mu$ lato, $100-115\ \mu$ alto, rima longitudinale aperto. Asci paraphysati, clavati v. ellipsoidei, $75-90\ \mu \times 23-26\ \mu$, octospori, paraphysibus numerosis, hyalinis, linearibus, $2.5\ \mu$ crassis. Sporae brunneae, oblongae, medio vel paullo supra medium septatae, leniter constrictae, ad apicem rotundatae, $28-30\ \mu \times 12-13.5\ \mu$.

Hab. in foliis *Cathae edulis*, Zoutpansberg Dist., Transvaal, 6.8.11, leg. Doidge [1797].

4. *Cocconia porrigo* (Cke.) Sacc.

Syll. Fung. VIII, p. 738.

Syn. *Rhytisma porrigo* Cke., Grevillea X, p. 129 (1882).

On leaves of *Olea capensis*, Inanda, August, 1881, Wood No. 660 [10702]; Van Stadens Pass, C.P., 13.11.17, Doidge [10888].

Stromata amphigenous, but mostly epiphyllous, not crowded, round, flat, dull black, up to $5\ \text{mm. diameter}$; central portion raised and rugose, $120-140\ \mu$ thick. In the smaller stromata there is a regular circular locus forming a ring round the centre, and within it a second smaller locus which may be circular, S-shaped, or irregular. In the larger stromata there are several concentric, circular loculi. At the periphery the covering membrane is thin and membranaceous with lobed margins, and consists of radiating, brown, septate hyphae which are $4-4.5\ \mu$ thick. Hypostroma extensive, in and under the epidermis, dark, breaking out at many points and spreading horizontally on the cuticle; thus it forms a layer of short, septate hyphae about $26-30\ \mu$ thick, which becomes gradually thinner towards the periphery. Covering membrane radial in structure, carbonaceous, $30-40\ \mu$ thick. Height of stroma $140-160\ \mu$. Asci broadly ellipsoid, $80-85\ \mu \times 27-30\ \mu$ with distichous or tristichous spores. Paraphyses filiform. Spores dark brown, $30-33\ \mu \times 11-13\ \mu$, two-celled, constricted; loculi unequal, upper cell about $14-16\ \mu \times 13\ \mu$, broader and shorter than the lower cell, which is $16-18\ \mu \times 11\ \mu$.

5. *Cocconia concentrica* Syd.

Ann. Myc. XIII (1915), p. 215.

Syn. *Hysterostomella concentrica* Syd. Ann. Myc. VII (1909), p. 544; Syll. Fung. XXII, p. 556.

Hysterostomella circularis, Har. et Pat. Bull. Mus. Hist. Nat., 1911, p. 368.

On *Trichilia emetica*, Lourenço Marques, 24.9.08, T. R. Sim [508]; Umbelusi R. Portuguese East Africa, 9.11.10, Howard [1019]; Rikatli, P.E.A., September, 1918 Junod [11728].

This fungus has only been recorded from East Africa on *Trichilia emetica*.

It has been described by Sydow (*loc. cit.*) as follows:—On the upper side of the leaf there appear numerous concentric rings of small, black, punctiform bodies, which gradually

develop into small round discs. These form the central part of the stroma, and they later coalesce, forming stouter linear stromata which still show the original ring formation. Meanwhile the concentric ring formation continues at the periphery. The mature stroma is crustaceous in the centre, the original disc-like bodies having become confluent radially as well as laterally, but the concentric circles are still evident; the outer circles still comprise discrete, round, immature discs. The separate stromata vary considerably in length.

A transverse section through a collective stroma, which also passes through single stromata, shows a chain of fruiting bodies with central attachment, and with a sterile central portion.

The single stromata are thus attached to the leaf by their long axis, lying free on both sides of the keel formed by the line of attachment. They are $320-420\ \mu$ high. Beneath the sterile central portion the hypostroma fills the epidermal cells of the host in an area extending $100-120\ \mu$ in diameter. The hypostroma is not continuous, but more or less interrupted; and it produces numberless fine brown hyphae which pierce the cuticle and appear on the surface as a compact stroma. Asci club-shaped cylindrical, thick-walled round the apex, very briefly pedicellate, $45-55\ \mu \times 14-18\ \mu$, eight-spored. Spores distichous, elongated, rounded at both ends, two-celled, brown, $14-18\ \mu \times 4-5\ \mu$. Paraphyses filiform exceeding the asci in length, united above, and forming a somewhat coloured epithecium.

6. *Cocconia capensis* n. sp.

On leaves of *Ochna arborea*, Howiesons Poort, near Grahamstown, 12.7.19, Doidge [12373].

Stromata epiphyllous, round, dull black, minute, up to $1.5\ \text{mm. diam.}$, in appearance rather like a scale insect, centre raised, $130-150\ \mu$ high, margin flat, appressed to leaf surface; centre sterile, surrounded by a single circular loculus, $240-270\ \mu$ broad; covering membrane black, opaque, about $13\ \mu$ thick over the loculus, at the edge becoming somewhat lobed, thin, flat, and light coloured, consisting of fuscous, radiating hyphae, $3-3.5\ \mu$ thick.

Hypostroma very well developed, penetrating right through the leaf and forming a tangled web of hyphae in the intercellular spaces. At certain points the hypostroma becomes much thickened in the epidermal and subepidermal cells, and the hyphal mass becomes dark coloured and cellular, the cells being $6-7\ \mu$ diam.

It breaks through the cuticle and forms a short column of elongated cells, certain of which spread out radially on the leaf surface to form the hypothecium, and others growing to a higher level form the central sterile portion of the stroma, and give rise laterally to the covering membrane. The hypothecium is brown, $30-36\ \mu$ thick and similar in texture to the hypostroma, but the cells are flattened and elongated in a radial direction. There are numerous secondary points of attachment along the axis of the loculus.

Asci clavate, eight-spored, pedicellate, thickened round apex, paraphysate, $65-70\ \mu \times 22-25\ \mu$, they do not react to iodine; paraphyses numerous, filiform. Spores two-celled, brown, $21-23.5\ \mu \times 9-10\ \mu$, slightly constricted; in immature spores the cells are markedly different in form, they are equal in length, but the upper loculus is spherical, $9\ \mu$ diam., and the lower clavate, tapering from $7\ \mu$ at the centre to $4\ \mu$ at the tip. In mature spores the difference is less marked.

Cocconia capensis Doidge nov. sp.

Stromata epiphylla, sparsa, rotundata, atra, minuta usque $1.5\ \text{mm. diam.}$, $130-150\ \mu$ alta, pluries affixa, radiato ex hyphis $3-3.5\ \mu$ cr. contexta; loculo unico annulari, $240-270\ \mu$ lato, hypostromate epidermali et sub-epidermali oriunda. Hypothecium brunneolum $30-36\ \mu$ cr. Asci clavati, paraphysati, octospori, pedicellati, ad apicem incrassati, $65-70\ \mu \times 22-25\ \mu$; paraphysibus numerosis filiformis. Sporae brunneae, sub-clavatae, 1-septatae, constrictae, $21-23.5\ \mu \times 9-10\ \mu$, loculo supero latiore.

Hab. in foliis *Ochnae arboreae*, Howiesons Poort, 12.7.19, leg. Doidge [12373].

7. *Hysterostomina tenella* Syd.

Ann. Myc. XIII (1915), p. 228.

Syn. *Hysterostomella tenella* Syd. Ann. Myc. X (1912), p. 442.

On *Asparagus striatus*, Despatch, near Uitenhage, 23.3.11; Doidge [1241].

On *Asparagus* sp., Kentani, 19.3.12, Pegler [2240]; 3.6.12, Pegler [2362].

Stroma flat, membranous, dull black, 2–10 mm. long, irregular in shape, 70–80 μ high, on both sides of the leaf, or more or less clothing the stem. Loculi very closely crowded, irregularly arranged, dehiscing by a longitudinal slit; covering membrane radial, dark brown to black, opaque, formed of hyphae 3–4 μ thick. Hypothecium very thin, light brown. Asci paraphysate eight-spored, at first broad, oval, with conglobate spores; later becoming elongated, with distichous spores; very thick walled at the upper end, 45–60 $\mu \times$ 16–20 μ . They do not react to iodine. Spores dark brown, strongly constricted, cells spherical, sub-equal, elliptic-oblong, rounded at the ends, 13–16 $\mu \times$ 6–8 μ . Hypostroma not well developed; it can only be detected in the stomata in the form of pegs, 20–25 μ high and 14–16 μ thick, which do not penetrate into the intercellular spaces.



Fig. 1.—Section through loculus of *Macowaniella congesta*, showing hypostroma.

8. *MacOwaniella* Doidge, nov. gen. *Polystomellacearum*.

Stromata superficialia, radiato-contexta ex hypostromati epidermali et sub-epidermali oriunda; loculis linearibus, rectis v. curvulis, inordinate nidulantibus. Asci paraphysati octospori. Sporae phaeodidymae. Mycelium superficiale ramosum, septatum.

This genus differs from *Hysterostomina* in the presence of free mycelium, and from *Lembosiodothis* in the absence of sub-cuticular bands. The hypostroma is deep-seated in the tissues and is not sub-cuticular.

MacOwaniella congesta (Wint.), Doidge.

Syn. *Lembosia congesta* Wint., Syll. Fung. IX, p. 1105 [Wint. Exot. Pilze in Flora, 1884, p. 9].

On leaves and stems of *Carissa arduina*, Lemana, Zoutpansberg, 14.8.11, Doidge [1786]; Isipingo Beach, Natal, 21.5.17, Doidge [10155]; Cango Valley, Oudtshoorn

Dist., 9.11.17, Doidge [10898]; Woodville Forest, George, 11.11.17, Doidge [10929]; Howiesons Poort, Grahamstown, 17.11.17, Doidge [10960]; Lovedale, Alice, 18.11.17, Doidge [10979]; Hoggs Back, C.P., 15.1.18, J. and M. Henderson [11345]; Grahamstown, 1917 [11369].

On *Carissa grandiflora*, Isipingo, Natal, 27.3.18, Bottomley [11380]; Krantzkloof, Natal, 26.5.15, Doidge [8983].

On *Carissa acuminata*, Stella Bush, Durban, 7.4.18, Bottomley [11381].

This fungus was originally described by Winter from material collected by MacOwan.

In his "Lembosia-studien" [Ann. Myc. XI (1913), p. 457], Theissen mentions this species, and dismisses it with a single sentence: "Der Beschreibung Winter's ist nur hinzuzufügen, dass die an hyalinen Hyphenbüschel entstehenden Konidien sichelförmig sind, einzellig, beid endig spitz, farblos, $20\mu \times 2\mu$." However, he omits it from his synopsis of species, and from this one would gather that he was doubtful about its systematic position.

The fungus on *Carissa arduina* is very common throughout the country, and with the abundant material at my disposal I have made a careful study of sections through the host, and find that it has a very abundant and well-developed hypostroma. It must therefore be placed in the Polystomellaceae. It differs from *Hysterostomina* in the presence of fairly well-developed superficial mycelium. I have therefore made it the type of a new genus *MacOwaniella*.

MacOwaniella congesta occurs on the younger branches of *Carissa* spp., less frequently on the leaves. Stromata, superficial, round or elliptic or irregular in shape up to 4 mm. diameter; loculi linear, straight, curved, or flexuous, closely crowded, irregularly arranged, $500-800\mu$ long and $180-220\mu$ broad; less frequently oval, attenuated towards both ends, $280-340\mu \times 160-175\mu$; rarely almost circular, $160-190\mu$ diam., $120-140\mu$ high, dehiscent by a longitudinal slit. Hypothecium pseudo-cellular, colourless or light brown in centre, consisting of thin-walled plectenchyma. Covering membrane radial in structure.

Hypostroma strongly developed, penetrating deep into the tissues, forming closely packed masses of interwoven hyphae in the intercellular spaces and in the cells. These are especially evident in the epidermal and sub-epidermal cells under the fruiting bodies, where at intervals the hyphal mass becomes thick walled and dark brown to black in colour, and is connected with the ascostroma by slender, brown to black, fungous pegs about 6μ thick, which pierce the cuticle. At or near the point where the cuticle is pierced the dark hyphae often penetrate laterally into the cuticle for a short distance.

Asci oblong to clavate, attenuated into a short foot, eight-spored, paraphysate, $60-70\mu \times 21-26\mu$. Paraphyses filiform, often forked at the apex, tips more or less conglutinate. Spores conglobate or sub-distichous, oblong, 1-septate, constricted, fuscous when mature, $17-18\mu \times 7-9\mu$.

Mycelium on the leaves well developed, fuscous, undulating, abundantly branched, and anastomosing, with a few sessile, dark brown, hemispherical hyphopodia, $6.5-8\mu \times 6.5\mu$ on the primary hyphae; mycelium on the branches not so well developed, hyphae straighter, more slender, $3.5-5\mu$ thick, branches parallel, with occasional elongated reticulations.

9. *Asterodothis solaris* (K. and Cke.) Theiss.

Ann. Myc. X (1912), p. 179; Ann. Myc. XIII (1915), p. 232.

Syn. *Asterina solaris* K. and Cke., Grevillea IX, p. 33; Syll. Fung. I, p. 42, on

Olea verrucosa, South Africa, Rabh. W., F. Eur. 365.

Lembosia albersii P. Henn., Bot. Jahrb. XXVIII, p. 39; Syll. Fung. XVII, p. 897, on *Elaeodendron* sp., East Africa.

Seyesia elegantula Syd., Engl. Bot. Jahrb., 1910, p. 463; Syll. Fung. XXII, p. 522, on *Xymalos* sp., Uganda.

On *Olea verrucosa* leg. MacOwan, 3991, Rabh. F. Eur. 3651; Wellington, C.P., 10.11.10, Doidge [1033]; Langholm Estates, Bathurst Dist., 14.7.19, Doidge [12346]; Barberton, 30.2.15, Thornicroft [8826].

On *Olea exasperata*, Belmont Valley, Grahamstown, 15.1.17, Doidge [10955].

On *Olea capensis*, Van Stadens Pass, 17.5.18, Pole Evans [11444]; Hoggs Back, C.P., 15.1.18, J. and M. Henderson [11339]; Paddock, Natal, 22.12.13, V. d. Bijl [8875].

On *Olea woodiana*, East London, 24.11.17, Doidge [10902].

On *Olea laurifolia*, Hoggs Back, C.P., 15.1.18, J. and M. Henderson [11343]; Kirstenbosch, December, 1916, Glover [10043]; Howiesons Poort, Grahamstown, 12.7.19, Doidge [12387]; Fort Cunynghame, March, 1915 [8896]; Schwarzwald, Victoria East, C.P., 11.8.15, V. d. Bijl [9464].

On *Elaeodendron croceum*, Kentani, 20.7.12, Pegler [2533]; Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12288]; Knysna, 3.6.12, Pienaar [2435].

On *Elaeodendron capense*, Grahamstown, 13.12.11, Burt-Davy [2070]; Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12295].

Stromata amphigenous but more frequently epiphyllous, forming dark brown to black, orbicular spots up to 2–4 mm. diam. These are usually quite distinctly radiate and fibrillar in structure, even to the naked eye.

The hypostroma may be detected within the leaf, lying between the epidermis and the palisade cells, and pushing down between the latter. In the *Elaeodendron* spp. there is more than one row of epidermal cells; but even then the internal mycelium penetrates between all the rows of cells as far as the palisade tissue. At certain points the hypostroma becomes much thickened and a hyphal mass is produced above the palisade cells which gives rise to a short, dark, cylindrical column of perpendicular prosenchymatous structure. This column or foot ruptures the epidermis and the hyphae composing it flatten out horizontally and form the stroma; certain of the hyphae form a hypothecium about 10 μ thick, and others growing to a higher level produce the radial covering membrane. The illustration of the stroma of *Asterodothis solaris* in the *Annales Mycologici* [13 (1915), Tafel I, Fig. 10] is somewhat misleading, as it depicts the loculi in close contact with the central column. This is not always the case, as in the large number of sections examined of the fungus on various hosts the central part of the stroma was sterile and the loculi were at some little distance, closely crowded, forming a sort of crown round central point or being irregularly scattered about the centre.

The central part of the stroma as seen in a surface section may be parenchymatous in structure, in which case the loculi show as small mounds; but frequently the loculi appear as dense black bodies borne on radiating hyphal strands which are more or less connected with one another. The loculi (external measurement) are round 200–250 μ diam., or oval, 250–300 $\mu \times$ 140–200 μ ; they dehisce irregularly by a stellate or longitudinal fissure.

In section the loculi are 120–160 μ high, and are connected with the intramatrical hypostroma by numerous dark coloured fungus pegs which pass through the stromata. This fact is not mentioned by Theissen.

Radiating from the stroma are numerous hyphae which are almost straight, dark brown, septate, 5–6 μ thick; these often become adnate by their radial walls and form hyphal strands. The hyphae produce branches, which form acute angles with the parent hypha.

On the hyphae are borne numerous outgrowths which may be regarded as rudimentary setae or bristles. These are sometimes very short and almost like hyphopodia in appearance; more frequently they are more or less erect, truncate, and up to 25 $\mu \times$ 7–8 μ . On the hyphae are also borne very dark brown, club-shaped conidia, which are 3–4 septate, blunt at the ends, and 36–42 $\mu \times$ 11–12 μ .

The fruiting layer is flat and the asci parallel, oval to clavate, eight-spored, 70–90 $\mu \times$ 24–28 μ . Paraphyses numerous, filiform. Spores distichous, ellipsoid, reddish-brown, smooth, 27–31 μ long, upper cell 10–12 μ broad, lower cell 8–10 μ broad.

The spores have a peculiar method of germination, the germ tubes appearing on the medial constriction between the two loculi, one on each side.

A fungus with a similar conidia to *Asterodothis solaris* has been collected on *Myrsine melanophleas* [10908] and [8995], but unfortunately neither of these two collections bear mature ascostromata.

10. *Polyrhizon bewsii* n. sp.

On living leaves of *Elaeodendron aethiopicum*, Bisley, near Maritzburg, Natal, 11.4.17, Bews [10087]; 20.7.18, Doidge [11592].

Stromata amphigenous, scattered, raised, round to irregular, about 1 mm. diam., simple or compound, in the latter case 2-3 or more stromata arising close together become confluent. There is no discoloration of the leaf tissues in the neighbourhood of the stroma.

Intramaterial mycelium copious, consisting of colourless hyphae 3.5-4 μ thick, filling the intercellular spaces and penetrating right through the leaf so that stromata are usually formed at corresponding points on the upper and lower surface, one being older than the other. Each stroma is attached to the leaf by a central foot, which is a continuation of the brown compact hypostroma formed in the epidermal and sub-epidermal cells. This foot is 80-160 μ thick, and is continued into the stroma, forming a dark, cellular hypothecium, around which the loculi are arranged. Apart from the central foot the stromata are unattached, lying free on the leaf surface. In old stromata the asci above the foot break away, leaving a cavity, so that the loculi have the appearance of being arranged in a circle around a sterile centre.

Covering membrane black, carbonaceous, radiating in structure at margin, rugose. Diameter of a single stroma is 700-1000 μ ; height, excluding foot, 200-270 μ .

Asci paraphysate, briefly pedicellate, eight-spored, 70-100 $\mu \times$ 30-45 μ , ovate, thick-walled round apex, with conglobate spores when immature; later becoming more elongated, clavate, with distichous spores. Paraphyses linear, about 3.5 μ thick. Spores two-celled, almost black, opaque, slightly constricted, ellipsoid, upper loculus slightly larger, 30-33 $\mu \times$ 13.5-15 μ .

Polyrhizon bewsii Doidge, nov. sp.

Stromata amphigena, orbicularia, atra, carbonacea, circ. 1 mm. diam., centro affixa, e 1-pluribus ascomatibus circinantibus concreta. Ascomata partialia, pede centrali 80-160 μ cr., epidermide innata, versus marginum radiato-contexta, 700-1000 μ diam., 200-270 μ alto, loculis rotundatis, immersis. Asci paraphysati, clavati, breviter pedicellati, octospori, 70-100 $\mu \times$ 30-45 μ , paraphysibus linearibus, 3.5 μ cr. Sporae didymae, brunneae, opacae, leniter constrictae, elongatae, loculo supero paullo majore, 30-33 $\mu \times$ 13.5-15 μ .

Hab. in foliis *Elaeodendri aethiopici*, Natal, 1.9.17, leg. Bews et Doidge [10087] et [11592].

11. *Placoasterella rehmsii* (P. Henn.) Theiss. et Syd.

Ann. Myc. XIII (1915), p. 237.

Syn. *Asterella rehmsii* P. Henn., Engl. Bot. Jahrb. XVII, p. 114; Sacc. Syll. Fung. XI, p. 257.

On *Aloe arborescens*, Ripplemead, Dohne, Stutterheim Dist., 25.2.15, Bell Edmonds [8899].

On *Aloe mitriformis*, Montague Baths, December, 1915, Pole Evans [9417].

On *Aloe natalensis*, Botanic Gardens, Maritzburg, 8.4.11, Pole Evans [1440].

On *Aloe* sp., New Hanover, Natal, 9.9.13, V. d. Bijl [8892].

This fungus was originally described as occurring in Abyssinia on *Aloe abyssinica* and *Aloe maculata*.

Aerial mycelium not very copious, consisting of brown hyphae, 3.5-5 μ thick, very tortuous, branched, and anastomosing. Stromata scattered or confluent in small groups, either round, 150-200 μ diam., or more frequently elliptic, 200-280 $\mu \times$ 120-150 μ ; in the latter case straight or curved; when two or three are confluent the stroma often appears to be forked; dehiscing by an irregular, round, or elongated fissure.

The cuticle of the leaf is stained with a red pigment where it has been attacked by the fungus, and the discoloration spreads into the palisade cells. The discoloured parts of the leaf tissue are swollen, so that the stromata lie on round, reddish or brown blisters, which are up to 5 mm. diam., and which are often so numerous as to become confluent.

Hypostroma epidermal, filling certain of the epidermal cells with dark hyphal balls, especially under the centre of the superficial fruiting bodies; mycelial strands also penetrate between the palisade cells.

Fruiting body usually unilocular, 60–80 μ high, with a thin black outer wall. Asci oval to clavate, paraphysate, eight-spored, apedicellate, thick-walled round apex, staining brick red with iodine, 36–40 $\mu \times 22\frac{1}{2}$ –25 μ . Spores brown, two-celled, constricted ellipsoid, cells almost equal, upper loculus somewhat broader, 18–23 $\mu \times 8$ –10 μ .

12. *Hysterostoma orbiculata* Syd.

Ann. Myc. XIII (1915), p. 239.

Syn. *Dothidasteromella orbiculata* Syd., Ann. Myc. X (1912), p. 41.

On *Olea verrucosa*, Wellington, C.P., 18.11.10, Doidge [1031]; Port Elizabeth, 17.10.09, West (immature) [1870].

In renaming this fungus, Theissen and Syd. (*loc. cit.*) remark that the species has been described from insufficient material, and that the diagnosis could be much improved after studying well-developed specimens.

There is in the National Herbarium abundant material of the same collection as the type [1031], and much of this is in excellent condition. I am, however, only able to add a few details to the description cited above.

Stromata usually epiphyllous, less commonly hypophyllous, circular in outline, dull black, 4–6 mm. in diameter, carbonaceous; the surface is rough and traversed by deep fissures or clefts. Each stroma is surrounded by a radiating fringe of hyphae, which are septate, 4–6 μ thick, and run parallel to one another. These hyphae are, as a rule, not branched, and are often fused by their lateral walls into hyphal strands, but are not compacted into a stroma.

The central part of the stroma consists of numerous loculi, which are irregularly arranged; they are round to irregular in shape and very closely crowded, 250–300 μ in diameter. The lateral walls are almost perpendicular, the total height of the ascostroma being 240–270 μ . The outer covering is radial in structure, and it breaks down irregularly at maturity.

There appear to be no stomata on the upper surface of the leaf of *Olea verrucosa*, but there are at intervals folds or clefts in the cuticle similar to those beneath which the guard cells are formed on the lower surface. The hypostroma of the fungus is found in the epidermal cells, between the epidermis and the cuticle and in the thickness of the cuticle itself; here it is pseudo-cellular and dark brown. There is also an almost continuous layer of colourless, interwoven hyphae lying between the epidermis and the palisade cells and penetrating into the upper part of the latter. The dark-coloured hypostroma breaks through the leaf, often at the clefts mentioned above; the cuticle is split irregularly in several directions, and through the fissures thus formed the hypostromal tissue grows and produces the fruiting bodies, the hypothecium being in direct connection with the hypostroma at several points under each loculus.

The hypothecium is brown, similar in texture to the hypostroma in the epidermal cells and cuticle, but less dense; it is about 90–100 μ high in the centre of each loculus, where it forms an irregular cushion, on which the asci are borne, and becomes much thinner at the periphery, measuring about 45 μ . Asci elongated, club-shaped, paraphysate, eight-spored, thick walled round the apex, 75–140 $\mu \times 16$ –32 μ . Paraphyses numerous, filiform, 2–3 μ thick, becoming more or less conglutinate at the tips and forming an epithelial layer. The asci do not react to iodine, but stain a brick-red colour. Spores usually distichous, oblong, medially uniseptate, brown, thick walled, 30–34 $\mu \times 12$ –15 μ , upper loculus very slightly broader and more broadly rounded than the lower.




Fig. 2.—Mycelium and spores of *Isipunga areolata*.

13. *Isipinga* Doidge, nov. gen. *Polystomellacearum*.

Mycelium superficialum, ramosum, septatum, hyphopodium. Stromata superficialia ex hypostromate subcuticulari oriunda, pluries affixa. Loculi inordinate dispositi, rotundati vel elliptici (non lineares). Asci paraphysati, octospori. Sporae phaeodidymae.

This genus differs from *Hysterostoma* Theiss. in the presence of a well-developed aerial mycelium, and from *Asterodithis* in the absence of a central column or foot.

Isipinga areolata n. sp.

On leaves of *Euclea natalensis*, Isipingo Beach, Natal, 21.5.17, Doidge [10153]; 13.5.13, Doidge [6647]; near Durban, Natal, 25.5.97, Medley Wood [9520] (Wood No. 6447); 25.5.15, Doidge [8986]; 13.1.18, V. d. Bijl [11363].

Hypophyllous, usually near the reflexed leaf margin, less frequently scattered over the whole under-surface of the leaf, forming circular, dull-black areas, 5–15 mm. diam., often coalescing and forming irregular black blotches, especially along the leaf margins: stromata scattered and visible as small denser black points.

Aerial mycelium well developed, hyphae fuscous, 3.5–5 μ thick, branching freely and anastomosing to form regular, circular, or slightly oval areolae, 30–50 μ diam.; hyphopodia unicellular, darker than the hyphae, ovate or ellipsoid, straight or curved, 8–10 $\mu \times$ 5 μ . Stromata scattered, circular or oval, opaque black, 300–350 μ diam. or about 400 $\mu \times$ 240 μ , formed of radiating cells; arising from a subcuticular hypostroma which breaks through the cuticle to form the stromata in the cavities above the sunken stomata; stromata attached to host at several points by dark brown or black fungous pegs, which are up to 20 μ thick where they penetrate the cuticle; height of fruiting body is 80–100 μ , each stroma containing several round to irregular loculi; the dark covering membrane slopes gradually towards the leaf surface at the circumference of the stroma. Loculi dehisce at maturity by irregular fissures in the covering membrane.

Paraphyses fairly numerous, hyaline, linear, about 3 μ thick, slightly exceeding the asci. Asci eight-spored, ovate or broadly ellipsoid, thick walled round apex, 50–65 $\mu \times$ 33–40 μ , not reacting to iodine, but staining a sort of brick-red colour. Spores tristichous or conglobate, brown, two-celled, 27–32 $\mu \times$ 13–13.5 μ , constricted, upper loculus larger.

Isipinga areolata Doidge, n. sp.

Mycelium hypophyllum, bene evolutum, ramosum, areolatum ex hyphis 3.5–5 μ crassis, anastomosantibus, hyphopodiis continuis, ovatis v. ellipsoideis, rectis v. curvatis, 8–10 $\mu \times$ 5 μ , compositum. Stromata sparsa, rotundata v. elliptica, atra, opaca, 300–350 μ diam.; v. elliptica, circ. 400 $\mu \times$ 240 μ , radiatim contexta, hypostromate subcuticulare oriunda, pluries affixa. Loculi inordinate dispositi, 80–100 μ alti. Asci octospori, ovati vel late ellipsoidei, 50–65 $\mu \times$ 33–40 μ , paraphysate; paraphysibus filiformis, hyalinis, 3 μ crassis. Sporae tristichae v. conglobatae, brunneae, 1-septatae, 27–32 $\mu \times$ 13–13.5 μ , constrictae loculo supero, paullo majore.

Hab. in foliis *Eucleae natalensis*, Isipingo, Natal, 21.5.17, leg. Doidge [10153].

14. *Isipinga contorta* Doidge.

Syn. *Dothidasteromella contorta* Doidge, Trans. Roy. Soc. of S.A. VIII, Part 3, 1920.

On leaves of *Trichocladus ellipticus*, Branders' High Forest, Victoria East, 14.8.15, V. d. Bijl [9462]; Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12298].

Forms small, dull-black, round, or irregular areas on upper surface of the leaves; the leaf tissues become discoloured in the neighbourhood of the fungus, right through to the lower epidermis. Aerial mycelium well developed, primary hyphae long, radiating, fuscous, 3–3.5 μ thick, with unicellular hyphopodia which are alternate or unilateral, subglobose, 6–7 μ diam.; between the radiating hyphae there is a densely interwoven mass of paler secondary hyphae which are very tortuous and crumpled, copiously branched, and anastomosing. Stromata scattered, opaque, black, 250–350 μ diam., with radiating structure, unilocular, attached at several points, and arising from a well-developed subcuticular hypostroma. Asci paraphysate, ellipsoid, or ovate, very briefly stipitate, straight

or curved, $55-80\ \mu \times 20-25\ \mu$; paraphyses numerous, linear, flexuose, or somewhat crumpled. Spores distichous or conglobate, 1-septate, slightly constricted, upper loculus broader, $16-20\ \mu \times 8-10\ \mu$.

15. *Polystomella caulicola* n. sp.

On stems of *Asparagus* sp., banks of Orange River, Aliwal North, 11.1.12, Pienaar [2093].

Stromata caulicolous, usually irregularly elliptic to linear in outline, elongated in a direction parallel with the stem axis, up to 8 mm. long, varying in width, $80-100\ \mu$ high; often confluent and quite irregular in outline, attached at many points; surface rough, dull black. Covering membrane black, carbonaceous, formed of radiating hyphae about $3.5\ \mu$ thick and consisting of numerous *Microthyrium*-like discs fused together at the edges; the membrane covers a number of flattened loculi which are $100-150\ \mu$ diam. and $80-90\ \mu$ high, and are separated by thin walled plectenchyma which are colourless or light brown. Hypothecium delicate, thin, colourless. Asci paraphysate, cylindrical-clavate, eight-spored, briefly pedicellate, $50-70\ \mu \times 18-20\ \mu$. Spores distichous, ellipsoid, hyaline, two-celled, very slightly constricted, cells sub-equal, $17-20\ \mu \times 6-8.5\ \mu$ (not quite mature).

Hypostroma well developed in the epidermis and penetrating into the sub-epidermal cells.

Polystomella caulicola Doidge, n. sp.

Stromata caulicola, elongata, usque 8 mm. longa, $80-100\ \mu$ alta, pluries affixa, membrana una radiato e cellulis $3.5\ \mu$ cr. contexta, carbonacea, ex pluribus quasi-Microthyriis conflata tecta, ex hypothecio epidermali oriunda. Loculi rotundati, $100-150\ \mu$ diam., $80-90\ \mu$ alti, hypothecio tenue, hyaline. Asci paraphysati, cylindracei v. clavati, octospori, breviter pedicellati, $50-70\ \mu \times 18-20\ \mu$. Sporae distichae, ellipsoideae, hyalodidymae, leniter constrictae, loculus subaequalibus, $17-20\ \mu \times 6-8.5\ \mu$ (vix maturis).

Hab. in caulibus *Asparagi* sp., Aliwal North, 11.1.12, leg. Pienaar [2093].

16. *Palawaniella* Doidge, nov. gen. *Polystomellacearum*.

Stromata superficialia, orbicularia, pluries affixa, hypostromate in epidermide bene evoluto, radiato-contexta, carbonacea; mycelio libero nullo; loculis e medio evoluto, plus minusve annulatum dispositis, sed discretis, rotundatis; hypothecium tenue; asci paraphysati, octospori; sporae phaeodidymae.

This genus differs from *Palawania* chiefly in the centrifugal development of the stromata, and in the epidermal, rather than sub-epidermal hypostroma. It differs from *Pleistomella* in the brown, two-celled spores; the loculi are less definitely arranged in concentric rings.

Palawaniella eucleae n. sp.

On living leaves of *Euclea macrophylla*, Howiesons Poort, near Grahamstown, 12.7.19, Doidge [12375].

Stromata epiphyllous, scattered, round, 4-7 mm. diam., loculi developing centrifugally and arranged in more or less definite concentric circles; these show first of all as minute black points, which increase in size up to $160-500\ \mu$ diam.; in the centre the loculi are very closely crowded; towards the circumference they are more scattered, but in either case they are round to oblong, discrete, $80-90\ \mu$ high, adnate with their whole base to the substratum. The outer wall is radial in structure and splits at maturity by irregular stellate or, less frequently, longitudinal fissures.

Hypostroma well developed, formed of fuliginous hyphae which form compact balls in the epidermal cells; hyphae may often be traced running between the palisade cells and the epidermal cells, and so connecting adjacent loculi. There are no stomata on the upper surface of the leaf and the cuticle is not ruptured, the ascus-bearing stroma being connected with the hypostroma in the epidermal cells by numerous fine, colourless filaments, which penetrate the cuticle; these are about $1-1.5\ \mu$ thick. Hypothecium thin, colourless.

Asci paraphysate, eight-spored, very briefly pedicellate, clavate to ovate, thick walled at apex, $43\text{--}57\ \mu \times 23\text{--}27\ \mu$; they do not react to iodine. Paraphyses very numerous, hyaline, filiform $2\cdot5\text{--}3\ \mu$ thick, slightly exceeding the asci. Spores distichous or conglobate, ellipsoid, two-celled, fuscous, later brown, very slightly constricted, cells almost equal, $20\text{--}21\ \mu \times 8\cdot5\text{--}10\ \mu$.

Palawaniella eucleae Doidge, nov. sp.

Stromata ascophora epiphylla, superficialia, sparsa, 4–7 mm. diam., radiato-contexta; hypostromati epidermali; loculi numerosi, e medio evoluti plus minusve concentrice dispositi, sed discreti, rotundati v. elliptici, $160\text{--}500\ \mu$ lati, $80\text{--}90\ \mu$ alti; hypothecio tenue. Asci paraphysati, octospori, brevissime pedicellati, clavati v. ovati, ad apicem incrassati, $43\text{--}57\ \mu \times 23\text{--}27\ \mu$; paraphysibus numerosis, hyalinis, filiformis, sporae distichae v. conglobatae, ellipticae, medio septatae, brunneae, leniter constrictae, loculis subaequalibus, $20\text{--}21\ \mu \times 8\cdot5\text{--}10\ \mu$.

Hab. in foliis *Eucleae macrophyllae*, Howiesons Poort, 12.7.19, leg. Doidge [12375].




Fig. 3.—Section through locus of *Palawaniella eucleae*.

17. *Pleiomastix* *Halleriae* n. sp.

On *Halleria lucida*, Louis Trichardt, 8.4.19, Putterill [11847]; Van Stadens Pass, C.P., 13.11.17, Doidge [10872]; Howiesons Poort, Grahamstown, 17.11.17, Doidge [10959]; Buccleuch, Natal, 17.7.18, Doidge [11585]; Howiesons Poort, 12.7.19, Doidge [12384].

On *Halleria elliptica*, Kentani, 4.12.15, Pegler (Pegler 2348) [9163].

Stromata epiphyllous, superficial, scattered, round, up to 3 mm. diam., obscurely brown, $100\text{--}115\ \mu$ high, attached at many points, radiate in structure, with an opaque, rugose covering membrane $30\text{--}45\ \mu$ thick; hypothecium brown, $30\text{--}36\ \mu$ thick, loculi numerous, densely crowded in concentric circles, but discrete, round, $190\text{--}250\ \mu$ diam., or becoming oval by crowding and then $250\text{--}280\ \mu \times 160\ \mu$. Hypostroma epidermal, but occasionally, in centre of stroma, more deeply seated and invading the palisade cells. Asci ovate, $50\ \mu \times 26\text{--}33\ \mu$ or clavate, $66\text{--}70\ \mu \times 20\text{--}23\ \mu$, thickened at apex, eight-spored; paraphyses hyaline, linear, about $1\cdot5\ \mu$ thick, soon disappearing. Spores distichous or tristichous, at first equally or sub-equally 1-septate, constricted, later muriform, usually transversely 5 (rarely) 7-septate, and with a longitudinal septum running the length of spore, rarely constricted at secondary septa; septa very delicate, plasma granular, hyaline (mature?) $22\text{--}24\ \mu \times 9\text{--}10\ \mu$. Asci stain brick-red with iodine.

Pleiomastix *Halleriae* Doidge, nov. sp.

Stromata ascophora superficialia, sparsa, rotundata, usque 3 mm. diam., obscure brunnea, $100\text{--}115\ \mu$ alta, radiato-contexta, pluries affixa; hypostromate epidermali; strato tegente opaco, $30\text{--}45\ \mu$ crasso; loculi numerosi, dense concentricque dispositi, sed

discreti, 190–250 μ diam., vel elliptici, 250–280 $\mu \times 160 \mu$. Asci ovati, 50 $\mu \times 26$ –33 μ , vel clavati, 66–70 $\mu \times 22$ –23 μ , ad apicem incrassati, octospori, paraphysati; paraphysibus filiformis, mox mucosis praesentibus. Sporae distichae v. tristichae, oblongae, utrinque rotundatae, hyalinae (an semper?) medio v. paullo supra medium septatae, deinde transverse 5, rare 7 septatae, septis tenuissimis, cellulis pluribus v. omnibus septo longitudinali divisus, plasmate granuloso, 22–24 $\mu \times 9$ –10 μ .

Hab. in foliis *Halleriae lucidae*, Louis Trichardt, 8.4.19, leg. Putterill [11847].

18. *Diplochorella amphimelaena* (Mont.) Theiss. and Syd.

Ann. Myc., 1914, p. 277.

Syn. *Dothidea amphimelaena* Mont.

Ann. Sc., Nat. II, Ser. t. XX, p. 372. Syll. Crypt., p. 222.

Homostegia amphimelaena Sacc. Syll. Fing. II, p. 650.

Phyllachora osyridis Cke., Grevillea XIII, p. 64.

Dothidella osyridis Berl. et Vogl., Syll. Fung. IX, p. 1038.

Dothidella osyridis var. *tassiana* Sacc., Syll. Fung. XIV, p. 676.

Microcyclus osyridis Sacc., Ann. Myc. II, 1904, p. 165.

Microcyclus tassianus Syd., Ann. Myc. II, 1904, p. 165.

On *Osyris compressa*, Muizenberg, C.P., 1883, MacOwan (Rabh. Wint. Fung. Eur. 3562) [3902]; Port Elizabeth, 23.3.11, Doidge [1243]; St. James, Capetown, 10.12.11, Pole Evans [1982]; Kloof Neck, Capetown, 24.2.12, Doidge [2155]; Muizenberg, 12.4.12, Beardmore [2292]; Belmont Valley, Grahamstown, 15.11.17, Doidge [10954]; Kalk Bay, January, 1918, Potts [11302].

Compound stromata, circular, 2–3 mm. diam., forming flat, raised discs on both sides of the leaf, with almost vertical edges, silver grey on the surface, thickly beset with erumpent punctiform, dull black bodies about 50 μ diam., which eventually cover the whole surface. The densely crowded loculi develop under the epidermis—the apex of each separately rupturing the epidermis—and become somewhat raised above it. Gradually the part of the epidermis intermediate between the apices of the loculi disappears and the surface becomes altogether black. The leaf is normally 500–550 μ thick, but through the action of the fungus on both leaf surfaces it is often up to 1300 μ thick. The central leaf tissue is colourless and free from fungous stroma to a thickness of 700 μ ; above and below this the stromatic layer develops, which is again differentiated into two layers, the outer composed of the loculi and the inner of the basal hyphal stroma. The first consists of closely crowded, perithecium-like loculi, which are 120 μ high and 85–95 μ diam., and which are more or less connected laterally. From the base of the loculi the hyphae of the vegetative stroma run vertically inwards, forming a fairly sharply defined stromatic plate 120–170 μ thick. The loculi are typically composed of grey-brown, laterally compressed cells which are 14–15 μ long, 5 μ thick, and 8–9 μ broad, and are in close contact by their flat sides. These hyphae sometimes penetrate inwards in more or less compact strands, but only to a limited depth, and the intermediate leaf tissue is not destroyed. Towards the base of the loculi the greater part of the flat cells become polyhedral or regularly elliptic and somewhat smaller and form a sort of wall round the loculi consisting of a few concentric layers of cells; they also fill up the spaces between the loculi; the cells are dark brown and thick walled. There is apparently no stromatic connection between the stromata on the upper and lower leaf surfaces.

Asci apophysate, basal, short cylindrical, 35 $\mu \times 12 \mu$, apedicellate, eight-spored. Spores polystichous, hyaline, two-celled, 12–13 $\mu \times 4 \mu$, constricted at the septum; lower cell twice the length of the upper.




Fig. 4.—*Montagnella maxima*.

19. *Montagnella maxima* Mass.

Gardener's Chronicle, 1899, p. 291, c.i.e.; Syll. Fung. XVI, p. 630.

Ann. Myc. XVI, p. 630; Ann. Myc. XIII (1915), p. 638, Fig. 4.

On *Aloe ferox*, Port Elizabeth, 24.3.12, Doidge [2274]; Uitenhage, 15.5.12, Pienaar [2496]; Graaff-Reinet, 1.4.12, Burt-Davy [5156].

On *Aloe platylepis*, Port Elizabeth, 23.3.11, Doidge [1312].

This fungus forms large raised galls about 1 mm. high; on both sides of the leaf, they are circular or oval and up to 1 cm. diam. The stroma breaks through the epidermis, forming a number of little furrows which are arranged in numerous concentric circles. The vegetative stroma extends perpendicularly downwards into the gall tissue, filling the intercellular spaces with smooth, light brown hyphae $3\frac{1}{2}$ –4 μ thick; before they break through the epidermis these become collected together and form a thick compact web, on which are formed in groups the narrow, almost free loculi. These in section are 135 μ high and 100 μ diam., the stromatic base of each group being 160–240 μ thick. Asci basal, paraphysate, broadly clavate to oval, very briefly pedicellate, 58–68 $\mu \times$ 30–40 μ ; spores brown, tristichous, oblong, 30 $\mu \times$ 6–8 μ , four-celled, constricted in the middle, rounded at both ends, straight.

20. *Elmerococcum peglerae* (Pole Evans) Doidge.

Syn. *Montagnella peglerae* Pole Evans, Ann. Bolus Herb. II, p. 5.

On leaves of *Myrsine melanophleas*, Kentani, 3.4.12, Pegler [2206]; May, 1913, Pegler [6620].

Stromata hypophyllous, black, punctiform, not more than 0.5 mm. diam., attached to the leaf by a central foot which breaks out through a stoma. The central foot gives rise to a thick basal cushion of sterile plectenchyma, on which are borne the small bolster-shaped ascostroma and one or two sub-spherical pycnidia, both being flattened laterally by contact. The stroma is often surrounded by a short sparse fringe of fuscous hyphae, which are septate and 3–3.5 μ thick. Ascostroma 300–430 $\mu \times$ 200–270 μ and about 80–90 μ high, unilocular; wall consisting of polyhedral plectenchyma. Asci paraphysate, oval to clavate, very briefly pedicellate, 60–70 $\mu \times$ 23–27 μ ; paraphyses not numerous, filiform, exceeding the asci. Spores distichous, hyaline, 3-septate, clavate to ellipsoid, not constricted, 27–30 $\mu \times$ 9–10 μ . Pycnidia 160–200 μ diam.; conidia very numerous, minute, bacilloid.

Elmerococcum peglerae Doidge, nov. sp.

Stromata erumpenti-superficialia, centro affixa, orbicularia v. elliptica, usque 5 mm. diam.; ascostroma unica et pycnidia unica vel dua in quaque stromati aggregata. Ascostromata nigra, perithecioidea, elliptica 300–430 μ longa, 200–270 μ lata, 80–90 μ alta. Asci ovati vel clavati, paraphysati, breviter pedicellati, 60–70 $\mu \times$ 23–27 μ . Sporae distichae, hyalinae, 3-septatae 27–30 $\mu \times$ 9–10 μ . Pycnidia rotundata 160–200 μ diam.; conidiis minutis bacillaribus.

Hab. in foliis *Myrsine melanophleas*, Kentani, 3.4.12, Pegler [2206]; May, 1913, Pegler [6620].

21. *Rosenscheldia horridula* n. sp.

On living leaves of *Olea capensis*, Umgeni Beach, Durban, 4.6.12, Pole Evans [3149]; East London, 19.7.19, Doidge [12411].

On *Olea woodiana*, East London, 24.11.17, Doidge [10903].

Stromata usually hypophyllous, less frequently epiphyllous, very numerous, but not large, up to 2 mm. diam. Often larger, irregular stromata are formed by confluence.

In the intercellular spaces under the epidermis the hyphae are colourless and closely packed, passing over into a darker coloured mass of round-polygonal cells 5–6.5 μ diam. This layer of dark, thick-walled cells is rather irregular and here and there penetrates deeper into the leaf tissues. From this stromatic base arise very numerous, closely crowded, cylindrical or club-shaped bodies 200–240 μ high, 96–160 μ thick at the base

160–200 μ thick at the apex. This column has the same prosenchymatous structure as the basal stroma. In the upper, broader part of each there is a single loculus sunk in the stroma; this has no separate wall, but is covered by the polygonal-celled hyphal web of the stroma column. The cells of this covering layer are larger than those at the base, being 7–10 μ diam.

In this species of *Rosenscheldia* the sterile part of the column is very short, the loculus being sub-spherical 100–150 μ diam. Asci basal, clustered, eight-spored, paraphysate, narrow-clavate, thickened round apex, 55–70 $\mu \times 16$ –17 μ . Paraphyses numerous, filiform. Spores distichous, four-celled, fuscous, clavate, 20–22 μ long, 6–6.5 μ thick at the broad end.

Rosenscheldia horridula Doidge, nov. sp.

Stromata plerumque hypophylla, minuta, usque 2 mm. diam., v. confluenta majores, columnata, columnis cylindraceutis v. clavatis, dense aggregatis, 200–240 μ altis, base 96–160 μ , apice 160–200 μ crassis. Loculi apicales singuli in quaque columna, stromate apicale tegentes, 100–150 μ diam. Asci paraphysati, octospori, clavati, ad apicem incrassati, 55–70 $\mu \times 16$ –17 μ ; paraphysibus numerosis, filiformis. Sporae distichae, 3-septatae, fuscae, clavatae, 20–22 $\mu \times 6$ –6.5 μ .

Hab. in foliis *Oleae capensis*, Durban, 4.6.12, leg. Pole Evans [5149].

22. *Auerswaldia examinans* (M. et B.) Sacc.

Syll. Fung. II, p. 626.

Syn. *Dothidea examinans* M. et B.—Bl. javan, 520.

Sphaeria examinans Mont. et Berk.—Hooker's Lond. Journ. Bot. I, p. 156.

On bark of old seedling citrus tree, Krantzklouf, Natal, 7.4.14, Bell [7733].

Auerswaldia examinans was originally described as occurring on the bark of a tree in Java. The South African specimen agrees with the description given in the *Annales Mycologici* (XIII, 1915, p. 298), except that the loculi are somewhat larger.

The bark is thickly beset with the erumpent, dull black, tuberculate stromata in all stages of growth, from the youngest which are punctiform to the mature columns with a length of 800 μ . The surface is rough, granular, covered with the somewhat mammillate protruding apices of the loculi.

The stroma develops under the bark, spreading considerably, and breaking through at several points develops the stromatal columns in which the loculi are formed. The hyphae forming the column are pendicular to the hypostroma, towards the apex becoming indistinctly cellular. In the young fruiting bodies only the hypostroma and the outer layer of the column is dark coloured. The loculi lie in the upper part of the column, and are 140–160 $\mu \times 120$ –150 μ , circular to ellipsoid, and sunken in the stroma, the latter being raised in a series of minute peaks over the apices of the loculi. The loculi are not ostiolate, dehiscing by the rupture of the outer stromatic wall. Asci basal, clavate-cylindrical, 65 $\mu \times 18$ μ , with a short, broad foot 12 μ long. Spores distichous, brown, one-celled, oblong-ellipsoid, rounded at both ends, 20–24 $\mu \times 10$ –11 μ .

23. *Dothidina disciformis* (Wint.) Theiss. and Syd.

Ann. Myc. XIII (1915), p. 304.

Syn. *Auerswaldia disciformis* Wint.—Hedwigia XXIII, 1884, p. 170; Syll. Fung. IX, p. 1033.

On leaves of *Myrica* sp., near Capetown, 1883, MacOwan (Rabh. Wint. Fung. Eur. 3063) [3403]; Krantzklouf, Natal, 14.8.14, V. d. Bijl [8396]; Uganda, March, 1916, Dummer [11986].

Stromata on both sides of the leaf, round or irregular, often angular, 2–5 mm. broad, forming black, comparatively smooth cushions, raised 200–300 μ above the leaf surface. The epidermis, which is filled with stroma, at first covers the true stroma, but later it becomes torn and pushed back, and is also torn away from the stroma, with which it was

at first united. It then surrounds the free erumpent fruiting bodies, being torn into broad black flaps. The inner part of the stroma is comparatively light coloured, but consists of thick-walled hyphae, 6–10 μ broad. Loculi crowded, spherical or spherical-ovate, 120–160 μ diam., not reaching the base of the stroma, without a true wall. Paraphyses filiform. Asci cylindrical, rounded above, narrower at the base, 100–135 $\mu \times$ 14–16 μ . Spores obliquely monostichous or almost distichous up to eight in an ascus, elliptic, one-celled, olive brown, 17–19 $\mu \times$ 8–9 μ .

24. *Parastigmatea* Doidge, nov. gen. *Stigmateacearum*.

Omnia ut in *Stigmatea*, sed sporae hyalinae, continuae.

Parastigmatea nervisita n. sp.

On leaves of *Stephania hernandifolia*, Maritzburg, Natal, 26.6.11, Doidge [1656]; 6.4.14, Doidge [8352].

Ascomata epiphyllous, minute, dull black, punctiform, circular, 240–300 μ diam., discrete, but in small groups on main and lateral veins, 80–90 μ high, smooth, remaining covered by the cuticle. Basal membrane brown, about 10 μ thick, composed of concentric layers of compressed hyphae. Covering membrane radial in structure, composed of sinuous hyphae, 3–3.5 μ thick, arched in centre, flat at edges, 6–7 μ thick, dehiscing by a circular central pore; central cavity containing asci about 150 μ diam. Asci paraphysate, ovate or broadly ellipsoid, thick walled, eight-spored, 50–53 $\mu \times$ 26.5–30 μ . Spores distichous or conglobate, hyaline, sub-clavate or somewhat piriform, rounded at both ends, continuous, 20–24 $\mu \times$ 8–10 μ .

Parastigmatea nervisita Doidge, nov. sp.

Ascomata epiphylla, minuta, atra, rotundata, in nervibus insidentia, 240–300 μ diam., 80–90 μ alta, laeves, cuticulare tecta, radiatim e hyphis undulatis 3–3.5 μ crassis, contexta. Asci paraphysati, ovati vel late ellipsoidei, 8-sporei, 50–53 $\mu \times$ 26.5–30 μ . Sporae distichae v. conglobatae, hyalinae, continuae, subclavatae v. sub-piriformae, utrinque rotundatae, 20–24 $\mu \times$ 8–10 μ .

Hab. in foliis *Stephaniae hernandifoliae*, Maritzburg, 26.6.11, et 6.4.14, leg. Doidge [1656] et [8352].

25. *Perischizon oleifolium* (Kalch and Cke.) Syd.

Ann. Myc. XII (1914), p. 265, XII (1915), p. 269.

Syn. *Dothidea oleifolia* Kalch et Cke., Grevillea IX (1880), p. 31.

On *Olea capensis*, 9.1.14, Kirstenbosch, C.P., Pearson [7374].

On *Olea laurifolia*, 21.6.16, Malalane, Eastern Transvaal, Hall [9749].

Stromata amphigenous, scattered or crowded, circular, raised, $\frac{1}{2}$ – $\frac{3}{4}$ mm, diam., with a central foot 100–200 μ diam. penetrating into leaf tissues. Surface rough, lumpy, with a single (occasionally double) peripheral wall, beset laterally as well as at the base with bristly, brown, truncate septate hyphae, 100 μ long, 5–8 μ thick, which later disappear. Stroma hard in texture, dark, prosenchymatous, composed of parallel hyphae. The loculi are sunken in the periphery and coalesce to form a circular hymenium, the stromatal cover becoming mucilaginous at maturity and disappearing. Asci clavate, thick-walled round the apex, 90–115 $\mu \times$ 28–38 μ , eight-spored, embedded in mucilage. Paraphyses simple, thick, clavate at the apex, and 3–4 μ broad, light brown. Spores distichous, ellipsoid-oblong, brown, 26–34 $\mu \times$ 11–14 μ , two-celled, constricted, abruptly rounded at both ends; upper cell somewhat broader than the lower.

26. *Phragmodothella nervisequens* n. sp.

On leaves of *Burchellia capensis*, Van Stadens Pass, 7.7.18, Pole Evans [11446].

Stromata amphigenous, but mostly hypophyllous, forming elongated, raised, rusty-brown streaks, up to 1 cm. long and 1 mm. broad, on mid-rib and main lateral veins; smaller, punctiform, elongated, or branched on smaller veins. Stromata developed under

the epidermis, causing a gall-like hypertrophy of the leaf tissues, later becoming erumpent; 120–160 μ high, the inner part of the stroma being composed of compact brown cells 3–5 μ diam., prosenchymatous between the loculi, and with a darker, roughened crust. Loculi numerous, immersed, sub-spherical, 80–100 μ diam. Asci aparaphysate, eight-spored, clavate or ellipsoid, thick-walled round apex, with a short foot, 45–60 $\mu \times$ 20–23.5 μ . Spores hyaline, distichous, 3-septate, oblong or somewhat clavate, slightly constricted at the centre, 18–20 $\mu \times$ 5–6.5 μ .

Phragmodothella nervisiquens Doidge, nov. sp.

Stromata amphigena, plerumque nervisequentes hypophylla, nervis majoribus usque 1 cm. longa et 1 mm. lata; nervis secundariis minores punctiforma, elongata v. ramosa; erumpentia dothideodea, 120–160 μ alta. Loculi numerosi, immersi, suborbiculares, 80–100 μ diam. Asci aparaphysati, octospori, clavati v. ellipsoidei, apice incrassati, breviter pedicellati, 45–60 $\mu \times$ 20–23.5 μ . Sporae hyalinae, distichae, triseptatae, oblongae v. subclavatae, medio leniter constrictae, 18–20 $\mu \times$ 5–6.5 μ .

Hab. in foliis *Burchelliae capensis*, Van Stadens Pass, 7.7.18, leg. Pole Evans [11446].



Fig. 5.—*Phragmodothis asperata*.

27. *Phragmodothis asperata* Syd.

Ann. Myc. XIII (1915), p. 345.

On the stem of *Euphorbia schintzii*, Wagenpadsnek, Pretoria Dist., 26.12.11, Pienaar [2159]; 31.12.13 [7353].

Stromata only growing in the outer layers of cells and becoming apparently quite superficial through the rupture of the periderm, almost spherical, 1½–3 mm. diam., 1½–2 mm. high, black, surface much wrinkled and cracked. Interior of stroma brown, the base and periphery being dark, opaque, formed of perpendicular, ascending hexagonal cells 15–19 μ diam. Loculi elliptical, 250 $\mu \times$ 200 μ , sunken under the periphery, often apparently free through the crumbling away of the stroma surface. Asci clavate, thick-walled, 140–200 $\mu \times$ 16–32 μ , 6–8 spored. Spores distichous or obliquely monostichous, oblong, four-celled, narrowed towards each end, but not pointed; straight, or one side straight and the other convex, verrucose at maturity. Paraphyses wanting.

28. *Trabutia evansii* Theiss. et Syd.

Ann. Myc. XIII (1915), p. 352.

On *Ficus* sp., Lourenço Marques, Portuguese East Africa, May, 1909, Howard [668].

Stromata only epiphyllous, on inconspicuous yellowish or brownish, discoloured spots, small, circular, 1 mm. diam., usually more or less regularly arranged in groups $\frac{1}{2}$ –1 cm. diam., but seldom becoming confluent. Not infrequently these groups are so numerous that they run together, and the whole leaf surface is more or less evenly and thickly beset with the small stromata. Stromata convex, almost hemispherical, unilocular, between the cuticle and the epidermis, composed of grey brown, vertical, parallel hyphae 4–5 μ diam. Clypeus opaque, 50–80 μ thick, loculus 300–450 μ diam., 150–240 μ high, with a wall 15–18 μ thick composed of delicate brown hyphae and concentric in structure. Asci cylindrical with monostichous spores, or clavate with distichous or somewhat conglobate spores (in the latter case the spores are often oblique or transverse), 70–90 $\mu \times$ 10–16 μ , eight-spored. Paraphyses numerous. Spores ellipsoid, abruptly rounded at both ends, continuous, hyaline, 11–14 $\mu \times$ 8–9 $\frac{1}{2}$ μ .

29. *Trabutia ficuum* (Niessl.) Theiss. et Syd.

Ann. Myc. XIII (1915), p. 352.

Syn. *Phyllachora ficuum* Niessl. Hedwigia XX. 1881, p. 99. Syll. Fung. II, p. 598.

On *Ficus* sp. Portuguese East Africa, 24.5.1908, Howard [520].

On *Ficus howardiana*, Lourenço Marques, P.E.A., 30.8.1909, Howard [719].

Stromata epiphyllous, shiny black, small, $\frac{1}{2}$ –1 mm. diam., raised, irregular, between the cuticle and the epidermis, becoming flat at the edge. In the centre there are a few loculi, 120–140 μ high, with opaque covering stromatal layer about 30 μ thick, the inner part of the stroma is lighter brown. Loculi flattened-spherical or lenticular, from 220 μ diam. and 100 μ high, to 320 μ diam. and 115 μ high. Asci paraphysate, cylindrical-clavate, 55–65 $\mu \times$ 14–18 μ , eight-spored. Spores hyaline, continuous, monodistichous, oval to ellipsoid, rounded at both ends, 11–13 $\mu \times$ 6 $\frac{1}{2}$ μ .

30. *Trabutia nervisequens* (Lingelsh) Theiss. et Syd.

Ann. Myc. XIII (1915), p. 353.

Syn. *Phyllachora schweinfurthii* P. Henn., var. *nervisequens* Lingelsh, Engl. Bot. Jahrb. XXXIX, 1907, p. 604.

On *Ficus* sp., Umgeni, near Durban, 21.3.10, Doidge [854]; Malvern, Natal, 28.12.11, Doidge [1994]; Winkle Spruit, Natal, 29.1.12, Pole Evans [2019]; Port Shepstone, Natal, 15.10.12, Pole Evans [5608]; Amanzimtoti, 14.6.14, Franks [7813].

This fungus was originally described on *Ficus hochstetteri*, var. *glabrior* in Abyssinia. It appears to occur quite commonly on the Natal coast.

Stromata epiphyllous, developed on inconspicuous yellowish coloured spots, chiefly along the mid-rib and lateral veins, being elongated in the direction of the vein and more or less confluent. Small stromata only 1 mm. long, larger ones up to 2 cm. long, very slightly shiny, between the cuticle and the epidermis, with one or many loculi, flat, somewhat wavy. The inner part of the stroma is prosenchymatous, composed of brown, parallel hyphae, perpendicular to the leaf surface, and 7–9 μ thick. Loculi lenticular, 400–600 μ diam., 250–300 μ thick. Locular wall comparatively stout, 12–18 μ thick, light brown, composed of numerous concentric layers of very delicate hyphae. Clypeus 40–60 μ thick, stout, opaque. Asci cylindrical, stipitate, paraphysate, 75–110 $\mu \times$ 11–14 μ . Spores usually monostichous, broadly ellipsoid, continuous, hyaline, 12–16 $\mu \times$ 8–10 μ .

On the surface of young stromata a conidial layer is often developed, which produces hyaline, filiform, crumpled conidia.

Var. *robusta* Doidge.

On *Ficus* sp., Port Elizabeth, 20.6.19, Gunn [12248].

Differs from the type in the thickness of the stroma and in the form and size of the loculi. Stroma 400–450 μ high, loculi sub-spherical, 330–370 μ diam., 300–400 μ high.

31. *Catacauma dalbergiicola* (P. Henn.) Th. et Syd.

Ann. Myc., XIII (1915), p. 388.

Syn. *Phyllachora dalbergiicola* P. Henn., Hedwigia XXXVI, 1897, p. 224; Syll. Fung. XIV, p. 664.

On leaves of *Dalbergia armata*, Barberton Dist., August, 1906, Pole Evans [588]; Stella Bush, Durban, 11.7.11, Doidge [1662]; Verulam, Natal, 3.7.13, Pole Evans [6808].

The type specimen is on leaves of *Dalbergia variabilis* and was collected in Brazil. The South African specimens agree exactly with the description of the original, except in the size of the loculi, which are somewhat smaller.

Epiphyllous, taking different forms on different leaves, sometimes very minute, sometimes producing round-angular stromata up to $1\frac{1}{2}$ –2 mm. diam., convex, somewhat shiny, between the epidermis and the palisade cells. Clypeus epidermal, opaque, about 35 μ thick, the inner part of the stroma being formed of vertical, parallel, light rusty-brown prosenchyma. Loculi sunken in the stroma, one or more according to the size of the stroma, 240–300 μ diam. and 180–200 μ high (in the original description 360–440 μ diam.), with a thin wall. Asci clavate, paraphysate, 60–70 $\mu \times 16$ –20 μ . Spores distichous, oblong-ellipsoid, hyaline, continuous, 12–14 $\mu \times 6$ –7 μ .

32. *Catacauma Pterocarp* Syd.

Ann. Myc. XIII (1915), p. 387.

Syn. *Phyllachora Pterocarp* Syd., Ann. Myc., 1912, p. 40.

On *Pterocarpus angolensis*, Letaba Drift, Zoutpansberg Dist., 6.8.11, Doidge [1807]; Barberton, 22.8.12, V. d. Bijl [5132].

Stromata epiphyllous, scattered, small, irregularly circular, 1–2 mm. diam., shiny black, on pale leaf spots, slightly convex, with somewhat undulating surface. Clypeus epidermal, opaque; loculi with their bases on the palisade tissue, 1–6 in each stroma, flattened-spherical, closely appressed to the leaf tissue at the base, with thin brown walls, 350–450 μ diam., 240–300 μ high. The upper parts of the loculi are connected by thick stromatic tissue. Asci paraphysate, clavate-cylindrical, 60–80 $\mu \times 16$ –24 μ . Spores monostichous or distichous, hyaline, one-celled, elliptic, rounded at both ends, 14–18 $\mu \times 8$ –9 μ .

The spores of both specimens appear to be somewhat immature. Theissen and Sydow (*loc. cit.*) state that it is doubtful whether there is a working difference between this fungus and *Phaeostroma pterocarp* Syd. with two-celled spores.

33. *Catacauma Peglerae* n. sp.

On leaves of *Eugenia capensis*, Kentani, 6.7.15, Pegler (Pegler No. 2340) [9099]; Umbogintwini, Natal, 9.5.13, Doidge [6636]; Scottsburgh, Natal, 5.7.13, Pole Evans [6841]; Warner Beach, Natal, 1.4.18, Bottomley [11667].

Stromata epiphyllous, 1–2 mm. diam., circular, often forming larger, irregular stromata by confluence. Central stroma often surrounded by a ring of secondary stromata at a distance of 2–5 mm. (radius), which may be small and discrete, or fuse to form a continuous ring. Stromata black, shining, somewhat conical or mammillate in section. Epidermal clypeus about 40 μ thick, black, opaque, the stroma lying in the cavity between the arched epidermal clypeus and the depressed and hollowed palisade tissue. Locular wall thin and very closely appressed to the palisade cells. Loculi flattened-hemispherical

or pear-shaped, 400–600 μ diam. and 350–500 μ high. The loculi are often connected by opaque black stromatic tissue about 150 μ thick. Asci paraphysate, eight-spored, clavate-cylindrical, 120–140 $\mu \times 17$ –20 μ , with a short foot, 10–13 μ long. Spores mono-distichous, one-celled, hyaline, oval to ellipsoid, 20–23 $\mu \times 12$ –13 μ .

Catacauma peglerae Doidge, nov. sp.

Stromata epiphylla, 1–2 mm. diam. v. conflundo majores, nonnunquam concentricae dispositae, atrae, nitidulae, subconicae, subepidermales. Loculi 400–600 μ diam., 350–500 μ alti, clypeo opaco, aterrimo, 40 μ crasso, parietibus tenuibus. Asci paraphysati, octospori, cylindraneo-clavati, 120–140 $\mu \times 17$ –20 μ , breviter pedicellati. Sporae monostichae v. distichae, continuae, hyalinae, ovatae v. ellipticae, 20–23 $\mu \times 12$ –13 μ .

Hab. in foliis *Eugeniae capensis*, Kentani, 6.7.15, leg. Pegler [9099].

34. *Catacauma grammicum* (P. Henn.), Theiss. et Syd.

Ann. Myc. XIII (1915), p. 382.

Syn. *Phyllachora grammica* P. Henn., in Flore du Bas et Moyen Congo in Ann.

Mus. du Congo, Vol. II, fasc. II, 1907, p. 98.

On leaves of *Ficus capensis*, Lemana, Zoutpansberg Dist., 14.8.11, Doidge [1829]; Kentani, C.P., 1.3.15, Pegler (Pegler No. 1993) [8884].

Stromata scattered, amphigenous, usually only on one side of the leaf, but occasionally an epiphyllous is found opposite to a hypophyllous stroma, following the nerves, forming dull black, linear streaks up to 7 mm. long and 0.8 mm. broad on a yellow brown leaf spot; from these short lateral streaks radiate or form thin connecting branches with neighbouring stromata. Stroma sub-epidermal, the base being sharply defined against the sub-epidermal cell-layer; the sub-epidermal layer and the rest of the mesophyll remaining unchanged even when two stromata are formed opposite to each other, on each of the leaf surfaces; only the region adjoining the base of the stroma is coloured red. The stroma consists of vertical, parallel prosenchyma, in the epidermal clypeus the cells are short and opaque black. Loculi broadly ellipsoid, raising the epidermis in which the clypeus has developed; at the base only resting on a thin line of stroma which is against the sub-epidermal cells, 300–350 μ diam., 140–170 μ high. Asci cylindrical, paraphysate, eight-spored, 45–58 $\mu \times 7$ –10 μ . Spores monostichous, one-celled, hyaline, ellipsoid, rounded at both ends, 8–10 $\mu \times 4$ –5½ μ .

35. *Phaeochorella parinari* (P. Henn.) Th. et Syd.

Ann. Myc. XIII (1915), p. 405.

Syn. *Cocconia parinari* P. Henn., Engl. Bot. Jahrb. XXX, 1901, p. 257; Syll.

Fung. XVIII, p. 159.

On leaves of *Parinarium capense*, Pretoria Dist., October, 1908, Doidge [613]; 10.4.11, Erasmus [1272]; 29.4.11, Doidge [1507]; 8.4.12, Doidge [2203]; Seven Oaks, Natal [10981].

On *Parinarium mobola*, Rhodesia, Howard [730]; Zoutpansberg Dist., 6.8.11, Doidge [1809]; Barberton, 16.8.11, V. d. Bijl [1922]; Zoutpansberg Dist., 19.10.14 [9173].

Stromata epiphyllous, more or less covering the leaf surface, round to irregular, 1½ to 4 mm. in diam., convex, somewhat shiny, black, with a rough surface, 400–450 μ high, with a flat base resting on the sub-epidermal layer of cells, which is not stromatized, covered with an epidermal clypeus, light brown, vertically prosenchymatous in structure. Loculi numerous, pear-shaped, 300–400 μ diam., with an indistinctly periphysate neck. Asci cylindrical, 85–100 $\mu \times 11$ –15 μ , with delicate, filiform, hyaline paraphyses. Spores up to eight in an ascus, mostly monostichous, elliptic to oblong, rounded at both ends, red-brown, one-celled, with a slender medial hyaline band, 12–17 $\mu \times 8$ –9 μ .

36. *Phragmocaula viventis* (Cke.) Theiss. et Syd.

Ann. Myc. XIII (1915), p. 41.

Syn. *Dothidea viventis* Cke., *Grevillea* V, p. 16.

Phyllachora viventis (Cke.) Sacc., Syll. Fung. II, p. 601.

Dothidea viventis var. *albizziae* Cke., *ibid*.

Phyllachora albizziae Cke., *Grevillea* XIII, p. 65.

Homostegia albizziae (Cke.) Berl et Vogl., Syll. Fung. IX, p. 1049.

On leaves of *Albizzia fastigiata*, Inanda, Natal, May, 1881, Medley Wood (Wood No. 583) [9492] and [16445]; Winklespruit, Natal, 13.4.11 and 2.7.11, Pole Evans [1401] and [1583]; Stella Bush, Durban, 11.7.11, Doigde [1615]; Verulam, Natal, 3.7.13, Pole Evans [6829].

Stromata usually hypophyllous, small, irregular elliptic or oval in outline, somewhat raised, dull black, 0.5 to 0.7 mm. in length, the long axis often lying along a vein or in small groups of 2-4. Stroma arising between the peridermal and the sub-epidermal cells, with its base at the level of the epidermis. Loculi, several, under a common, arched, epidermal clypeus, spherical to lenticular, 190-200 μ diam. at maturity. Asci paraphysate, cylindrical, 80 $\mu \times$ 10-12 μ , with a foot 15-18 μ long. Spores distichous, colourless, four-celled, constricted at the middle, 20-24 $\mu \times$ 5-6 μ . The asci do not stain blue with iodine.

37. *Scolecodothis capensis* n. sp.

On leaves of *Olea* (?) *foveolata*, Howiesons Poort, Grahamstown, 17.1.17, Doigde [10963]; 12.7.19, Doigde [12379]; Van Stadens Pass, 13.11.17, Doigde [10869].

Stromata amphigenous; the epiphyllous stromata develop first, they are scattered, round to irregular, 1-1½ mm. diam., raised and convex above leaf surface, occasionally larger and irregular in shape by confluence, black and shining. Hypophyllous stromata smaller and flatter, and usually more or less undeveloped and sterile. The leaf tissue is not discoloured. Stroma between the epidermis and the palisade tissue. Clypeus epidermal, thick, opaque, 90-100 μ thick. The inner part of the stroma consisting of lighter brown, prosenchymatous plectenchyma. Loculi 4-5 in each stroma, immersed, lenticular or somewhat irregular by compression, 550-650 μ diam., 200-250 μ high. Asci paraphysate, eight-spored, narrow, ellipsoid, tapering to both ends, pedicellate, 100-120 $\mu \times$ 13-14 μ . Spores parallel, hyaline, one-celled, narrow, fusiform, pointed at both ends, 60-70 $\mu \times$ 5-6 μ in centre.

Scolecodothis capensis Doigde, nov. sp.

Stromata amphigena, sparsa, rotundata v. irregulares, 1-1½ mm. diam., convexa, atra, nitidula. Loculi 4-5 in quoque stromate, 550-650 μ diam., 200-250 μ alti; clypeo opaco, aterrimo, 90-100 μ crasso. Asci paraphysati, octosporo, anguste ellipsoidei, utrinque attenuati, pedicellati, 100-120 $\mu \times$ 13-14 μ . Sporae parallelae, hyalinae, continuatae, fusoidae, 60-70 $\mu \times$ 5-6 μ .

Hab. in foliis *Oleae* (?) *foveolatae*, Grahamstown, 17.1.17, leg. Doigde [10963].

38. *Phyllachora caffra* Syd.

Ann. Myc. XV (1917), p. 548.

Syn. *Physalospora caffra* Syd., Ann. Myc. X (1912), p. 39.

On leaves of *Cordia caffra*, Amanzimtoti, Natal, 10.7.11, Doigde [1631].

Stromata epiphyllous, minute, gregarious on irregular rusty-brown spots, 250-300 μ in diam., unilocular; on the upper leaf surface round, punctiform, convex, shining; on the lower surface the epidermis becomes somewhat convex, and later the stroma becomes visible as a minute, dull-black point. Loculi occupying the whole thickness of the leaf, spherical, 250-300 μ diam., with a short epidermal clypeus at the apex, the lateral and basal wall consisting of a web of hyaline or yellowish threads; where the basal wall is

near the lower epidermis, a short clypeus is formed. The leaf, which is normally 200–220 μ thick, becomes arched over the loculus to a thickness of 360 μ . Asci cylindrical, paraphysate, 75–90 $\mu \times$ 9–12 μ . Spores monostichous, ellipsoid, hyaline, broadly rounded, one-celled, 11–13 $\mu \times$ 6–8 μ .

39. *Phyllachora dombeyae* Syd.

Ann. Myc. XV (1917), p. 532.

Syn. *Physalospora dombeyae* Syd., Ann. Myc. X (1912), p. 441.

On *Dombeya rotundifolia*, Equeefa, Natal, 24.4.11, Fuller [1539].

On *Dombeya schimperiana*, Woodbush, Zoutpansberg Dist., 4.8.11, Doidge [1762].

Stromata epiphyllous, gregarious on yellow leaf spots, $\frac{1}{3}$ – $\frac{1}{2}$ mm. diam., convex, somewhat shiny, not visible on the lower leaf surface, or only visible as minute dull black spots; with one or few loculi. Loculi spherical, 160–190 μ diam., covered at the apex with a dark epidermal clypeus, almost without lateral walls, not touching the lower epidermis, or forming a very short clypeus in the lower epidermal cells. Asci clavate, paraphysate 60–75 $\mu \times$ 15–20 μ . Spores distichous or tristichous, one-celled, hyaline, rounded at both ends, asymmetrical, straight or slightly curved, 25–35 $\mu \times$ 3–4 $\frac{1}{2}$ μ . Conidia filiform, curved 18–26 $\mu \times$ 1–1 $\frac{1}{2}$ μ .

40. *Phyllachora melianthi* (Thuem.) Sacc.

Syll. Fung. IX, p. 1013; Ann. Myc. XIII (1915), p. 528.

Syn. *Rhytisma melianthi* Thuem., Flora, 1876, p. 569.

Cryptomyces melianthi (Thuem.), Sacc., Syll. Fung. VIII, p. 707.

On *Melianthus major*, near Capetown, January, 1884, MacOwan (Rabh. Wint. Fung. Eur. 3557) [3897].

On *Bersama lucens*, Kentani, 3.7.15, Pegler [9017].

Stromata on both sides of the leaves, numerous, scattered, small, unilocular, hemispherically or conically convex, flat at the periphery, $\frac{1}{2}$ mm. diam., black; on the opposite side of the leaf the grey epidermis bulges out, and later becomes filled with stroma and becomes black. Between the two leaf surfaces lies a single large stroma (1–1 $\frac{1}{2}$ mm. diam.), which either remains unilocular, surrounded only by a broad, flat, stromatic border, or contains several loculi, and is then not surrounded by sterile stroma. Clypeus epidermal opaque, 15–25 μ thick, inner part of stroma and locular wall consisting of light brown prosenchyma. Locular wall thin, delicate. Loculi flattened spherical, 240–280 μ diam. (in stromata on *Bersama*, 320–450 μ diam.), 240–300 μ high, occupy from three-quarters to the whole of the thickness of the leaf; occasionally irregular in shape where a vascular bundle impinges on locular wall, dehiscing towards the upper leaf surface by an apical pore. Asci paraphysate, cylindrical to clavate, 60–70 $\mu \times$ 15–18 μ . Spores mono- or distichous, ellipsoid, hyaline, one-celled, rounded at both ends, 13–14 $\mu \times$ 7–8 μ .

41. *Phyllachora hieronymi* P. Henn.

Pilze Ostaf. apud. A. Engler, Die Pflanzenwelt-Ostafrikas, p. 34; Syll. Fung. XIV, p. 673.

On *Cyathea dregei*, Winters Kloof, Natal, 27.6.11, Doidge [1666].

Stromata epiphyllous, shiny black, often on the veins, slightly convex, 0.4–0.5 mm. diam., usually unilocular, penetrating right through the leaf. Clypeus epidermal, well developed, opaque, 30–33 μ thick. Locular wall brown, thin, 12–14 μ thick, often thinner at the sides. Loculus flat ellipsoid, 300–320 μ diam., 170 μ high. Asci paraphysate, clavate-cylindrical, briefly pedicellate, 65–70 $\mu \times$ 16–20 μ . Spores distichous, hyaline, oblong, 20–24 $\mu \times$ 7–8 μ .

42. *Phyllachora aberiae* P. Henn.

Engl. Bot. Jahrb., XLI (1908), p. 272; Syll. Fung. XXII, p. 411.

On *Doryalis caffra*, Driefontein, Zoutpansberg Dist., 12.8.11, Doidge [1813]; Uitenhage, C.P., 31.5.12, Pienaar [2420]; 17.10.06 [224].

Stromata epiphyllous, cellular, $\frac{3}{4}$ –1 mm. broad, shiny, black, subrotund to irregular, slightly raised, showing on the under side of the leaf as minute raised points on the discoloured leaf tissue. Leaf normally 280 μ thick, hypertrophied in the region of the stroma, 400–500 μ thick. Clypeus epidermal epiphyllous, continuous, 25–34 μ thick. Loculi under the clypeus in the mesophyll, flask-shaped, or spherical with a wide neck, the base of the loculi being one-half or two-thirds of the distance through the thickness of the leaf; surrounded by a brown, thick wall, which is 25–38 μ thick and formed of loosely interwoven hyphae; usually opening towards upper leaf surface, occasionally towards the lower, when it is covered by a short hypophyllous epidermal clypeus: occasionally two loculi on opposite sides of the leaf come into contact at their bases and are consequently flattened. Loculi 240–270 μ diam., or 250–280 μ high with a breadth of 200–220 μ . Between the necks of neighbouring loculi a loose hyphal stroma spreads inwards from the clypeus.

Asci ellipsoid, with monostichous, transverse, or partly distichous spores; seldom cylindrical with distichous or imbricate monostichous spores, shortly pedicellate, foot usually 8–10 μ long, rarely 22–30 μ long; asci 80–90 μ long and 12–20 μ broad, according to the arrangement of the spores, eight-spored, paraphysate. Paraphyses numerous. Spores cylindrical, rounded at both ends, 26–33 $\mu \times 4\frac{1}{2}$ –5 μ , hyaline, one-celled.

43. *Phyllachora amaniensis* P. Henn.

Engl. Bot. Jahrb. XXXVIII, 1905, p. 113; Syll. Fung. XXII, p. 420; Ann.

Myc. XIII (1915), p. 473.

On leaves of *Ficus capensis*, Amanzimtoti, Natal, 19.3.14, Franks [7812]; Barberton Dist., 16.10.13, V. d. Bijl [7378]; Wyebank, Natal, 29.8.15, Doidge [9537].

This fungus was originally described on leaves of *Ficus* sp. collected at Amani, Usambara; the South African specimens agree with the description in all important particulars, but the spores are slightly narrower, the spores of the type being 7–8 μ wide.

Stromata epiphyllous, scattered or loosely arranged in groups $\frac{1}{2}$ –2 cm. diam., round or irregular, slightly raised, 1–1.5 mm. diam., dull black, not visible on upper surface, or showing as minute black pin-spots. Stroma originating in the lower part of the mesophyll, dividing the leaf into two very unequal portions, the lower narrower portion being pushed up by the developing stroma and forming an arch over it. Clypeus epidermal opaque, 25–35 μ thick. Loculi sub-spherical, less frequently flask-shaped, the narrowing in the latter case being due to crowding, deeply sunken in the leaf tissue, and occupying three-quarters or even more of the mesophyll. The base of the loculi usually rests on the palisade cells and the latter become filled with black, opaque stromatic tissue, which may or may not involve the upper epidermis. Loculi 240–400 μ diam., lateral walls delicate, 10–15 μ thick, consisting of light brown, rather thin-walled prosenchyma. Asci cylindrical with numerous paraphyses, 90–130 $\mu \times 9$ –11 μ , straight or curved, eight-spored. Spores monostichous, ellipsoid, rounded at both ends, hyaline, continuous, 14–17 $\mu \times 6$ –7 μ .

44. *Phyllachora Peltophori* Syd.

Ann. Myc. X (1912), p. 40; XIII (1915), p. 507.

On *Peltophorum africanum*, Ledzee, Zoutpansberg, 7.8.11, Doidge [1810].

Stromata on both surfaces of the pinnules, scattered or in groups, minute, about $\frac{1}{2}$ mm. diam., slightly convex, unilocular, smooth, shining. Loculi flattened, spherical, 200–250 μ diam., 160–180 μ high, covered above and below by the epidermal clypeus, which only extends a short distance beyond the loculi. Lateral locular wall weak or wanting. Asci clavate, paraphysate, 65–80 $\mu \times 14$ –18 μ . Spores distichous, ellipsoid, hyaline, rounded, 13–17 $\mu \times 6$ –8 μ .

45. *Phyllachora Lessertiae* n. sp.

On leaves of *Lessertia tenuifolia*, Smits Kraal, Boshof Dist., June, 1911, Burt-Davy [1568].

Stromata amphigenous and petiolar, but mostly epiphyllous, scattered or crowded, minute, up to 0.5 mm. diam., somewhat raised, dull black, mostly unilocular. Stroma reduced to a clypeus above each loculus, 100–120 μ broad, and certain hyphal strands and knots under the clypeus and extending beyond it in the sub-epidermal cells. Loculi sub-spherical to pyriform, 169–190 μ diam., 200–240 μ high, occupying about one-third of thickness of leaf. Locular wall about 15 μ thick, consisting of delicate, compressed, colourless hyphae, broader near apex and darker coloured, fusing above with clypeus. Asci eight-spored, paraphysate, cylindrical or narrow-ellipsoid, with a very short foot, 80–90 $\mu \times 12$ –16.5 μ . Periphyses present. Spores one-celled, hyaline, ellipsoid, rather thick-walled, obliquely monostichous or sub-distichous, 15–17 $\mu \times 8$ –10 μ .

Phyllachora Lessertiae Doidge, nov. sp.

Stromata plerumque epiphylla, sparsa, minuta, usque .5 mm. diam., vix vel leniter convexa, atra, clypeo epidermale, 100–120 μ lato, unilocularia. Loculi subglobosi v. pyriformi, 160–190 μ diam., 200–240 μ alti, parietibus circ. 15 μ crassis. Asci octospori, paraphysati, cylindracei, v. ellipsoidei, brevissime pedicellati, 80–90 $\mu \times 12$ –16.5 μ . Sporae continuae, hyalinae, ellipsoideae, utrinque rotundatae, 15–17 $\mu \times 8$ –10 μ .

46. *Phyllachorella rikatiensis* n. sp.

On leaves of *Andradia arborea*, Rikatli, P.E.A., September, 1918, Junod [11736].

Stromata minute, punctiform, about 0.3–0.5 mm. diam., amphigenous, penetrating right through the leaf, scattered over greater part of leaf surface or in round irregular patches about 5 mm. diam., black shining, slightly convex. Epidermal clypeus on both sides of leaf, black, opaque, about 20 μ thick, only extending a small distance on each side of the loculi. Inner part of stroma of usual prosenchymatous structure; in this the loculi are embedded. Stroma may contain a single flattened spherical loculus, 240–250 μ diam., 140–150 μ high, or a single pycnidium of similar dimensions. Very frequently the flattened spherical loculus is about 300 μ diam. and has a small lenticular pycnidium cut off from one surface; exceptionally, there are two smaller loculi and a pycnidium, the shape of each being an irregular cone on a curved base. Locular wall about 10 μ thick.

Asci aparaphysate, clavate, eight-spored, 50–60 $\mu \times 13$ –15 μ . Spores usually distichous, crowded in upper half of ascus, hyaline, one-celled, oblong, rounded at both ends, 10–13.5 $\mu \times 3$ –5 μ .

Conidia brown, oblong, continuous, 13.5–15 $\mu \times 6$ –7 μ .

Phyllachorella rikatiensis Doidge, nov. sp.

Stromata amphigena, minuta, circ. 0.3–0.5 mm. diam., atra, nitidula, leniter convexa. Loculi 1–3 in quoque stromate, sub-globosi v. compressi, 240–300 μ diam., 140–150 μ alti, clypeo atro, opaco, 20 μ crasso, parietibus 10 μ crassis. Asci aparaphysati, clavati, octospori, 50–60 $\mu \times 13$ –15 μ . Sporae plerumque distichae, hyalinae, continuae, oblongae, utrinque rotundatae, 10–13.5 $\mu \times 3$ –5 μ . Pycnidia loculis ascogenis similes, conidiis brunneis, oblongis, continuis 13.5–15 $\mu \times 6$ –7 μ .

Hab. in foliis *Andradiae arboreae*, Rikatli, P.E.A., 1918, leg. Junod [11726].

47. *Endodothella natalensis* n. sp.

On leaves of *Dalbergia armata*, Winklespruit, Natal, 6.7.12, Doidge [2513]; Verulam, Natal, 3.7.13, Pole Evans [6809].

Stromata epiphyllous, minute, punctiform about 0.3 mm. diam. or becoming larger by confluence, somewhat convex, black shiny, showing on the under-surface only as minute, brown blisters, uni- or bilocular. Clypeus epidermal, dark brown, sub-opaque, ca. 15 μ thick, spreading to a diameter of about 50 μ . Loculi sub-spherical to lenticular, 240–320 μ

diam. and 190-240 μ high, the base of the loculi being 30-45 μ from the lower surface of a leaf which is normally about 130 μ thick. Locular wall opaque, black, 13-16 μ thick, or somewhat lighter in colour and evidently prosenchymatous in structure. Loculi dehiscing by a apical pore opening towards upper side of leaf. In the bilocular stromata the space between the apices of the loculi is entirely filled with opaque, black, stromatal tissue. Asci paraphysate, eight-spored, ellipsoid or narrow cylindrical, straight or curved, 80-87 $\mu \times 15-17 \mu$, with a short foot 6-7 μ long. Spores distichous or obliquely monostichous, hyaline, two-celled, not constricted, fusiform, 20-23.5 $\mu \times 8.5-10 \mu$, wall about 1.5-2 μ thick, cells sub-equal, or upper cell 10 μ long, lower cell 13.5 μ long.

Endodothella natalensis Doidge, nov. sp.

Stromata epiphylla, minuta, rotundata, cir. 3 mm. diam. v. confluendi majores, leniter convexa, atra, nitidula. Loculi 1-2 in quoque stromati, sub-globosi v. globosi-depressi, 240-320 μ diam., 190-240 μ alti, totam fere folii crassitudinem occupantes, clypea, atro-brunneo, subopaco, ca. 15 μ crasso, 500 μ lato, parietibus opacis, atris, 13-16 μ crassis. Asci paraphysati, octospori, ellipsoidei v. cylindracei, recti v. curvati, 80-87 $\mu \times 15-17 \mu$, pede breve 6-7 μ long. Sporae distichae v. oblique monostichae, hyaline, 1-septatae, haud constrictae, fusiformae, 20-23.5 $\mu \times 8.5-10 \mu$, cellulis subaequalibus, v. supero 10 μ longo, infero 13.5 μ longo.

Hab. in foliis *Dalbergiae armatae*, Winklespruit, Natal, 6.7.12, leg. Doidge [2513].

48. *Endodothella strelitziae* (Cke.) Theiss. et Syd.

Ann. Myc. XIII (1915), p. 587.

Syn. *Dothidea strelitziae* Cke., Grevillea X., p. 120.

Phyllachora strelitziae Sacc., Syll. Fung. II., p. 606.

On leaves of *Strelitzia augusta*, Inanda, Natal, May, 1881, Medley Wood (Wood No. 580) [9468] and [10442]; Isipingo, Natal, 13.5.13, Doidge [6639]; Scottsburg, Natal, 5.7.13, Pole Evans [6831]; Inanda, Natal, 13.9.13, V. d. Bijl [6958]; Kentani, 8.1.16, Pegler (Pegler No. 2384) [9422].

Stromata round, conical to hemispherical, 0.7-0.85 mm. diam., in groups of two to three, dull black, epiphyllous, unilocular, surrounded by a red brown zone in the leaf tissue; on the under side of the leaf only a wine-red discoloration is visible. The stroma fills the epidermis and the two sub-epidermal layers of elongated, flat cells, and raises these up to form an opaque clypeus, tearing them away from the adjacent layers of parenchyma. In the cavity thus formed the loculi are produced; they are covered by a clypeus formed from three layers of cells and about 100 μ thick: they are 650 μ diam. and 180 μ high; being separated at the base from the leaf parenchyma by a thin brown stromatic layer, the hyphae penetrating with the parenchyma in a more or less compact mass. Asci clavate, briefly pedicellate, paraphysate, 55-75 $\mu \times 13-16 \mu$, eight-spored. Spores distichous, oblong or clavate, hyaline, septate somewhat above the middle, not or slightly constricted, 14-17 $\mu \times 3-4 \mu$, upper cell shorter but broader, rounded, lower cell tapering somewhat.

49. *Oligostroma maculiformis* (Wint.) Doidge.

Syn. *Didymella maculiformis* Wint., Rab. Wint. Fung. Eur. 3056 (1884).

Oligostroma proteae Syd., Ann. Myc. XII (1914), p. 265; XIII (1915), p. 592.

On *Protea grandiflora*, near Capetown, June, 1884, MacOwan (Rab. Wint. Fung. Eur. 3056) [3396].

On *Protea flanaganii*, Kentani, 17.7.12, Pegler, 5163. (This is the type collection of *Oligostroma proteae* Syd.)

On *Protea neriifolia*, Bains Kloof, near Wellington, 19.11.10, Doidge [1026]; 5.4.12, Stoneman [2231].

On *Protea* spp., Wellington, 10.6.11, Mally [1589]; without locality [943].

(?) Stroma not fully developed on *Protea* sp., Kentani, 19.10.12, Pegler [5618], and on *Protea abyssinica*, Diepkloof, near Dullstroom, 19.9.10, Doidge [931].

This fungus is very variable in its external appearance, but after careful comparison I think there can be no doubt that *Oligostroma protea* Syd. on *Protea flanaganii* [5163] is identical with Winter's *Didymella maculiformis*. The variations mentioned in macroscopic characters are due to variations in the thickness of the cuticle of the host, rather than to any variations in the structure of the fungus.

Stromata amphigenous, forming discoloured, roughened areas on both sides of the leaf; these are usually irregular in outline and up to 10 mm. in diam. or larger by confluence; less frequently they are more or less circular and develop centrifugally. The stroma is reduced to an epidermal clypeus, 150–250 μ diam. over each loculus and the short scattered hyphal strands or knots under it. Sometimes the loculi are closely crowded and the clypeus over adjoining loculi becomes confluent, forming a more or less continuous stromal plate, or the loculi may be more scattered, solitary, or in twos and threes.

Loculi immersed, globose or ovate globose, 100–150 μ diam., locular wall consisting of small brown cells and fused at the apex with the epidermal clypeus. Asci sessile, cylindrical-clavate or clavate, 75–100 $\mu \times$ 16–20 μ , rounded at the apex, paraphysate, eight-spored. Spores distichous, oblong-cuneate, unequally uniseptate, not constricted, 24–28 μ long; upper loculus shorter, but more broadly rounded or ovate 8–10 μ long, 7–9 μ broad, lower loculus longer and narrower, 15–19 $\mu \times$ 6–8 μ , hyaline or sub-hyaline.

50. *Ophiodothella edax* (B. and Br.) V. Höhn.

Fragm. XII No. 630 (1910).

Syn. *Dothidea edax* B. et Br., Journ. Linn. Soc., 1873, p. 135.

Ophiodothis edax Sacc., Syll. Fung. II, p. 653.

On leaves of *Tephrosia elongata*, Olifantsfontein, Pretoria Dist., 21.2.20, Pienaar [12822].

This fungus was originally described on leaves of *Tephrosia suberosa* from Ceylon. The South African specimen agrees with the description of the original, except in the size of spores. These are said to average 33–35 $\mu \times$ 2.8–3.2 μ , but it is also stated that the asci are not quite ripe. It is probable, therefore, that the difference in the measurements of the spores is due to the fact that the present specimen is in a more mature condition.

Stromata minute, black, punctiform, formed in groups on large yellow leaf spots. There is an epidermal clypeus on both upper and under surfaces of the leaf, which is black, opaque, about 20 μ thick in the lower epidermis, 25–30 μ in the upper. There are 1–3 loculi in each stroma; these are flattened-spherical, 200–250 μ diam., 150–200 μ high, at base and apex united with the clypeus. The locular wall is 8–10 μ thick and consists of hyaline very much flattened cells. Asci numerous, paraphysate, eight-spored, thin-walled, pedicellate, clavate-ellipsoid, 70–80 $\mu \times$ 10–13 μ . Spores parallel, filiform, continuous, hyaline, straight or curved, thin-walled, cylindrical, somewhat attenuated towards the blunt ends, 45–60 $\mu \times$ 3–3.3 μ .

THE GENUS BERSAMA.

By E. P. PHILLIPS, M.A., D.Sc., F.L.S., Botanist in Charge of the
National Herbarium.

THE genus *Bersama* was monographed by Mr. E. G. Baker in 1907 (Journ. Bot., XLV, p. 12), and two species, viz., *B. lucens*, Szyszl., and *B. tysoniana*, Oliv., were described from South Africa. From information received from the Union Forest Department, it was suspected that other species occurred in the South African forests, and, at the suggestion of the Chief of the Division of Botany, I undertook the examination of all the material in the local herbaria. This has led me to separate two plants as distinct species from specimens hitherto always called *B. tysoniana*.

The interest in the various species was first aroused by Mr. C. C. Robertson, M.F., the Senior Research Forest Officer, who sent samples of the bark of *B. tysoniana* to the Imperial Institute for examination, as it was reported that the natives used the bark medicinally. The information supplied by Mr. Robertson to the Imperial Institute did not tally in some respects with the actual samples forwarded, and he took a good deal of trouble to clear up the matter. We are now in a position to state that the information Mr. Robertson received applied to a distinct species which I have named *B. Swinnyi*, while the bark supplied was that of *B. tysoniana*. Dr. T. R. Sim,* in his description of *B. tysoniana*, falls into the same error, as the tree referred to by Mr. Henkel as having "a very bitter bark, like quinine in taste," is not this species, but quite a distinct plant, which I have named *B. Stayneri*, so his remarks on *B. tysoniana* do not refer to this species alone. Sim suspected the Port St. Johns' plants to be an allied but distinct species, and I have been able to confirm this. It is the plant now named *B. Swinnyi*.

Mr. J. J. Kotze, B.Sc., of the Forest Department, who has given me every assistance in this work, allowed me to go through the files relating to the genus, and, having the actual specimens in the Forestry Herbarium, I was able to clear up some doubtful points. All the correspondence referred to plants which went under the name of *B. tysoniana* and the information was confusing until I was able to definitely recognize that three distinct species were involved. The information derived from the above source I have tabulated under the specific names.

B. tysoniana.

The Conservator of Forests, Transkeian Conservancy, states: "Trees are scattered singly, fairly plentifully over most of the mountain forests and where accessible. It is a prolific seeder, and I am of opinion that it could be easily propagated artificially under mountain forest conditions in its habitat. In collecting the bark, natives do not cut down the whole tree, but merely take their requirements from time to time from living trees. I can confirm the statement: it is a well-known fact that local natives use the bark medicinally." (Stated that native doctors use the bark as a specific against fevers.—(E. P. P.)

* Sim, "Forest Flora, Cape Colony," p. 175.

A sample of bark was collected by Forester Van de Vijver from the Ngadu Patrol, and he writes: "This bark is used as medicine by some natives, but not all. In cases of calves being thin, they dose the animal with it powdered and mixed with other roots, also lambs." Forester Dawson submitted a sample of bark from the Amanzanyama Patrol, with the note: "Bark has a slightly bitter taste. I have only seen one tree in my patrol and very few natives know it here. They do not seem to use it themselves, but on their cattle for gall-sickness after boiling bark and dosing from one to two bottles." Forester Fegen collected samples of bark from the Notinsela Patrol, and reported: "Bark has a slightly bitter taste. Natives state that the bark is used as a medicine for hysteria, which is common amongst the native girls. Other natives state that it is used indiscriminately for both cattle and human beings. Judging by the number of trees barked, it must be used more extensively for medicine than natives will admit."

Forester Kriel, who collected the species in the Wilo Forest, Mqanduli District, reports it to be a tree about 25 ft. high with a girth of 42 in.

The tree is locally known as "Bitter Bark" and "Bastard Sneezewood" and by the natives as "Endiaza" or "Andianda."

B. Swinnyi.

Forester Swinny reports this species to be a tree up to 90 ft. high, with a stem diameter of 2-3 ft., and not common at Port St. Johns. He further states that there appears to be two forms of the species which the natives distinguish, though they have only one name, "Sendiandia." The smaller form, when full grown, has a trunk 1-1½ ft. in diameter.

Mr. Robertson first detected the difference in the barks of this species and *B. tysoniana*, and noted that the bark had "a strong bitter taste producing a sort of burning sensation in the mouth lasting for several hours, during which I could taste nothing else distinctly." He forwarded a sample of the bark to the Imperial Institute in March, 1915. Mr. Ross, then Conservator in the Eastern Conservancy, writes:* "I distinctly remember this tree being pointed out to me many years ago when stationed at Kokstad as being of value medicinally, an infusion of the bark being extensively used by the natives for certain ailments. It is found somewhat sparingly throughout the mid-Pondoland forests and close to the Natal border in the Mount Ayliff District. It is a very conspicuous object in the forests, of large girth, with deeply furrowed bark, and with foliage not unlike sneezewood. It is usually unsound near the base and shows signs of much damage by removal of bark. This tree is by some foresters confused with 'Coffee Pear' (*Pleurostylia capensis*), but a keen observer will easily distinguish a great difference."

Forester Pretorius collected samples of bark from the Cwebe Patrol and supplied the following information:—"Bark has a slightly bitter taste (Mr. Robertson remarks 'but burning taste'). The bark of this tree in these parts is very thin and can only be removed in small bits. The natives use it for different medicines, but they refuse to say what it is really used for." Specimens of bark from the Port St. Johns Patrol were received with similar information to the above; in each case Mr. Robertson noted the burning taste, more so in some specimens than in others.

B. Stayneri.

This species was sent to Mr. Robertson by Forester C. W. Chilvers in June, 1915. Mr. Robertson noted the differences between these specimens and *B. tysoniana* as follows:—

1. Bark has not got the "strong bitter taste" which I referred to in the case of one of the specimens from the Transkei (i.e. *B. Swynni* E.P.P.).
2. Under sides of the leaves and the petioles are very hairy, whereas they are glabrous on specimens received from the Transkei.
3. The specimens were in flower when collected in May. Sim states that this species † flowers in August to September.

* It is just possible that some of Mr. Ross's remarks may apply to *B. tysoniana*.

† Mr. Robertson, although he noted these differences, still considered the plant *B. tysoniana*.

Mr. C. F. Stayner, in a minute to the Conservator of Forests, Pietermaritzburg, states : " One large tree isolated by heavy fellings was in flower as late as September, and I have come to the conclusion that this species (but he confused his specimen with *B. tysoniana* E.P.P.) is inconsistent in its period of blossoming. I find that *Bersama* is well known to native herbalists and is frequently employed, but for what disease it is a specific I cannot ascertain." Mr. Stayner submitted his specimen to the Bolus Herbarium, and these were reported on as follows:—" Very nearly allied to *B. tysoniana*, Oliv., and probably that species, but the type of *B. tysoniana* has more slender, longer, and less tomentose petioles."

Mr. J. S. Henkel, who also collected the species at Mkazeni Forest Reserve, Riverside, Natal, in March, 1917, states " that the tree is about 30 ft. high—a bole about 70 in. in girth and 12 ft. long, somewhat crooked. Natives use the bark as medicine and the tree has been injured by the removal of bark. Bark has a bitter quinine-like taste." Mr. Henkel also noted that the leaves were hairy, and not glabrous. Forester Honshold informs Mr. Henkel that in the reserve there are about three to four other trees, all smaller than the one he saw. In the bush many seedlings were noticed up to about 4 ft. high. The young shoots are frequently reddish in colour and leaves serrated and mucronulate, but older leaves have margins entire.

Mr. Robertson took considerable pains to sift all the evidence he had accumulated, and tabulated his results* as follows:—

Mr. Robertson distinguished the various samples of bark he examined as (1) those with a bitter taste and (2) those with a bitter and burning taste. This latter property he referred to as "X."

Forest.	Large or Small Variety.	Sample of Bark.	Taste.	Remarks.
MOUNTAIN FORESTS.				
Kambi.....	Presumably the large variety	Sent June, 1914, and February, 1915	Not X.....	<i>B. tysoniana</i> , sent to Imperial Institute.
Nqadu.....	Large.....	Collected March, 1915..	Not X.....	<i>B. tysoniana</i> .†
Amanzimnyama.	Large.....	Collected March, 1915..	Not X.....	<i>B. tysoniana</i> .
COAST FORESTS.				
Port St. Johns..	Large.....	Sent July, 1914.....	Has X.....	<i>B. Swinnyi</i> .
Bulolo Forest..	Large.....	Collected March, 1915..	Has X, though not strongly	<i>B. Swinnyi</i> .
Mboleni Forest..	Large.....	Collected March, 1915..	Not X.....	<i>B. tysoniana</i> .
Cwebe Forest..	Large.....	Collected March, 1915..	Has X, though not strongly	<i>B. Swinnyi</i> .
Cwebe Forest..	Small.....	Collected March, 1915..	Has X, though not strongly	<i>B. Swinnyi</i> .
Port St. Johns..	Small.....	Sent July, 1914.....	Has X.....	<i>B. Swinnyi</i> .
Bulolo Forest..	Small.....	Collected March, 1915..	Has X.....	<i>B. Swinnyi</i> .†

Mr. Robertson came to the following conclusions:—

1. In mountain forests the large variety (i.e. *B. tysoniana* E.P.P.) has not "X." Yet the natives do use the bark in these forests, so that its medicinal qualities do not depend on its having "X." It appears that only the large variety is found in the mountain forests.

* In the "Remarks" column I have inserted the name of the species which was given in the original by a herbarium number only.

† I have not seen specimens of these, but do not doubt the correctness of the names given.—(E. P. P.)

2. In the coast forests the large variety * usually has "X" to a greater or less extent, but apparently not so strongly as the small variety (*B. Swinnyi* E.P.P.). The latter always has it and usually very strong.

The results of my investigation may be summarized as follows:—

1. Foresters had hitherto only recognized two species, viz., *B. lucens* and *B. tysoniana*.
2. Under the name of *B. tysoniana*, three species have been included—
 - (a) *B. tysoniana*, found in both the mountain and coast forests.
 - (b) *B. Swinnyi*, found only in the coast forests.
 - (c) *B. Stayneri*, only recorded from Natal.
3. The bark of *B. tysoniana*, *B. Swinnyi*, and *B. Stayneri* is used medicinally by the natives.
4. The bark of *B. Swinnyi* has a bitter burning taste and differs in this respect from that of *B. tysoniana* and *B. Stayneri*, in which the bark, while bitter, has not the burning taste.
5. There appear to be two forms of *B. Swinnyi* in the forest of Port St. Johns, known to the foresters as the large and small varieties. The natives are said also to distinguish these, but have the same name for both.

I have been unable to furnish any additional information with regard to the other species recorded.

In conclusion I should like to refer again to Mr. C. C. Robertson's share of this work. Since 1914 he has been slowly accumulating evidence from foresters about the species, and my remarks are only a digest of the information he collected.

With the assistance of the Forest Department, the Division of Botany is undertaking an examination of the barks of the various species as regards their medicinal value.

KEY TO SPECIES.

- Inflorescence quite sessile, ovoid or globose, surrounded at the base by numerous large ovate acuminate silky bracts..... *Swinnyi*.
 Inflorescence distinctly peduncled, rarely sub-sessile, but then bracts at the base never densely silky.
 Mid-rib on lower surface of leaflets covered with long straight hairs. *Stayneri*.
 Mid-rib on lower surface of leaflets glabrous.
 Fruits without woody protuberances; inflorescence silky or tomentose with fine adpressed hairs.
 Leaves usually 2-3-jugate; leaflets obovate, usually rounded at the apex, obtuse..... *lucens*.
 Leaves usually 4-jugate; leaflet oblong-lanceolate, acuminate, acute..... *abyssinica*.
 Fruits with woody protuberances; inflorescence tomentose with spreading hairs appearing almost shaggy..... *tysoniana*.

1. *B. Swinnyi*, Phillips.

Arbor. *Rami* glabri vel juniores serici. *Folia* 10-20 cm. longa, pinnata, 5-jugata; foliola 2-8 cm. longa, 0.9-3.5 cm. lata, oblongo-lanceolata, ovato-oblonga vel ovata, apice obtusa, glabra. *Racemus* sessilis, 2.5-5 cm. longus, circa 3 cm. latus, subglobous vel ovatus. *Bracteae* numerosae, 2 cm. longae, ovatae, acuminatae, sericae.

A large or small tree, with rough bark. *Branches* glabrous or in very young branches velvety. *Leaves* 10-20 cm. long, grouped at the ends of the branches, petiolate, pinnately compound, 5-jugate; petiole 2.5-3.5 cm. long, pubescent in young leaves, glabrous in older leaves; leaflets petiolulate, 2-8 cm. long, 0.9-3.5 cm. broad, oblong-lanceolate or ovate-oblong, or ovate, obtuse or slightly retuse at the apex, rounded at the base, with

* Mr. Robertson here confuses *B. tysoniana* and *B. Swinnyi*, both of which occur in the coast forests.—(E. P. P.)

the mid-rib prominent beneath and the lateral veins distinct, glabrous. *Inflorescence* sessile, 2.5–5 cm. long, about 3 cm. in diameter, subglobose or ovoid in shape, surrounded at the base with numerous ovate acuminate densely silky bracts about 2 cm. long. *Floral-bract* 7 mm. long, ovate, acuminate, very densely silky. *Pedicel* about 4 cm. long, terete, silky. *Sepals* 6 mm. long, ovate, acute, distinctly keeled, silky. *Petals* 1.3 cm. long, about 3 mm. broad at the middle, spathulate-linear, tomentose without, sometimes with two marginal processes at the middle. *Stamens* monadelphous; filaments terete and narrowing above, flattened below and hairy on the flattened portion. *Ovary* 1.5 mm. long, sub-globose, villous; style 1.3 cm. long, terete and densely villous in the lower $\frac{3}{4}$ becoming more or less four-angled and glabrous above; stigma globose. *Fruit* not seen.

Pondoland: In monte propoxe ostium flum St. Johns R., c. 1000 ft., February, *Bolus* 8724; 20 ft., January, *Flanagan* 2884; Port St. Johns, *Swinny in Herb. Forest Dept.*, 1038, 1037; Pongwani Forest, Port St. Johns, January, *Coetzee in Herb. Forest Dept.*, 1866.

2. *B. Stayneri* Phillips.

Rami glabri. *Folia* pinnata, 7–16 cm. longa, 4–5-jugata; petiolus pilosus; foliola 1.7–5.2 cm. longa, 0.8–1.5 cm. lata, oblongo-lanceolata vel elliptica, apice acuta mucronataque, basi paullo cuneata, subtus pilosa vel pubescentia. *Racemus* 3.5–6.5 cm. longus. *Fructus* 2.7–3 cm. longus, ligneus.

Bark thick and rough. *Branches* glabrous, rough with longitudinal furrows. *Leaves* pinnate, petioled, 7–16 cm. long, 4–5-jugate; petiole 1–2 cm. long, very densely pilose; rachis densely pilose; leaflets 1.7–5.2 cm. long, 0.8–1.5 cm. broad, oblong-lanceolate or elliptic, acute and mucronate at the apex rarely rounded, slightly cuneate at the base, the mid-rib deeply sunk on the under surface which is frequently somewhat rugose, prominent beneath and with the lateral veins distinct beneath, glabrous above, pilose or pubescent beneath, especially on the mid-rib. *Inflorescence* a shortly peduncled many-flowered raceme, 3.5–6.5 cm. long; peduncle 1–1.5 cm. long, very densely pilose. *Bracts* 3 mm. long, linear, tomentose, *Pedicels* 5 mm. long, terete, tomentose. *Calyx* 6 mm. long, densely tomentose, divided almost to the base. *Petals* 1.3 cm. long, 3 mm. broad, spathulate-linear, tomentose, reflexed in open flowers. *Filaments* 5 mm. long, semiterete, monadelphous and villous at the base. *Ovary* densely villous; style 3 mm. long, terete; stigma subglobose. *Fruit* 2.7–3 cm. long dehiscing by four valves; valves woody, covered with woody protuberances. *Seeds* reddish in colour, 1.1 cm. long, 6 mm. in diameter, ellipsoid, with a waxy yellow arillus at the base.

Natal, without locality, *Stayner in Herb. Bolus*; Stinkwood Forest, Ingeli, May, *Chilvers in Herb. Forest Dept.*, 1518; Mkanzeni, Riversides, *Henkel in Herb. Forest Dept.*, 2421.

3. *B. lucens* Szysz.

A shrub 8–10 ft. high (ex *Wood*). *Branches* glabrous, with wrinkled greyish bark. *Leaves* petioled, compound, imparipinnate, 2–4-jugate, 10–20 cm. long; petiole 1.5–5.5 cm. long, glabrous, rarely pubescent; leaflets 1.8–9 cm. long, 1.3–4.6 cm. broad, obovate or obovate-elliptic, rarely elliptic, very rarely subacute, slightly narrowed at the base, glabrous, with the mid-rib distinct above, prominent beneath, and with thickened wavy margins. *Inflorescence* a lax axillary or terminal raceme, 6–14.5 cm. long, many-flowered; peduncle pubescent. *Pedicels* 0.4–1 cm. long, terete, densely tomentose with fine adpressed hairs; bracts 0.5 mm. long, ovate, pubescent. *Calyx* gamosepalous; lobes 3–4 mm. long, 2 mm. broad, ovate or ovate-elliptic, obtuse, finely but densely pubescent without and with fine adpressed hairs within, ciliated, the two posterior lobes connate and bifid at the apex. *Petals* dull yellow, 8.5 mm. long, 2.5 mm. broad above, oblong, obtuse, narrowed into an evident claw, pubescent. *Disc* unilateral. *Filaments* all connate at the base, 1 mm. long, terete, glabrous except at the base, the anterior filaments ciliated at the base and forming a tomentose shield 3 mm. long and 2.5 mm. broad; the posterior filaments

ciliated and pubescent at the base; anthers 2.5 mm. long, 1.5 mm. broad, oblong. *Ovary* 1 mm. long, 1.5 mm. broad, subglobose, densely villous, four-locular with a single ovule in each loculus; style 7 mm. long, cylindric, pubescent on the lower half; stigma globose, faintly lobed. *Fruit* 1.7 cm. long, 1.7 cm. in diameter, finely but densely pubescent, four-celled, each cell with a single seed, splitting into four valves when ripe. *Seeds* bright scarlet, 7 mm. long, 7 mm. in diameter, subglobose, wrinkled, flat on one side, convex on the other, with a fleshy yellow-green arillus. *Wood*, *Natal Plants*, 88; *Sim*, *Forests and Forest Flora*, 155; *Journ. Bot.*, XLV, p. 13. *Natalia lucens*, *Hochst.*, *Harv. and Sond.*, *Fl. Cap.*, 1, 369.

Komgha Div., Kei Mouth, 200 ft., August, *Flanagan*, 421; Transkei, Kentani, 20 ft., December, *Miss Pegler*, 726; Pondoland, mouth of St. Johns River, *February*, *Bolus*, 8836; *Flanagan*, 2568; 20 ft., January, Manino Forest, Engcobo Dist., August (leaf only). *Zahn in Herb.*, *Forest Dept.*, 2046; *Natal*, near Durban, 200 ft., February, *Wood*, 7392; March, *Wood*, 7518; Verulam, 100-200 ft., January, *Wood*, 10207; near Durban, *Wood*, 2570; 12666; Port Natal, *Gueinzuis*; Groenberg, February, *Wood*, 808; without precise locality, *Saunders*; near, Durban, 200-300 ft., March, *Wood*, 6307; Swaziland, kloof near Dalriach Mbabane, c. 4800 ft., December, *Bolus*; *Transvaal*, Barberton Dist., Unevoti Creek, Barberton, 3000 ft., February, *Galpin*, 849; *Thorncroft in Herb. Transvaal Mus.*, 18292.

Flanagan, 421, mentioned by Baker as a form with smaller leaflets than the type is slightly confusing, inasmuch as Flanagan gave the same number to two different collections. His No. 421 in the Bolus Herb. was collected at the Kei Mouth at an altitude of 200 ft. in August, 1894, and is the true *B. lucens*. His No. 421 in the Cape Government Herb. was also collected near the Kei Mouth at an altitude of 300 ft., but in January, 1890. The inflorescence in these specimens is compact, oblong in outline, and at the end of a long naked peduncle 9-12 cm. long. The peduncle is fasciated to a slight extent, which will account for the compact inflorescence. There is no difference in the floral structures of these specimens and other specimens of *B. lucens* which I have examined.

4. *B. abyssinica* Fresen.

A tree. *Leaves* petioled, 18-22 cm. long, pinnate; petiole 4-5 cm. long, terete, glabrous; rachis sometimes winged between the uppermost pair of leaflets; leaflets usually in four pairs, 2.5-4 cm. apart, 4-9 cm. long, 1.7-3.5 cm. broad, oblong-lanceolate or ovate-lanceolate, acuminate, acute, entire, or the margin sometimes serrated in the upper half, glabrous. *Inflorescence* racemose, 10-20 cm. long, on a woody peduncle 5-7 cm. long. *Calyx* campanulate, silky. *Petals* five, ligulate-spathulate, twice as long as the calyx, densely and finely silky, finally reflexed. *Stamens* five, in the male flower nearly as long as the petals; filaments monadelphous, silky. *Capsule* about 1 cm. long, 1 cm. in diameter, subglobose 3-4 valved, densely pubescent, without woody protuberances. *Seeds* 6 mm. long, black.

This description was drawn up partly from a fruiting specimen collected by Schimper in Abyssinia and partly from the description in the "Flora Tropical Africa."

Baker, in his monograph on the genus (*Journ. Bot.*, XLV, 12), makes no mention of this species occurring in Natal, though the record is made in the "Flora Tropical Africa," I, 434.

= *B. integrifolia*, Richard, *Fl. Abyss.*, I, 107. This is a tropical African species.

5. *B. tysoniana* Oliv.

A tree 25 ft. high, with a girth of 42 in. (Krid). *Branches* glabrous. *Leaves* petioled, pinnately compound, 4-5-jugate, rarely 3-jugate, 5-20 cm. long; petiole 2-5 cm. long, subtomentose, pubescent to almost glabrous, sometimes white silky; leaflets 1.5-6 cm. long, 0.7-3 cm. broad, oblanceolate, oblong, oblong-elliptic, rarely obovate, rounded or shortly apiculate at the apex, cuneate more rarely rounded at the base, entire, rarely serrated in the upper half, with the mid-rib prominent beneath, glabrous. *Inflorescence* a

dense raceme, distinctly peduncled or subsessile, 6-17 cm. long, cylindric. *Peduncle* usually 1-3.5 cm. long, tomentose with spreading hairs. *Pedicels* 3 mm. long, tomentose with spreading hairs. *Sepals* 5-7 mm. long, 3 mm. broad, ovate, concave, acute or obtuse, tomentose with spreading hairs, two of the sepals united. *Petals* white 1.1-1.5 cm. long, 2 mm. broad, oblong, linear-oblong or linear, narrowing sometimes into a linear claw below, subostuse or acute, villous. *Filaments* 0.8-1.2 cm. long, terete and glabrous in the upper half, expanded villous and monadelphous in the lower half; anthers 1.5 mm. long, oblong in outline. *Ovary* densely villous; style 8 mm. long, densely villous below, glabrous or almost so above; stigma subglobose or globose. *Fruit* 2.2 cm. long, about 2.2 cm. in diameter, subglobose; the valves covered with woody protuberances. *Seed* 1.2 cm. long, about 6 mm. in diameter, more or less keeled on one face, convex on the other, reddish with a yellow arillus at the base.

Komgha Div., Komgha, on Alan Page's farm "Goldust," May (in leaf only), *Galpin*, 8470; Prospect, near Komgha, September, *Flanagan*, 309; Instsubani Forest, February (fruit), *Leigh in Herb. Forest Dept.*, 1840; Wilo Forest, Mqanduli Dist., *Kriel in Herb. Forest Dept.*, 1772; Mboleni Forest, Notineela, October, *Fegen in Herb. Forest Dept.*, 1771; Manina Forest, Engcobo Dist., August, *Zahn in Herb. Forest Dept.*, 2047 (leaf); Cwebe Forest, January (fruit), *Pretorius in Herb. Forest Dept.*, 1854.

Gerrard 1428 from Natal. Description of branches and leaves by Baker in *Journ. Bot.*, Jan., 1907, p. 15.

Arbor vel frutex? Ramulis cortice griseo tectis; foliis saepissime 4-jugis cum impari petiolulato, foliolis petiolulatis oblongis apice apiculatis lateralibus aliquantulum inaequilatereibus oppositis vel passim alternis basi cuneatis vel interdum rotundatis coriaceis margine integris vel plus minus serratis costa superne impressa subtus subconspicua nerviis secundariis tenuibus inter se juxta marginem anastomosantibus foliolis terminalibus oblongo-oblancoelatis foliolis proximis quam distalibus minoribus; rachi omnino exalata.

I have seen a specimen in leaf only (*Zahn in Herb. Forest Dept.*, 2037) which agrees with the above description, except that the leaves are 6-jugate. The leaflets which appear to be young have a long hairy apiculus, and are either entire or serrated in the upper half. In *Tyson*, 6216, in the Bolus Herb. on which Oliver founded his *B. tysoniana*, I find that some of the leaflets have this hairy apiculus and some are also toothed, so that it would appear that Baker is correct in regarding *Gerrard*, 1428, as near *B. tysoniana*, and I am of opinion that it is this species. The distribution of *B. tysoniana* suggests that it will probably be found in Natal.

A REVISION OF THE AFRICAN SPECIES OF SESBANIA.

By E. P. PHILLIPS, Division of Botany, Pretoria, and J. HUTCHINSON (Kew).

THE present paper is an attempt to revise the African species of the genus *Sesbania*, family *Papilionaceae*. Imperfect as it may prove to be, it is long overdue, chiefly because of the great accumulation of herbarium material since the publication of the second volume of the "Flora of Tropical Africa" in 1871, much of this material having remained unnamed or imperfectly determined. The results of this investigation might very well have been more satisfactory to the authors had there been more field notes available regarding the situation, habit, floral colouring, etc., of the specimens accumulated in the various herbaria which they have been able to consult.* That this information is vital in the determination and limitation of the species of *Sesbania*, at least, has been well demonstrated by Prain† in his critical elucidation of the Indian species.

The genus *Sesbania* contains about fifty species which occur in the warmer parts of the world, mainly in or by the sides of streams, lakes, and swamps. It appears to attain its greatest development in tropical Africa, a few of the species extending into South Africa as far as Natal, and into various parts of the Transvaal and Bechuanaland. In the present revision of the species from this area twenty-three are recognized to be distinct. They belong to two sections, nineteen to *Eusesbania* and four to *Daubentonia*, the latter characterized by its four-winged fruits. The third and purely American section into which *Sesbania* was divided by Bentham and Hooker (Gen. Pl., I, 502) is now recognized by American botanists‡ as a distinct genus, *Glottidium*, and the present authors' views coincide in respect to this. They cannot, however, go so far as to accept the American view regarding the generic status of *Daubentonia* which occurs in the three widely separated areas, namely, the south-eastern United States and Mexico, sub-tropical South America, and tropical East Africa. This broken distribution seems to point to a separate origin of the species of *Daubentonia* from the basal stock, *Eusesbania*, species of which occur in all these areas. *Glottidium*, on the other hand, a native of Florida, is well separated from *Sesbania* by its short fruits with only two seeds and the manner of dehiscence, the seeds remaining inside the dry bladder-like endocarp which detaches itself as a whole from the outer shell.

In the case of the African species we have found a most useful and constant character in the nature of the appendages on the claw of the vexillum. In the first five species shown in the key these are long and quite free from the vexillum in their upper half. In

* For the privileges of examining the specimens under their charge, the authors tender their thanks to Sir David Prain, Director, Royal Botanical Gardens, Kew; Dr. A. B. Rendle, Natural History Museum, South Kensington; Dr. I. B. Pole Evans, Director, Botanical Survey, South Africa; Mrs. Bolus, Bolus Herbarium, Capetown; Dr. L. Perinquey, South African Museum; Dr. Schönland, Albany Museum, Grahamstown; the Director, Transvaal Museum, Pretoria; and Mr. Fred Eyles, Salisbury, Rhodesia.

† Prain in Journ. Anat. Soc., Bengal, LXVI, 366-370 (1897).

‡ Small, Fl. South-Eastern Unit. States, Ed. II, 615 (1913).

all the other species they are sessile and adnate to the claw, whilst they are not perceptible in one species, *S. leptocarpa*. These appendages on the claw of the vexillum in *Sesbania* are probably closely connected with pollination, and no doubt, as in some other *Leguminosae*, their function is to clasp and lift open the free stamen in order to give insects ready access to the nectaries.

We have considered it sufficient to restrict ourselves in regard to synonymy to those names to which have been applied to the African species.

KEY TO THE AFRICAN SPECIES OF *SESBANIA*.

I. *Eusesbania*.—*Fruits not winged on the margins of the valves (in S. sphaerocarpa fruits acutely keeled down the middle of the valves).*

Appendages of claw of vexillum free in their upper half or third:

Vexillum yellow, plain or finely speckled on the back with dark spots:

Stems and leaves rather densely villous pubescent:

Flowers solitary or in pairs; leaflets 5–10 pairs, stipules deciduous; appendages to vexillum obtuse.....

1. *S. Goetzei*.

Flowers several in each inflorescence; leaflets more than 10 pairs; stipules persistent; appendages to vexillum acutely acuminate.....

2. *S. speciosa*.

Stems and leaves glabrous or nearly so:

Inflorescence branched; peduncles smooth; West African species.....

3. *S. punctata*.

Inflorescence not branched:

Peduncles smooth.....

4. *S. aegyptiaca*.

Peduncles aculeate.....

5. *S. Dummeri*.

Vexillum and tips of wings suffused throughout with dark purple.....

4. *S. aegyptiaca*
var. *bicolor*.

Appendages of vexillum wholly adnate to the claw or absent:

Flowers very large, 5–10 cm. long; fruits about 30 cm. long; leaflets 3–3.5 cm. long; cultivated species.....

6. *S. grandiflora*.

Stipules foliaceous, broad, about 1 cm. long, more or less persistent:

Peduncles smooth:

Flowers pale blue, not speckled.....

7. *S. caerulescens*.

Flowers yellow, vexillum speckled.....

8. *S. cinerescens*.

Peduncles strongly aculeate.....

9. *S. macrantha*.

Stipules more or less subulate, never foliaceous, and mostly soon falling off:

Stems and leaves or leaves only rather densely pubescent:

Calyx glabrous outside:

Pod not torulose.....

10. *S. pubescens*.

Pod strongly torulose.....

11. *S. Dalzielii*.

Calyx pubescent outside.....

12. *S. Wildemannii*.

Stems and leaves glabrous or only slightly pubescent:

Leaflets usually well over 1 cm. long; leaves nearly always as long or longer than the inflorescence:

Valves of the fruit not keeled down the middle:

Appendages absent from the claw of the vexillum; fruits torulose, much constricted between the seeds.....

13. *S. leptocarpa*.

- Appendages present on the claw of the vexillum :
 Margins of the fruits straight :
 Fruits about 7 mm. broad ; seeds separated
 by partitions of the endocarp ; stems and
 leaf-rachis never aculeate..... 14. *S. pachycarpa*.
 Fruits 3 mm. thick, seeds not separated ;
 stems and leaf-rachis often aculeate.... 15. *S. aculeata*.
 Margins of fruits much constricted between the
 seeds..... 16. *S. arabica*.
 Valves of the fruits sharply keeled on the back,
 Angolan, species..... 17. *S. sphaerocarpa*.
 Leaflets very small, averaging about 5 mm. long ; leaves
 much shorter than the inflorescence :
 Vexillum mottled with purple spots..... 18. *S. microphylla*.
 Vexillum striate, not mottled..... 19. *S. mossambicensis*.

II. **Daubentonia**.—*Fruits broadly winged on the margins of the
 valves ; peduncles aculeate towards the base.*

- Stems not aculeate ; peduncles aculeate only near the base :
 Leaves much shorter than the fruits ; inflorescence very
 few (1–3) flowers ; N.E. Tropical Africa..... 20. *S. tetraptera*.
 Leaves much longer than the fruits ; inflorescence several
 flowered ; Zambezi basin..... 21. *S. Kirkii*.
 Stems aculeate :
 Leaflets about 18 pairs ; inflorescence up to 11 cm. long ;
 staminal sheath shorter than the free parts of the filaments.. 22. *S. hamata*.
 Leaflets about 10 pairs or less ; inflorescence 1–3 flowered,
 very short ; staminal sheath longer than the free part of
 the filaments..... 23. *S. Rogersii*.

EUSES BANIA.

1. **Sesbania Goetzei**, Harms., in Engl. Bot. Jahrb., XXX, 327 (1902).

A shrub 2–3 m. high, branchlets rather densely and softly grey-pubescent, the older ones becoming striate or somewhat angular. *Leaves* shorter than the inflorescence, 3–5 cm. long ; rachis and leaflets densely and softly grey-pubescent ; leaflets about 6–10 pairs, shortly stalked, oblong, rounded and unequal-sided at the base, rounded and very minutely mucronate at the apex, 1–1.5 cm. long, 3–6 mm. broad ; stipules soon falling off, triangular-lanceolate, acute, 3–4 mm. long, pubescent ; stipules minute. *Inflorescence* mostly two-flowered, axillary ; peduncle softly pubescent ; pedicels about 5 mm. long, with two linear-lanceolate fugacious bracteoles towards the apex. *Flower buds* ellipsoid, turbinate at the base, subacute at the apex. *Calyx* widely campanulate, abruptly contracted into a solid turbinate base, about 5 mm. long, with five acute subulate-triangular lobes scarcely 1.5 mm. long, sparsely pubescent outside. *Vexillum* large, orbicular, about 3 cm. long and broad, strongly and closely mottled with purple, with two free linear curved obtuse appendages towards the base about 7 mm. long. *Alae* about 2.5 cm. long and 8 mm. broad. *Carina* 3 mm. long, the claw slightly longer than the broadly boat-shaped limb. *Staminal-sheath* 2 cm. long, free parts of filaments curved upwards and about 8 mm. long. *Ovary* with a very thin line of hairs along the top ; style glabrous. *Fruits* not known.

TROPICAL AFRICA : Tanganyika Territory, Ubungu, Rukwa Lake, moist banks, 800 m., July, W. Goetze, 1115 (type). British East Africa, between Kikuyu and Eldama Ravine, 1898, A. Whyte ; Eldama Ravine, A. Whyte.

A very beautiful species with short leaves, few hairy leaflets, large handsome flowers, the standard petals densely mottled with purple.

2. *Sesbania speciosa*, Taub., ex Engl. in Abh. Preuss. Akad. Wiss., 1894, 21, 42, et in Engl. Pflanzenw. Ost-Afr. C. 213 (1895); *S. Hildebrandtii* Taub., l.c.

Branchlets flexuose, obtusely angular, shortly pubescent. *Leaves* about as long as the inflorescence up to 15 cm. long; rachis and leaflets on both sides fairly densely pubescent; leaflets about 15 pairs, stalked, narrowly oblong, slightly narrowed and almost equal-sided at the base, acutely mucronate at the apex, 2-2.5 cm. long, 6-8 mm. broad; stipules persistent, obliquely lanceolate, very acutely acuminate, 1-1.3 cm. long, about 4 mm. broad at the base, adpressed-pubescent on both sides; stipels minute. *Inflorescence* 6-10 flowered, axillary; peduncle stout towards the base, pubescent; pedicels about 1.5 cm. long, nearly glabrous; bracts persistent, linear, about 6 mm. long, pubescent, margins membranous towards the base; bracteoles deciduous. *Flower buds* not seen. *Calyx* widely campanulate from an acute turbinate base, about 8 mm. long, with five subulate acute teeth about 1.5 mm. long, glabrous outside. *Vexillum* (probably yellow) finely mottled with purple, rather broadly elliptic, shortly clawed, 3 cm. long, 2 cm. broad, with two linear free appendages at the base. *Alae* 3-5 cm. long, 8 mm. broad. *Carina* 3 cm. long, the boat-shaped limb a little longer than the claw. *Staminal-sheath* 2.5 cm. long. *Ovary* slightly hairy. *Fruit* up to nearly 33 cm. long, about 8 mm. broad, rather thick, with undulate edges, nearly glabrous. *Seeds* somewhat kidney-shaped, dark brown.

TROPICAL AFRICA: East African Protectorate, Usambara; Maschena, in meadows, July, 1893, C. Holst, 3508; banks of the Kingani, 6 ft. high, fl. and fr., Hildebrandt, 960 (Herb. Mus. Brit.).

3. *Sesbania punctata* D.C. Prodr., ii, 265 (1825); Baker in Oliv. Fl. Trop. Afr., ii, 133 (1871), partly.

A shrub or slender tree, growing on banks of streams; branches pale straw-coloured when dry, slightly flexuous, sometimes slightly muricate below the leaves, otherwise glabrous. *Leaves* shorter than the inflorescence, 7-10 cm. long; rachis pubescent on the upper side, soon becoming nearly glabrous, not prickly; leaflets 12-22 pairs, shortly stalked, oblong-linear, a little narrowed and sub-equal-sided at the base, shortly and acutely mucronate at the apex, 1.2-2 cm. long, 3-5 mm. broad, glabrous or slightly pubescent on the thickened mid-rib and margins, stipules deciduous, recurved, about 3 mm. long, slightly pubescent; stipels very minute. *Inflorescence* mostly branched, longer than the leaves, up to 20-flowered, often about 12-flowered; peduncle glabrous or very slightly pubescent, especially at the base, not prickly; pedicels slender, up to 2 cm. long, glabrous, with two small caducous bracteoles towards the apex; bracts small and early caducous. *Flower buds* broadly ellipsoid, obtuse, glabrous. *Calyx* broadly campanulate from a turbinate base, about 6 mm. long, with five rather short broadly triangular lobes slightly puberulous on the margins. *Vexillum* finely mottled with purple, about 2.5 cm. long, rather shortly clawed, with two large free acuminate appendages about 5 mm. long at the base. *Alae* 2-3 cm. long, about 6 mm. broad. *Carina* more or less broadly spoon-shaped, 2.5 cm. long, the claw slightly longer than the limb. *Staminal sheath* 2 cm. long, free part of the filaments about 5 mm. long. *Ovary* and style glabrous. *Fruits* up to 24 cm. long, slender, at first torulose, but sometimes becoming nearly straight when ripe, beaked for some time by the persistent style. *Seeds* dark brown, about 4-5 mm. long.

TROPICAL AFRICA: Senegambia: Without definite locality, Perrotet 231 (type); 1839, Guillemin (H.K.) Hendelot (H.K.). Roger in Herb. Gay (H.K.); Richard-tol, 26th January, 1823, Dollinger (H.K.). Northern Nigeria. Sokoto Province, shrub or slender tree overhanging rivers, December, 1910, J. M. Dalziel 324; Abinsi and vicinity, banks of streams, common, 15th December, 1912, J. M. Dalziel 613; Katagum District, banks or streams, J. M. Dalziel 7, Benne River, October, 1907, C. A. Parsons 154, Nupe, sand banks, small shrub with yellow flowers, 1858, C. Barter, 970.

Vernacular: "Alambo" (Dalziel 324, 613).

Sesbania punctata * D.C., has been much misunderstood in herbaria, and has been considered to be widely distributed over tropical Africa and other countries. This is not, however, the case, and it appears to be confined to the banks of streams, in the Savannah country from Senegambia to Northern Nigeria, probably as far as Lake Chad or may be further. Parsons (No. 154) says it is a very common river shrub, 10 ft. high or so, overhanging and sometimes forming dense impenetrable hedges.

The species may be at once recognized by its small *branched inflorescence*, *mottled standard-petal* with very long *free-appendages* and *smooth* (not aculeate) peduncles.

4. *Sesbania aegyptiaca* Poir. Encyc., vii, 128; Pers. Syn., ii, 316; D.C. Prodr., ii, 264 (1825); Baker in Oliv. Fl. Trop. Afr., ii, 134 (1871); Prain in Journ. As. Soc., Bengal, lxi, 367 (1897).

A tall shrub or small tree. *Branches* pilose or pubescent, rarely almost glabrous, usually ribbed. *Leaves* 3·5–13 cm. long, petiole 0·3–1 cm. long, pubescent or villous; rachis flattened or grooved above, pilose, rarely glabrous; leaflets 9–27-jugate; subsessile, 0·6–2·8 cm. long, 1·5–6 mm. broad, oblong, truncate, rounded, or sometimes retuse at the apex, apiculate, scarcely narrowed to the base, pubescent beneath, pubescent or glabrous above, not punctate with minute black dots, with the mid-rib distinct beneath; the terminal leaflets somewhat oblong-obovate. *Stipules* about 5 mm. long, ovate, acuminate, pubescent, usually deciduous, if persistent then coiled. *Inflorescence* an axillary raceme, 2–10 cm. long, 3–8-flowered, usually longer than, sometimes shorter than, the subtending leaf, the base of the peduncle densely or scantily pilose, otherwise usually glabrous. *Bracts* 3–4 mm. long, ovate, acuminate, acute, pilose, deciduous; bracteoles 1·5 mm. long, linear, pubescent without, deciduous. *Pedicels* 0·5–1·5 cm. long, glabrous. *Calyx-tube* 3–5 mm. long, 5–7 mm. in diameter, glabrous, truncate, with five small teeth or lobes 1 mm. long, ovate, usually minutely ciliate. *Vexillum* 1·1–1·9 cm. long, 1·2–1·9 cm. broad, obovate, suborbicular, or transversely oblong, rounded and usually retuse at the apex, with a short claw 2–3 mm. long and with two free appendages above the claw; alae 1·3–1·8 cm. long, 3–6 mm. broad, oblong, rounded or retuse at the apex, with a curved linear claw 5–6 mm. long, and with a projecting tooth from the base of the lamina forming a horse-shoe-shaped loop with the claw; carina 1·4–1·5 cm. long, 5–6 mm. broad, almost plano-convex in outline, with a linear claw 6–8 mm. long, and a projecting tooth from the lamina forming a horse-shoe-shaped, or distinct, loop with the claw. *Staminal-sheath* 0·8–1·1 cm. long, 2–3·5 mm. broad at the base, with evident or distinct veins; free portion of filaments arcuate; anthers 0·75 mm. long, oblong; free stamen bent near the base. *Ovary* 1–1·3 cm. long, sub-compressed, glabrous, with two longitudinal bands; style 3–5 mm. long, arcuate; stigma small, capitate. *Fruit* twisted, 6–14 cm. long, 3–4 mm. thick, subterete, long-acuminate, acute, glabrous, divided into compartments within. *Seeds* oblong, truncate at both ends, reddish-brown, 4 mm. long, 2 mm. broad, glabrous.

TROPICAL AFRICA: Senegambia: *Perrotet*, Sierra Leone, *Vogel* 24. Northern Nigeria: Kontagora, shrub or tree by streams, *J. M. Dalziel* 38; 50 miles from Maifoni, Bornu, *A. C. Parsons*. Lake Chad and Bornu, *Talbot* 1242. Attah, branched shrub, *Vogel* 45. Congo: *Burton*; Stanley Pool, *Hens* 348. Angola Kiteve, Kunene River, *Baum* 955. Eastern Sudan: Kordofan, *Pfund* 404; *Kotschy* 72, 223, 347, 539, White Nile, *Brownell*; *Lynes*; *Schweinfurth* 998; *Muriel* 107; Kassala, *Schweinfurth*, tree near wells, Mut Oasis; Dakila, *MacDougal* and *Sykes* 187 (Herb. Mus. Brit.). Khartoum, *Schweinfurth* 865. Sedan, *Schweinfurth* 537. Blue Nile, *Muriel*, 69; *Schweinfurth* 962; near Matamma, *Schweinfurth* 1865. Colonia Eritrea: Adi Gana, *Pappi* 192; Abyssinia: *Cockburn*; *Quartin-Dillon* and *Petit* 167; *Wellby*; *Schweinfurth* and *Riva*, 681. British East Africa: various localities; *James*; *Elliot* 313; *Whyte-Grenfell*; *Dümmer* 1781, 2008; *Dowson* 336;

* De Candolle's description reads as follows:—

S. punctata, herbacea glabra, foliis oblongo-linearibus obtusissimis mucronatis, 10–30-jugis, petiolo laevi superne pubescente, racemis multifloris, corolla calyce sextriplo longiore, leguminibus compressis subtorulosis rachi duplo triplove longioribus—in Senegal. (*Perrottet*), Flores flavi lin. 8–9 longi, vexillo punctis purpureis extus maculato (v.s. comm. a cl. *Perrottet*).

Kassner 623, 628, Uganda; *Speke and Grant* 710; *Whyte*; *Scheffler* 227; *Scott Elliot* 7547, 6560, *Dümmer* 225; *Bagshawe* 491, 99, Tanganyika Territory: *Hildebrandt* 961; *Holst* 3442; *Riddelsdell* 150; *Goetze* 53; Rhodesia: Livingstone, *Rogers*, 7227; Victoria Falls, *Allen* 7; *Flanagan* 3084, 3113; *Galpin* 7046; *Kolbe* 3132; *Lamb* 2293; *Rogers* 5099, 5095, 7407, 13127, 13190, 13289, 13465; Matoppos Hills, *Rogers* 5159. Portuguese East Africa: *Kirk*; *Rikatla*, *Jumod*, 229; Delagoa Bay, *Scott*, Lourenço Marques, *Schlechter* 11585; *Bolus* 7728. Maputa River, *Maputaland Exped. in Herb. Transvaal Mus.* 14402. Gazaland: Lower Buzi, *Swynnerton* 1393.

SOUTH AFRICA: Transvaal: Messina, *Rogers* 19404, Komatipoort, *Rogers* 423, 440, and in *Herb. Transvaal Mus.* 2604, Crocodile Poort; *Galpin* 1074; *Bolus* 7727, Barberton; *Pole Evans* 2944; Sheba Siding, *Thornecroft in Herb. Transvaal Mus.* 18235. Barberton, *Rogers* 20314. Swaziland, *Miss Stewart in Herb. Transvaal Mus.* 8975. Natal: Umilalasi, *Wylie in Herb. Wood* 8551; near Stanger, *Wood* 3859; near Durban, *Gueinzus*; *Gerrard* 1070; Umgeni, *Wood* 9562.

Var. *bicolor*, *Wight and Arn.* Prodr. 214 (1834); Prain in Journ. As. Soc., Bengal, lxvi, 367 (1897). *Sesbania atropurpurea*, Taub. in Engl. Bot. Jahrb., XXIII, 188 (1896); like the typical form in vegetative characters, but the standard petal entirely suffused with purple.

TROPICAL AFRICA: Senegal: *Perrottet* 232. Egyptian Sudan: Khartoum, in the garden of the Catholic Monastery and cultivated in the gardens below the town on the Blue Nile, 20–30 ft. high, *Schweinfurth* 796, 779. Cordofan, *Kotschy* 35.

SOUTH AFRICA: Natal: near Durban, *Wood* 8890.

A common species in most tropical countries. The tropical form of *Sesbania aegyptiaca* is said to have uniformly yellow flowers; a second form described by Persoon as *S. picta* has the standard finely mottled with purple. Whilst we have not considered it worth while to distinguish these two conditions, owing to the difficulty of determining them from dried specimens, we have maintained the var. *bicolor* of Wight and Arnot, which is probably only found in cultivation, and has the standard petal entirely suffused with purple. In Africa, as in parts of India, the mottled form appears by far the most common.

According to Prain (*l.c.*), *S. aegyptiaca* is a very familiar hedge plant in Indian fields and gardens. Its wood is still (in India) highly reputed as a source of charcoal for gunpowder manufacture.

5. *Sesbania Dummeri** n. sp.

A slender shrub up to 5 cm. high, branchlets glabrous except when quite young; nearly terete. *Leaves* shorter or nearly as long as the inflorescence, 8–11 cm. long; rachis glabrous or slightly pubescent, not prickly; leaflets 14–16 pairs, stalked, oblong-linear, slightly narrowed and sub-equal-sided at the base, rounded and distinctly mucronate at the apex, 1.5–2 cm. long, about 4 mm. broad, glabrous or with a few minute hairs on the margin; stipules sub-persistent, linear or linear-lanceolate, acutely acuminate,

* *Sesbania Dummeri* Phillips et Hutchinson, sp. nov.

Affinis *S. punctata* DC. et *S. aegyptiaca* Poir., sed ab. ambilus pedunculis basin versus aculeatis inflorescentiis simplicibus differt.

Frutex gracilis usque ad 5 m. altus; ramuli maturi glabri, fere teretes. *Folia* inflorescentia breviora vel fere aequilonga; rachis glaber vel leviter pubescens, haud aculeatus; foliola 14–16-juga, petiolulata oblongo-linearia, basi leviter attenuata et subaequilata, apice rotundata et mucronata, 1.5–2 cm. longa, circiter 4 mm. lata, marginibus glabris vel minute pubescentibus; stipulae subpersistentes, lineares vel lineari-lanceolatae, acutae acuminatae, 7–10 mm. longae, sicco-brunneae, parce ciliolatae; stipellae minutae. *Inflorescentia* racemosa, folia aequilonga vel longior, usque ad 8-flora; pedunculi basin versus aculeati, pedicellis gracilibus, usque ad 1 c. longis, apicem versus bracteolis parvis caducis munitis; bractaeae mox caducae. *Alabastra* oblique ellipsoidea, apice obtuse contracta. *Calyx* e basi turbinate late et leviter oblique, campanulatus, circiter 7 mm. longus, lobis 5 late triangularibus mucronulatis marginibus intra puberulo-ciliatis. *Vexillum* flavum, brunneo maculatum, circiter 2.5 cm. longum, breviter unguiculatum, ungue appendicibus oblique lanceolatis liberis circiter 7 mm. longis munito. *Alae* 2.5 cm. longae, circiter 7 mm. latae. *Carina* 2 cm. longa, limbo rhomboideo. *Tubus staminis* 2 cm. longus, filamentorum partibus liberis 6 mm. longis. *Ovarium* et stylus glaber. *Fructus* junior leviter torulosus, maturus non visus.

7–10 mm. long, brown when dry, sparingly ciliolate; stipels minute. *Inflorescence* racemose, as long or longer than the leaves, up to about eight-flowered; peduncle finely prickly, especially towards the base; pedicels slender, up to 1 cm. long, with two small caducous bracteoles towards the apex; bracts very early caducous. *Flower-buds* obliquely ellipsoid, obtusely contracted at the apex. *Calyx* broadly and slightly obliquely campanulate from a turbinate base, about 7 mm. long, with five broadly triangular mucronulate lobes puberulous-ciliolate within the margin. *Vexillum* yellow, finely mottled with brown, about 2.5 cm. long, rather shortly and broadly clawed, with two large obliquely lanceolate free appendages about 7 mm. long. *Alae* 2.5 cm. long, about 7 mm. broad. *Carina* 2 cm. long, the claw about as long as the more or less rhomboid limb. *Staminal sheath* 2 cm. long, free parts of the filaments 6 mm. long. *Ovary* and style glabrous. *Young fruits* slightly torulose; mature ones not seen.

TROPICAL AFRICA: Uganda: Kimerema, 4000 ft., margins of swamps, flowers yellow, spotted with brown, September, 1913, *R. Dümmer* 225. Without definite locality, *A. Whyte*.

6. *Sesbania grandiflora* Poir. *Encycl.*, vii, 127 (1806); *Pers. Syn.*, ii, 316 (1807).

A small tree or shrub; branchlets rather thick, very shortly and softly pubescent when young. *Leaves* much longer than the inflorescence, up to about 20 cm. long; rachis subterete, slightly pubescent or glabrous; leaflets about 15–20 pairs, stalked, rather elongate-oblong, rounded and slightly unequal-sided at the base, rounded and minutely mucronate at the apex, about 2.5–3 cm. long and 0.5–1 cm. broad, thinly chartaceous, shortly pubescent or glabrous, often glaucous, with about six pairs of lateral nerves; stipules deciduous; stipels subulate, minute. *Inflorescence* few-flowered, axillary; peduncle softly pubescent. *Flowers* large and showy, usually about 10 cm. long, pink or red. *Fruits* about 30 cm. long, flat.

This well-marked species, the native country of which is doubtful, is widely cultivated in the tropics for its handsome flowers. The leaves are often used as a substitute for spinach. We have seen African specimens from Senegambia (*Herb. Gay*), Gold Coast (*Axim, T. F. Chipp*, 391), and Lagos (*Dr. J. W. Rowland*).

7. *Sesbania caerulescens* Harms in Warb. Kunene-Sambesi Exped., 260 (1903). *S. Hockii*, De Wild, in Fedde Rep., XI, 544 (1913).

An erect plant about 3–5 m. high (*Eyles*); stems ribbed, glabrous, marked with brown lines. *Leaves* 5–13 cm. long; petiole 0.8–1.1 cm. long, glabrous; rachis channelled above, glabrous; leaflets 10–18-jugate, subsessile, 0.3–1.8 cm. long, 1.5–5 mm. long, oblong, obtuse, with an acute apiculus 0.5 mm. long, slightly narrowed at the base, punctate above with minute black dots, glabrous. *Stipules* 7 mm. long, oblanceolate, with a long point striped with brown, glabrous, deciduous. *Inflorescence* an axillary raceme, 4–6.5 cm. long, 1–4-flowered, as long as or shorter than the subtending leaf; bracts 7 mm. long, lanceolate, long-acuminate, striped with reddish-brown, glabrous, deciduous; bracteoles 3 mm. long, linear, otherwise similar to the bracts; pedicels 0.8–2.3 cm. long, glabrous. *Calyx-tube* spotted, 6–7 mm. long, 6.5–7 mm. broad, campanulate, glabrous; teeth 3 mm. long, ovate, acuminate, acute, ciliated. *Vexillum* a "washed-blue" (*Eyles*), spotted, 2.5–2.9 cm. long, 2.4–2.85 cm. broad, rounded, retuse at the apex, with a claw 3 mm. long. *Alae* spotted, 2.5–2.6 cm. long, 0.9–1 cm. broad, obtuse, with a linear claw 5 mm. long, 2 mm. broad, and with a downward projecting tooth. *Carina* spotted, 2–2.5 cm. long, 0.75–1.1 cm. broad, obtuse, with a linear claw 1–1.2 cm. long and a tooth from the lamina slightly projecting upwards. *Staminal sheath* 1.7–2 cm. long, 3–4 mm. broad at the base, with the free portion of the filaments arcuate; free stamen curved at the base. *Ovary* spotted, 1.7–2 cm. long, subcompressed; style 7–8 mm. long, arcuate, glabrous; stigma small, ellipsoid. *Fruit* 23–25 cm. long, 3.5–4 mm. broad, linear, long acuminate, with a broad suture on one side, somewhat keeled on the other, glabrous. *Seeds* dark reddish-brown, 5 mm. long, 2.5 mm. broad, oblong, slightly narrowed at each end, glabrous.

TROPICAL AFRICA: Angola: Miane River at Kavamba, 1175, m.; swampy ground, March, 1900, *Baum* 782 (type). River Kumpulua-Kuito, *Gossweiler* 3195. Luco plantations, an annual, erect, glaucous green herb up to 15 ft. high, fls. pale blue, in inundated marshes, February, 1906, *Gossweiler* 3745 (Herb. Mus. Brit.). Congo State: between Chiniamia and Elisabethville, *Hock*. Rhodesia: Salisbury, April, *Flanagan*, 3121; Cleveland Dam. In colonies on most land 5000 ft., 10 ft. high, February, *Eyles*, 600.

8. *Sesbania cinerascens* Welw. ex Baker, in Oliv. Fl. Trop. Afr., ii, 134 (1871); Hiern in Cat. Afr. Pl. Welw., i, 231, under *Sesban* (1896).

A tall arborescent herb or shrub up to 5 m. high, ashy grey in colour. Branches glabrous, smooth. Leaves 10–24 cm. long, 14–30-jugate; petiole 1–2.5 cm. long, glabrous; leaflets 14–30 pairs, subsessile, oblong or oblong-linear, rounded above, minutely punctate, usually unequal at the base, 0.7–2.8 cm. long, 3–8 mm. broad, glabrous, with the mid-rib distinct beneath. Stipules foliaceous, 1–2 cm. long, ovate, acuminate, soon deciduous. Inflorescence axillary, racemose, 11–24 cm. long, rarely only 7 cm. long, longer or shorter than the subtending leaf; peduncle glabrous; pedicels 1–2 cm. long, glabrous; bracts 1.5–5 mm. long, setaceous, caducous. Flowers yellow. Calyx-tube 5–6 mm. long, campanulate, with five distinct veins, glabrous; lobes 1.5–2 mm. long, ovate, acute, woolly-ciliate. Vexillum spotted, 1.8–2.1 cm. long, 1.7–2.2 cm. broad, obovate, reflex at the apex, narrowed into a short claw, with two ridges above the claw; alae 1.8–2 cm. long, 6–7 mm. broad, oblong or oblong-obovate, rounded above, produced into a curved claw 5–6 mm. long, with a downward-projecting tooth above the claw; carina 1.7 cm. long, obtuse, produced into a linear claw 1 cm. long, and with a downward projecting tooth above the claw. Staminal sheath 1.1–2 cm. long, 3.5–6 mm. broad at the base; free portion of filaments arcuate; anthers 0.5 mm. long, oblong. Ovary 0.9–1.4 cm. long, linear, with two longitudinal bands; style 5 mm. long, at right angles to the ovary; stigma small, capitate. Fruit 26 cm. long, semi-terete, acuminate. Seed 9 mm. long, 3 mm. in diameter, oblong, rounded at each end, glabrous.

TROPICAL AFRICA: Angola: Pungo Andongo, in thickets near the River Lombe, fl. and fr. Mar. *Welwitsch*, 1999; in bushy places by streams, near Pedra Cabondo, fl. and fr. Apr., *Welwitsch*, 2000. Rhodesia: Matoppos, c. 4000 ft., May, *Rogers* 7924; Mazoe near Salisbury, April, *Flanagan* 3122, Victoria, C. F. H. *Monro*, 1932; near Bulawayo, *Rand* 448; *Eyles* 1235, 1514.

SOUTH AFRICA: Natal: Itafamasi, *Wood*, 882, Insuzi, April, *Wood*, 5311, and in Natal Govt. Herb. 8677; Natal, *Gerrard* 32. Transvaal: Barberton Div., banks of Kaap River, 2000, September, *Galpin* 1074 B; Barberton, *Rogers* 21567, 23891; Pietersburg Div., Macoutsie River, *Breijer* in Herb. Transvaal Mus. 18664.

Mr. Eyles says in regard to his specimen No. 1235: "In December, 1917, I planted here several of the seeds of No. 1235 (over fourteen years old) after soaking forty-three hours in water. Germination was 100 per cent., and one of the plants has been growing since. It is about 12–15 ft. high, and in March it was profusely covered with medium-sized yellow flowers which never opened until 4.30 p.m. daily."

9. *Sesbania macrantha* * Welw., MSS. *S. punctata* Hiern. in Cat. Afr. Pl., Welw., i, 231, partly, non D.C.

* *Sesbania macrantha* Welw. mss., sp. nov. affinis *S. cinerascens* Welw. sed. caulibus et pedunculis dense aculeatis differt.

Herba usque ad 5 m. alta; rami leviter costati, plerumque aculeati. Folia 5.5–18 cm. longa; petioli 0.7–2 cm. longi, glabri vel parce aculeati; foliola 5–34-jugata, petiolulata, 0.9–2.4 cm. longa, 2–6 mm. lata, oblonga vel oblongo-linear, apice rotundata et mucronata, basi inaequalia, glabra, punctis minutis nigris; stipulae usque ad 1.4 mm. longae, lanceolatae, acuminatae foliaceae, superersistentes. Inflorescentia 13–20 cm. longa, 6–14-flora; pedunculi inferne aculeati; bractae mox deciduae; pedicelli 0.6–1.5 cm. longi, glabri. Calycis tubus 5–6 mm. longus, campanulatus glaber, dentibus 1 mm. longis, ovatis, subacuminatis, lanato-ciliatis. Vexillum 1.7 cm. longum, 2 cm. latum, rotundatum, retusum. Alae 1.8 cm. longae, 8 mm. latae, obovatae, obtusae. Carina 1.5 cm. longa, 5.5 mm. lata. Vagina staminialis 1.2 cm. longa. Ovarium compressum. Fructus 25–29 cm. longus, leviter compressus, circiter 6 mm. latus, acuminatus marginibus incrassatis. Semina 6 mm. longa, 3 mm. lata, oblonga, utrinque rotundata, glabra.

Herb. up to 5 m. high. *Branches* slightly ribbed, usually covered with prickles, glabrous. *Leaves* 5.5–18 cm. long; petiole 0.7–2 cm. long, glabrous, sometimes with a few prickles; leaflets 5–24-jugate, petiolulate, 0.9–2.4 cm. long, 2–6 mm. broad, oblong to oblong-linear, rounded and mucronate at the apex, unequal at the base, glabrous, punctate with minute black dots; petiolule 1 mm. long; stipules up to 1.4 mm. long, lanceolate, acuminate, foliaceous, subsistent. *Inflorescence* 13–20 cm. long, 6–14-flowered; lower portion of peduncle covered with prickles; bracts deciduous, not seen; pedicels 0.6–1.5 cm. long, glabrous. *Calyx-tube* 5–6 mm. long, campanulate, five-veined, glabrous; teeth 1 mm. long, ovate, subacuminate, woolly-ciliate. *Vexillum* 1.7 cm. long, 2 cm. broad, rounded, retuse above, produced into a claw 2 mm. long; alae 1.8 cm. long, 8 mm. broad, somewhat obovate, obtuse, produced into a curved linear claw 5 mm. long, and with an inclined tooth forming a horse-shoe-shaped loop with the claw; carina 1.5 cm. long, 5.5 mm. broad, obtuse, with a downward projecting claw. *Staminal sheath* 1.2 cm. long, 6 mm. broad at the base; the free portion of the filaments arcuate; free stamen kneed near the base. *Ovary* compressed, with two longitudinal bands; style arcuate; stigma capitate. *Fruit* 25–29 cm. long, somewhat compressed, about 6 mm. broad, acuminate; the valves thickened. *Seeds* 6 mm. long, 3 mm. broad, oblong, rounded at each end, glabrous.

TROPICAL AFRICA: Uganda: Kampala, tall handsome shrub, *Scott Elliot* 7267. "Elephant grass" land near Kasala Forest, slender, up to 15 ft. high, fls. yellow, *Dümmer* 742. British East Africa: Aberdare Mts., 5000–6500 ft., *Battiscombe* 8; two days' march from Mumias, *A. Whyte*; Nyasaland: Nyika Plateau, 6000–7000 ft., *A. Whyte*; Congo: Elizabethville, *Rogers* 26224; Southern Rhodesia: Odzani River Valley, Manica Dist., *A. J. Teague* 15; Angola: Huilla, in bushy places along the banks of the River Monino, fls. and fr., *Welwitsch* 1995; Pungo Andongo, in palm groves on the banks of the River Cuanza, near Sansamanda, rather rare, *Welwitsch* 1997.

SOUTH AFRICA: Transvaal: Waterval Onder, *Jenkins in Herb. Transvaal Mus.* 6724.

This appears to be a very fine species, resembling in its aculeate stems and peduncles *S. aculeata*, but with much larger stipules, inflorescence, and fruits than in that species.

10. *Sesbania pubescens*, D C. Prodr. ii, 265 (1825); Baker in Oliv. Fl. Trop. Afr. ii, 135 (1871); Hiern. in Cat. Afr. Pl. Welw. i, 231 (under *Sesban*) (1896). *Emerus pubescens*, Schum. Guin. Pl., 354 (1827).

A slender shrub growing near water; branches ascending, slightly angular when older, finely and rather laxly pubescent; young growths densely silky-pubescent. *Leaves* three or four times as long as the inflorescence, about 12 cm. long; rachis thinly pubescent; leaflets 12–18 pairs (usually about 17 pairs), shortly stalked, elongated-oblong, rounded on one side, slightly narrowed on the other, rounded and with a prominent subulate mucro at the apex, 1.5–2.3 cm. long, 5–6 mm. broad, glabrous but densely covered with minute black spots above, adpressed silky-pubescent below; stipules soon falling off, linear-subulate, very acute, 6–7 mm. long, with narrowly membranous edges, slightly pubescent up the middle; stipels minute. *Inflorescence* about 3 cm. long, or up to 8 cm. very slender, simply racemose or rarely branched; peduncle thinly pubescent; pedicels about 5 mm. long; bracts like the stipules, but with wider membranous margins, soon falling. *Flowers* yellow, about 1 cm. long; buds ellipsoid, beaked at the apex, about 5 mm. long, glabrous. *Calyx* campanulate from a broadly turbinate base, 5 mm. long, with five broadly triangular acute lobes slightly puberulous on the margin. *Vexillum* probably not spotted, about 8 mm. long, with two wholly adnate appendages on the claw. *Alae* about 8 mm. long. *Carina* limb as long as the claw. *Ovary* and style glabrous. *Fruits* curved, about 15 cm. long, 3 mm. broad, with thickened margins, acutely beaked, glabrous. *Seeds* oblong, reddish-brown, about 3 mm. long.

TROPICAL AFRICA: Senegambia: Marshes near Nboro, 1840, *Brunner*; without locality, *Thiery*; *Perrotet* 228. Gold Coast: Accra, *T. W. Brown* 419; Axim, *T. F. Chipp* 426; Aburi Gardens, *W. H. Johnson* 950. Dahomey: Cotanou, *Debeaux* 150

(Herb. Mus. Brit.). Western Sudan: between Madoo and Berirem, *A. Chevalier* 10095. St. Thomas Island, *A. Moller* 86. Angola: Loanda: rather rare in moist sandy places (near Praia de Zamba Grande, fl. and fr. Mar., Apr., *Welwitsch* 1992. Tanganyika Territory, *Stuhlmann* 7906.

This species is distinguished especially by its very small flowers and slender inflorescences. *S. sericea* D.C. from Ceylon is a very closely allied species, with fruits usually broader than those of *S. pubescens*.

11. *Sesbania Dalzielii* * n. sp.

A slender shrub up to 2 m. high; branches weak, angular when young, slightly pubescent, at length striate. *Leaves* usually longer than the inflorescence, 15–25 cm. long; rachis and leaflets silky pubescent; leaflets up to 40 pairs, shortly stalked, linear-oblong, rounded and acutely mucronate at the apex, obliquely acute at the base, 0.8–2.5 cm. long, 2–3.5 mm. broad, coriaceous, appressed silky-pubescent on both surfaces; stipules linear-subulate, acute, very soon falling off; stipels very minute. *Inflorescence* slender and laxly flowered; the uppermost ones longer than their subtending leaves; peduncle pubescent towards the base; pedicels slender, glabrous, bibracteolate towards the apex; bracteoles linear, 2 mm. long, falling off before the flower opens. *Flower-buds* ellipsoid-turbinate, shortly acuminate, glabrous. *Calyx* campanulate, acutely turbinate at the base, 5 mm. long, glabrous outside, ciliate within the five acutely acuminate lobes. *Vexillum* with two small sessile appendages on the claw. *Ovary* glabrous. *Fruit* about 20 cm. long, strongly torulose, shortly and acutely beaked, about 25-seeded. *Seeds* oblong-ellipsoid, 5 mm. long, green and shining.

TROPICAL AFRICA: Northern Nigeria: Abinsi and vicinity in marshy places, *J. M. Dalziel* 614 (type). Katagum District, marshy places, *J. M. Dalziel* 8, "Niger," *Bakie* 9. Jeba, on the Niger, *Barter*. Western Sudan: Koulikoro, *Chevalier* 3390.

12. *Sesbania Wildemannii*, nom. nov.

Sesbania affinis De Wild, in Ann. Mus. Cong. Ser. V, 1, 141 (1904), non Schrad.

Stems terete, finely longitudinally grooved, glabrous. *Leaves* up to about 9 cm. long; rachis angular, shortly pubescent; leaflets about 20–25 pairs, linear-oblong, rounded and mucronate at the apex, obliquely unequal-sided at the base, about 8 mm. long and 2 mm. broad, grey, at length glabrous above, finely appressed-pubescent below; stipules small, deciduous. *Inflorescence* few-flowered, very slender; peduncle slightly pubescent; pedicels nodding, 7 mm. long; bracts and bracteoles deciduous. *Flowers* about 1.5 cm. long. *Calyx* ribbed, pubescent outside, with five acutely triangular teeth. *Vexillum* finely mottled outside, with two sessile appendages on the claw. *Fruits* very narrow, about 15 cm. long and 2.5 mm. broad, glabrous, straight or very slightly torulose. *Seeds* oblong, rounded at the ends, 3 mm. long, brownish.

TROPICAL AFRICA: Belgian Congo: Between Leopoldville and Mombazi, *Gillet*. 2626.

We have seen rather an imperfect specimen of this species, which appears distinct on account of its hairy calyx tube.

* *Sesbania Dalzielii*, *Phillips et Hutchinson*, sp. nov., affinis *S. pubescenti* DC., sed fructibus torulosis differt.

Frutex gracilis usque ad 2 m. altus; rami debiles, primum angulares, leviter pubescentes, demum striati. *Folia* inflorescentia plerumque longiora, 15–25 cm. longa; rachis et foliola sericeo-pubescentia; foliola usque ad 40-jugata, breviter petiolulata, lineari-oblonga, apice rotundata et acute mucronata, basi oblique acuta, 0.8–2.5 cm. longa, 2–3.5 mm. lata, coriacea, utrinque appresse sericeo-pubescentia; stipulae lineari-subulatae, acutae, mox deciduae; stipellae minutissimae. *Inflorescentia* gracilis et laxiflora; pedunculi basin versus pubescentes; pedicelli graciles, glabri, bibracteati; bracteolae lineares, 2 mm. longae, mox deciduae. *Alabastra* ellipsoideo-turbinata, breviter acuminata, glabra. *Calyx* campanulatus, basi acute turbinate, 5 mm. longus, extra glaber, lobis 5 acute acuminatis intra ciliatis. *Vexillum* basin versus appendicibus sessilibus instructum. *Ovarium* glabrum. *Fructus* circiter 20 cm. longus, valde torulosus, breviter et acute rostratus. *Semina* circiter 25, oblongo-ellipsoidea, 5 mm. longa, viridia et nitida.

13. *Sesbania leptocarpa*, D.C. Prodr. ii, 265 (1825); Guill. et Perrott. Fl. Seneg. i, 199, (1830-33); Baker in Oliv. Fl. Trop. Afr., ii, 135, partly (1871).

Stem herbaceous, sparingly branched, glabrous, finely grooved. *Leaves* up to 9 cm. long, longer than the inflorescence; rachis very slightly aculeate towards the base; leaflets 8-18 pairs, distinctly stalked, narrowly oblong, very unequal-sided at the base, rounded and finely mucronate at the apex, 1-1.3 cm. long, about 2.5-3 mm. broad, finely punctate with numerous black spots on both surfaces, otherwise glabrous; stipules very soon falling off, linear-lanceolate, acute, about 2.5 mm. long, slightly hairy towards the tips; stipels minute, like little points. *Inflorescence* a simple raceme, about three-flowered; peduncle slender, slightly aculeate towards the base; bracts like the stipules, very soon falling off; bracteoles small, at the base of the calyx tube. *Flowers* small, about 1 cm. long. *Calyx* turbinate-campanulate, 4 mm. long, glabrous, with five triangular subacute teeth slightly puberulous on the margin. *Vexillum* suborbicular, striate and slightly punctate, about 1 cm. long; claw without appendages at the base or only slight thickenings representing them. *Alae* and keel striate. *Ovary* and style glabrous. *Fruits* 12-15 cm. long, 3 mm. broad, torulose, with a dark blotch between the joints, acutely acuminate, 14-24 seeded. *Seeds* black, 3 mm. long.

TROPICAL AFRICA: Senegambia: Moist plains around St. Louis, at north Point, near the Walo River, Richard-Tol, Dagana, etc., *Hendelot* 489; *Perrottet* 233.

This species is distinguished by its slightly aculeate peduncles, by the *absence of appendages* from the claw of the vexillum, and the torulose blotched fruits. It is apparently confined to Senegambia.

14. *Sesbania pachycarpa* D.C. Prodr. ii, 265 (1825); Guill. et Perrott. Fl. Seneg. 200, t. 50 (1830-33); Baker in Oliv. Fl. Trop. Afr., ii, 134, partly (1871). *Sesban phaeocarpus*, Hiern. in Cat. Afr. Pl. Welw. i, 232, partly, non Welw.

Stems herbaceous, pithy, glabrous. *Leaves* 40 cm. long, many times longer than the inflorescence; rachis contracted at the base, glabrous; leaflets very numerous, linear-oblong, truncate or rounded and mucronate at the apex, slightly oblique at the base, 1-2 cm. long, 3-5 mm. broad, glaucous-green, glabrous, very minutely punctate with black spots; stipules linear, nearly 1 cm. long, glabrous. *Inflorescence* short, slender, few-flowered, or sometimes reduced to two or three flowers; peduncle smooth; pedicels up to 1 cm. long, with two early deciduous bracteoles towards the apex. *Calyx* broadly campanulate, from a sharply turbinate base, 5 mm. long, finely puberulous within the margins of the acutely triangular lobes. *Petals* yellow; the vexillum densely mottled with dark spots. *Appendages* to vexillum adnate to the claw. *Fruits* erect, curved, about 15 cm. long and 7 mm. broad, when ripe acutely beaked, with straight margins to the valves, but with slight constrictions between the seeds between the margins. *Seeds* separated by partitions of the sub-membranous endocarp, broadly oblong, truncate at each end, 3.5 mm. long, rarely 3 mm. thick, pale brown.

TROPICAL AFRICA: Senegambia: *Perrottet* 230 (Herb. Mus. Brit.). Northern Nigeria: Bornu, *E. Vogel* 79; Katagum District, *J. M. Dalziel* 9; near Abinsi, *J. M. Dalziel* 615. Eastern Sudan: South Kordofan, *Pfund* 134; *Kotschy* 72. "Upper Nile." *Freeman and Lucas*. Abyssinia: Terrefera Valley, *Schimper* 775. Angola: Loanda, *Welwitsch* 1991 (Herb. Kew. non Herb. Mus. Brit.); Mossamedes, *Welwitsch* 1996b.

Known in Nigeria as "Zamarke" (*Dalziel*).

15. *Sesbania aculeata* Pers. Synops. ii, 316, partly (1807); D.C. Prodr. ii, 265 (1825); Prain in Journ. Asiat. Soc., Bengal, lxvi, 369 (1897). *Sesban punctatus*, Hiern. in Welw. Cat. Afr. Pl. i, 230, partly. *Sesban aegyptiacus* Hiern, *l.c.*, 231, non Poir.

A weed of moist places often up to 5 m. high; stem patently branched from the base to apex, often aculeate, but sometimes smooth. *Leaves* up to 25 cm. long; rachis generally aculeate below; leaflets about thirty pairs, generally about 1.5 cm. long and

3 mm. broad, glabrous and often glaucous; stipules linear-lanceolate, very acute, about 7 mm. long, glabrous, often sub-persistent. *Inflorescence* mostly rather few-flowered and much shorter than the subtending leaf; peduncle aculeate or smooth; pedicels slender, about 5 mm. long, glabrous. *Flowers* yellow, about 1.5 cm. long. *Calyx* tube 3 mm. long, campanulate, glabrous; teeth 1-2 mm. long, subacute. *Vexillum* yellow spotted on the back, with two sessile linear appendages on the claw. *Alae* oblong or oblong-obovate. *Carina* 1.3 cm. long, 4 mm. broad. *Staminal sheath* 1.2 cm. long. *Ovary* glabrous. *Fruits* 15-20 cm. long, straight or slightly curved, about 3 mm. thick, tipped with a long slender acute beak. *Seeds* oblong-ellipsoid, rounded at both ends, about 3 mm. long.

TROPICAL AFRICA: Senegal, *Perrottet* 229. Sierra Leone: Falaba, *Scott Elliott* 5220; near Lusenya, about 15 ft., with purplish flowers, *Scott Elliott* 4219. Gold Coast: Accra, *Krauss*; Lagos, *J. M. Dalziel* 1227; *Macgregor* 13. French Congo: Ft. Sibut, Krebedje, *Chevalier* 5675. Cameroons: Yaunde, *Zenker and Staudt* 501. Angola: Quiballa, *Monteiro*. Mossamedes, *Welwitsch* 1996. Between Huilla Mission Station and Chibia, *H. H. W. Pearson* 2699, 2724. Golungo Alto, *Welwitsch* 1994. Mossamedes. Eastern Sudan: *Schweinfurth* 1046, 1869; *Kotschy* 539; *Brown* 1075. Abyssinia: *Quartin-Dillon and Petit* 169. Colonia Eritrea: Beni-Amer, *Pappi* 161. British East Africa: Mombassa, in damp meadows, *Hildebrandt* 1990; *Kassner* 471; *Usambara, C. Holst* 3212. Nyasa Highlands, *Stolz* 267. Nyasaland: *Johnson* 361; *Buchanan* 910. Rhodesia: Matopos, *Allen* 731. Ngamiland: Kwebé Hills, swampy places, *Lugard* 161, 161a, 168. Portuguese East Africa: Lourenço Marques, *Schlechter* 11585; *Wilms* 422.

SOUTH AFRICA: Transvaal: Shilovane, *Junod* 1131. Crocodile Poort, shrub 8 ft., *Galpin* 1074a. Aapires River, north of Pretoria, *A. Haager* in *Herb. Conrath* 1185. Barberton Division: Nelspruit, *Rogers* 23858. Natal, *Drege, Sutherland*; *Gerrard* 32; *Keun, Wood* 287, 1593; *Gerrard and McKen*. 734.

A common weed in most tropical countries.

16. *Sesbania arabica*, *Hochst. ex Steud.*, *Nom. ed.* II, ii, 572 (1840).

Stems glabrous, rather slender, finely grooved. *Leaves* up to 15 cm. long, much longer than the inflorescence; rachis smooth; leaflets up to thirty pairs, distinctly stalked, linear-oblong, rounded and minutely mucronate at the apex, obliquely one-sided at the base, about 1.5 cm. long and 2.5-3 mm. broad, glaucous-green; stipules lanceolate, with a long linear acuminate apex, slightly pubescent on the margin, sub-persistent; stipels very minute, like little points. *Inflorescence* mostly about two or three flowered, slender; peduncle glabrous or nearly so; bracteoles paired at the base of the calyx, narrowly lanceolate, with membranous margins. *Flowers* apparently yellow; the vexillum finely mottled with purple. *Calyx* broadly campanulate from a shortly turbinate base, about 5 mm. long, slightly pubescent within the margins of the acute triangular teeth. *Vexillum* with two linear sessile appendages on the claw. *Fruits* erect, nearly straight, up to about 25 cm. long, strongly constricted between the seeds, segments about 6 mm. long. *Seeds* oblong-ellipsoid, rounded at both ends, 4.5 mm. long, dark brown, slightly shining.

TROPICAL AFRICA: Northern Nigeria: Benne Valley, Muri Province, Nov., *P. H. Lamb* 69. Eastern Sudan: S. Kordofan, *Pfund* 132, 308; *Kotschy* 47. Blue Nile, near "Camp Ardeba," *Muriel* 65. Mestra Zeraf Wood Station, *Brown* 1737. Colonia Eritrea: Beni-Amer, *Pappi* 166.

Occurs also in Southern Arabia. Arabic name, "Tawri" (*Muriel*).

17. *Sesbania sphaerocarpa* *Welw.* *Apont.* 590, No. 82 (1858): *Hiern.* in *Cat. Afr. Pl. Welw.*, i, 232, partly. *Sesbania sphaerosperma*, *Welw. ex Baker* in *Oliv. Fl. Trop. Afr.*, ii, 135 (1871).

Herb, rather woody at the base, about 30 cm. high or more; stems smooth or sparingly aculeata. *Leaves* up to 15 cm. long; rachis slightly aculeate below; leaflets up to about twenty pairs, elongate-oblong, rounded and minutely mucronate at the apex, unequal-sided at the base, 1.5–2.5 cm. long, about 5 mm. broad, pale glaucous-green, glabrous; stipules linear-subulate, acutely acuminate, about 8 mm. long, slightly hairy on the margins, soon falling off. *Inflorescence* very small, 2–3-flowered, about 1.5 cm. long; peduncle slender, slightly hairy towards the base; pedicels very slender, about 5 mm. long, glabrous; bracts and bracteoles caducous. *Calyx* widely campanulate, acutely turbinate at the base, with five short acute lobes. *Vexillum* broadly orbicular, with two sessile linear appendages on the claw. *Fruits* about 16 cm. long, 3 mm. broad, linear, acutely acuminate, with an undulate wing on the back of the valves. *Seeds* nearly rounded, small.

TROPICAL AFRICA: Angola: Loanda District, *J. Gossweiler* 253, 1479; Loanda, *Welwitsch* 1991, 1993.

18. *Sesbania microphylla*, *Harms* in *Kunene-Samb. Exped.*, 260 (1903).

An erect branching herb, up to 2 m. high; branches ribbed, glabrous. *Leaves* 2–12 cm. long; petiole 0.2–1 cm. long, glabrous; rachis flattened on the upper surface, glabrous, leaflets 5–30-jugate, subsessile, 2–7 mm. long, 0.5–2.5 mm. broad, oblong or elliptic, obtuse, shortly mucronate, slightly unequal at the base, punctate above with minute black dots, glabrous, with the mid-rib distinct beneath. *Stipules* 2.5–3 mm. long, setaceous, deciduous, or persistent. *Inflorescence* axillary, racemose, three or four flowered, 2–6 cm. long, longer or shorter than the subtending leaf; base of peduncle glabrous; bracts deciduous, not seen; bracteoles 2.5 mm. long, setaceous, persistent in mature flowers; pedicels 1–2 cm. long, glabrous. *Calyx-tube* 3 mm. long, 3 mm. in diameter, campanulate, with five broad brown bands; teeth 1 mm. long, ovate, with brown acuminate tips. *Vexillum* yellow, mottled with numerous fine purple spots on the back, 1.5 cm. long, 1.1 cm. broad, ovate-oblong, rounded above, with two adnate linear-appendages within a short claw 1.5 cm. long. *Alae* 1.6 cm. long, oblong, obtuse, with a linear claw 2.5 cm. long, and a downward projecting tooth from the base of the lamina. *Carina* 1.65 cm. long, 5 mm. broad, almost plano-convex in outline, with a linear claw 6 mm. long, and a downward projecting tooth from the lamina. *Staminal sheath* 1 cm. long, 2 mm. broad at the base, with the free portion of the filaments arcuate; free stamen slightly bent at the base; anthers 0.75 mm. long, ovate. *Ovary* 1.3 cm. long, sub-compressed, with two faint longitudinal bands; style 4 mm. long, arcuate; stigma capitate. *Fruit* 11.5–12.5 cm. long, 2–2.5 mm. broad, linear, acutely acuminate, laterally compressed, moniliform. *Seeds* dark brown, 3 mm. long, 1.5 mm. broad, oblong, rounded at both ends, glabrous.

TROPICAL AFRICA: Angola: Mouth of the Longa River, 3480 ft., 22nd December, 1899, *H. Baum* 569. Eastern Sudan: Bahr. el Ghazal, flowers yellow mottled with black, *A. F. Brown*. Rhodesia: Lomagundi, Darwendale on pierite formation, in vleis, April, *Eyles*, 695; Bulawayo, *J. S. MacDonald* in *Herb. Dept. Agric.* 4942.

19. *Sesbania mossambicensis*, *Klotzsch* in *Peters, Reise Mossamb. Bot.* 45 (1861). *Sesbania macowaniana* *Schinz*, in *Verh. Bot. Ver. Brand*, XXX, 165 (1888).

A weak sparingly branched herb about 1 m. high. *Branches* striate, glabrous. *Leaves* 1.5–11 cm. long; petiole 2–7 mm. long, glabrous; rachis flattened or channelled above, glabrous, rarely sparsely covered with hairs; leaflets 7–33-jugate, subsessile, oblong or oblong-linear, rounded at the apex, shortly apiculate, unequal at the base, glabrous, 0.3–1 cm. long, punctate with minute black dots on the upper surface, with the mid-rib distinct beneath; stipules usually persistent, 2–5 mm. long, ovate or ovate-lanceolate, acuminate, acute. *Inflorescence* an axillary raceme, 1–7 cm. long, 3–8-flowered, longer or shorter than the subtending leaf; peduncles glabrous; bracts generally persistent, 2 mm. long, linear; bracteoles persistent, 1.5 mm. long, linear; pedicels 3–7 mm. long,

glabrous. *Calyx-tube* 3-4 mm. long, 2-3 mm. in diameter, campanulate, glabrous, with five dark-brown bands; teeth 1-1.5 mm. long, acuminate from a triangular base, acute, with black tips, usually with cottony hairs on the margins. *Vexillum* striped with dark veins, but not mottled, 0.8-1.1 cm. long, 0.8-1.1 cm. broad, subrotund, obovate or transversely oblong, rounded and retuse at the apex, with a claw 2 mm. long. *Alae* 0.8-1.2 cm. long, 3-4 mm. broad, obovate or obovate-oblong, obtuse, with a linear claw 1.5-3 mm. long, and a short downward projecting tooth from the lamina. *Carina* 0.9-1.2 cm. long, 3-4 mm. broad, almost plano-convex in outline, obtuse, with a linear claw 5-7 mm. long, and a downward projecting claw from the lamina. *Staminal sheath* 0.7-1 cm. long, 1-3 mm. broad at the base, with the free portions of the filaments arcuate; free stamen dilated at the base. *Ovary* 0.7-1.1 cm. long, subcompressed, with two faint longitudinal bands; style 3-4 mm. long, arcuate, glabrous; stigma capitate. *Fruit* 4-6 cm. long, moniliform, long-acuminate. *Seeds* dark reddish-brown, 2.5 mm. long, 1.25 mm. broad, oblong, rounded at each end, glabrous.

TROPICAL AFRICA: Maniniam: On the Tubbo River, *G. Schweinfurth* 3674. Portuguese East Africa: Cabaceira Peninsula and near Quillimane, *Peters* (type). Congo: Elisabethville, *Rogers* 10872; without locality, *Rogers* 13641.

SOUTH AFRICA: Herbert Div., Biddulphis, nine miles north of Orange River at Hopetown, February, *Bolus* 1835; Gariep River, *Pappe*: Transvaal: Makapansberge, Streydpoort, *A. Rehmann* 5543. Pretoria Div., near the Aapies River, 4500 ft., January, *Schlechter* 4179; *Miss Leendertz* 3205; Pretoria, March, *Mrs. Pott in Herb. Transvaal Mus.* 11333; *Miss Leendertz* 4413; Hatherley, near Pretoria, *Rogers* 23943; Rustenburg Div., Rustenburg, January, *Playford in Govt. Herb.* 17268; Crocodile River, March, *Burtt-Davy*, 9297; Zeerust Div., Zeerust, *Rogers* 22672; Potchefstroom Div., Potchefstroom, *Miss Leendertz* 3205; Pietersburg Div., grassy plains near Pietersburg, 4000 ft., February, *Bolus* 10879. Bechuanaland: Masupa River, *E. Holub*. S.W. Protectorate: Klein Namutoni, *J. W. F. Breijer in Herb. Transvaal Mus.* 20618.

§ DAUBENTONIA.

20. *Sesbania tetraptera*, *Hochst. ex Baker* in *Oliv. Fl. Trop. Afr.*, ii, 136 (1871).

Stems ridged, otherwise glabrous, about 4 mm. thick. *Leaves* up to 12 cm. long, nearly twice as long as the inflorescence; rachis sparingly pubescent when young, soon becoming quite glabrous or with a few small prickles on the lower side towards the base; leaflets about fifteen pairs, shortly stalked, narrowly oblong, unequal-sided at the base, truncate-rounded and sharply mucronate at the apex, 1.2-2 cm. long, 3-4.5 mm. broad, glabrous, spotted with minute black dots on the upper side, with no visible nerves or veins; stipules subsistent, subulate-lanceolate, very acute, 5-6 mm. long, very slightly pubescent inside; stipels very minute and subulate. *Inflorescence* a simple raceme, few (up to seven) flowered; peduncle up to 6 cm. long, thickened at the base and there closely beset with subulate-triangular prickles; bracts linear-subulate, about 3 mm. long, persistent; pedicels about 7 mm. long, filiform, with a pair of linear deciduous bracteoles at the apex about one-third the length of the calyx. *Calyx* campanulate, turbinate at the base, 5 mm. long, with five short triangular lobes, the latter minutely pubescent inside the margin. *Vexillum* broadly orbicular, 1.2 cm. long, cuneate into a short claw about 2 mm. long, with two linear sessile ridge-like appendages about 2 mm. long near the base. *Alae* oblong; limb 8.5 mm. long; claw bent, 5 mm. long. *Carina* broadly semilunar; limb 8 mm. long, with an acutely triangular tooth on the upper side; claw curved, 5 mm. long. *Staminal sheath* 0.8 cm. long, free part of filaments 3-4 mm. long; anthers 0.75 mm. long. *Ovary* 8 mm. long, four-winged; style bent at right angles, 3 mm. long, puberulous towards the top. *Fruits* much longer than the leaves, 15 cm. long, 1 cm. broad, broadly four-winged, stipitate, for about 1 cm. at the base, acutely acuminate at the apex. *Seeds* reddish-brown, oblong-elliptic, broadly rounded at both ends, 6 mm. long, with a small circular hilum in the middle.

TROPICAL AFRICA: Nubia: In dry limestone of Arasch-Cool, 7th October, 1839, Kotschy 131 (H.K.). "Upper Egypt and Cordofan," without definite locality, 1837-38, Kotschy 83 (H.K.).

21. *Sesbania Kirkii* * n. sp.

Stems slightly ridged or nearly smooth, not prickly, rather stout, about 5-6 mm. thick. *Leaves* up to 30 cm. long, more than twice as long as the inflorescence; rachis sparingly pilose when very young, soon glabrous or with one or two small prickles on the lower side towards the base; leaflets up to about 35 pairs, shortly stalked, elongate-oblong or linear-oblong, slightly unequal-sided at the base, rounded and mucronate at the apex, 2.5-3.5 cm. long, about 5-6 mm. broad, spotted with minute black dots on the upper surface, otherwise glabrous, with several faint lateral nerves; stipules persistent, lanceolate or linear-lanceolate, acutely acuminate, 1 cm. long, pubescent inside; stipels distinct, subulate, about 1 mm. long. *Inflorescence* several flowered, racemose, slender, up to 14 cm. long; peduncle slender, broadened towards the base and there provided with numerous subulate spreading prickles; bracts persistent, longer than the pedicels, linear-lanceolate, very acute, 5-6 mm. long, sparingly ciliate; pedicels about 4 mm. long, slender, glabrous, with a pair of persistent bracteoles a little more than half as long as the calyx. *Calyx* campanulate, turbinate at the base, 4 mm. long, with five triangular lobes about 1 mm. long, the latter minutely ciliate. *Petals* as in *S. tetraptera* Hochst. *Fruits* much shorter than the leaves, about 15 cm. long and nearly 1.5 cm. broad, stipitate for about 1 cm. at the base, very acutely beaked, acuminate, broadly four-winged. *Ripe seeds* not seen.

TROPICAL AFRICA: Zambezi Basin: Tette, fls., February, 1859, Dr. J. Kirk (H.K.); near Sena, fr., June, 1859, Dr. J. Kirk (H.K.). Lower Shire, elephant marsh, especially on the banks of the river, fls. and fr., February, 1887, L. Scott (H.K.).

22. *Sesbania hamata*,† n. sp.

* *Sesbania Kirkii*, Phillips et Hutchinson, sp. nov.; affinis *S. tetrapterae*, Hochst., sed foliis, fructus brevioribus foliolis numerosissimis majoribus inflorescentiis multifloris differt.

Caulis leviter costatus vel fere teretes, haud aculeati, tatis robusti, circiter 5-6 mm. crassi. *Folia* usque ad 30 cm. longa, inflorescentiae ultra duplo-longiora; rachis primum parce pilosus, mox glaber vel inferne aculeis paucis instructus; foliola usque ad 35-jugata, breviter petiolulata, elongato-oblonga vel lineari-oblonga, basi leviter inaequilatera, apice rotundata et mucronata, 2.5-3.5 cm. longa, circiter 5-6 mm. lata, supra punctis nigris minutis maculata, ceterum glabra, nervis lateralibus subdistictis; stipulae persistentes, lanceolatae vel lineari-lanceolatae, acute acuminatae, 1 cm. longae, intra pubescentes; stipellae distinctae, subulatae, circiter 1 mm. longae. *Inflorescentia* pluriflora, racemosa, gracilis, usque ad 14 cm. longa; pedunculi graciles, basin versus latiores et aculeis numerosis subulatis patulis instructi; bractae persistentes, pedicellis longiores, lineari-lanceolatae, acutissimae, 5-6 mm. longae, parce ciliatae; pedicelli circiter 4 mm. longi, graciles, glabri, apice bracteolis calyce dimidio brevioribus instructis. *Calyx* campanulatus, basi turbinatus, 4 mm. longus, lobis triangularibus 5 circiter 1 mm. longis minute ciliolatis. *Petala* ut in *S. tetrapterae*, Hochst. (supra). *Fructus* foliis multo brevior, circiter 15 cm. longus et fere 1.5 cm. latus, basi 1.5 cm. stipitatus, apice acutissime rostrato-acuminatus, late 4-alatus. *Semina* matura non visa.

† *Sesbania hamata*, Phillips et Hutchinson, sp. nov.; affinis *S. tetraptera* Hochst., sed caulibus aculeis hamatis ornatis inflorescentiis plurifloris differt.

Caulis longitudinaliter costatus, aculeis hamatis. *Folia* 9-18 cm. longa; rachis interdum aculeis paucis ornatus; foliola 14-20-jugata, subsessilia, oblonga vel lineari-oblonga, apice rotundata, distincte mucronata, 1.2-2.7 cm. longa, 2.5-6 mm. lata, supra dilute viridia, glabra; stipulae persistentes, lineis rubro-brunneis striatae, 0.6-1 cm. longae, oblique lanceolatae, acuminatae, glabrae. *Inflorescentia* axillaris, racemosa, 9-11 cm. longa, 6-9-flora, folio subtendente circiter aequilonga; pedunculi inferne aculeis numerosis hamatis instructi; bractae persistentes, circiter 5 mm. longae, lanceolatae, acuminatae, acutae, leviter ciliatae; bracteolae bracteis similes sed breviores; pedicelli 0.9-1.5 cm. longi, glabri *Calycis* *tubus* 3 mm. longus, 4 mm. latus, campanulatus, glaber, dentibus 1 mm. longis ovatis acutis glabris. *Vexillum* maculatum, 1-2 cm. longum, 1.3 cm. latum, transverse oblongum, ungue 3 mm. longo; alae 1.25-1.3 cm. longae, 4 mm. latae, oblongo-ovatae, superne rotundatae, ungue lineari circiter 3 mm. longo et dente recurvato; carina 1.2 cm. longa, obtusa, ungue 5 mm. longo. *Tubus staminialis* 3 mm. longus, 3 mm. latus, filamentis superne arcuatis liberis; stamen liberum 8 mm. longum. *Ovarium* subsessile, 8 mm. longum, 4-alatum, glabrum; stylus 3 mm. longus, ab ovario sub angulo 45° divergens, pubescens, stigmatibus capitato. *Fructus* immaturi usque ad 9.5 cm. longi, 4.5 mm. lati, lineares, 4-alati, stylo persistente coronati.

Stem with longitudinal ridges, covered with hook-like prickles. *Leaves* 9-18 cm. long; rachis sometimes with a few hook-like prickles on the lower half; leaflets 14-20 pairs, subsessile, oblong or linear-oblong, rounded at the apex, distinctly mucronate, 1.2-2.7 cm. long, 2.5-6 mm. broad, pale green above, glabrous, with the mid-rib distinct beneath; stipules persistent, striped with reddish-brown marks, 0.5-1 cm. long, obliquely lanceolate, acuminate, glabrous. *Inflorescence* an axillary raceme 9-11 cm. long, 6-9-flowered, about as long as the subtending leaf, the lower half of the peduncle covered with numerous hook-like prickles; bracts persistent, about 5 mm. long, lanceolate, acuminate, acute, slightly ciliate; bracteoles similar to the bracts but a little shorter; pedicels 0.9-1.5 cm. long, glabrous. *Calyx-tube* 3 mm. long, 4 mm. broad, campanulate, glabrous; teeth 1 mm. long, ovate, acute, glabrous. *Vexillum* spotted, 1.2 cm. long, 1.3 cm. broad, transversely oblong, with a claw 3 mm. long, widening from the base upwards. *Alae* 1.25-1.3 cm. long, 4 mm. broad, oblong-ovate, rounded above, with a linear claw about 3 mm. long and a downward projecting tooth from the limb forming a loop with the claw. *Carina* 1.2 cm. long, obtuse, with a linear claw 5 mm. long, and with an acute tooth projecting at right angles to the limb and forming a horseshoe-shaped loop with the claw. *Staminal sheath* 4 mm. long, 3 mm. broad, with the upper half of the free filaments arcuate; free stamen 8 mm. long, dilated at the base above a short bent claw. *Ovary* subsessile, 8 mm. long, four-winged, glabrous; style 3 mm. long, bent almost at right angles to the ovary, pubescent; stigma capitate. *Young fruits* up to 9.5 cm. long, 4.5 mm. broad, linear, four-winged, tipped with the persistent style.

SOUTH AFRICA: Transvaal: Moist places near the Brak River, 2800 ft., 9th March, 1894, *Schlechter* 4620.

We have seen only one specimen of this apparently distinct species—in the Bolus Herbarium, Capetown. It extends the distribution of the group *Tetrapterae*, formerly only known by the solitary species (*S. tetraptera*) from the Nile districts, through the Zambezi Basin (*S. Kirkii* and *S. Rogersii*) into the sub-tropical Transvaal.

23. *Sesbania Rogersii* * n. sp.

Branched or unbranched plants 30-69 cm. high. *Stem* ribbed, covered with prickles, otherwise glabrous. *Leaves* 1.5-6 cm. long; rachis 3-4 cm. long, subterete; leaflets less than ten pairs, subsessile, 0.5-1.3 cm. long, 2-4 mm. broad, oblong, obtuse, shortly and bluntly mucronate, punctate above with minute black dots, otherwise glabrous, with the mid-rib distinct below; stipules setaceous, 4.5 long. *Inflorescence* an axillary raceme, 2-5 cm. long, 1-4-flowered, longer or shorter than the subtending leaf; peduncle with a few short prickles at the base; bracts 5 mm. long, lanceolate, acuminate, deciduous; bracteoles similar, soon falling off; pedicels 0.5-1.2 cm. long, glabrous. *Calyx-tube* 4 mm. long, campanulate, glabrous; teeth about 1 mm. long, ovate, acute, minutely ciliate. *Vexillum* mottled, 1-1.3 cm. long, 7-8 mm. broad, obovate, with a linear claw

* *Sesbania Rogersii*, *Phillips et Hutchinson*, sp. nov.; affinis *S. hamatae*, *Phillips et Hutchinson*, sed foliis, paucijugis inflorescentiis 1-3-floris, tubo staminorum filamentorum partibus liberis longiore differt.

Plantae simplices vel ramosae usque ad 30-70 cm. altae; caulis costatus, aculeis ornatus, ceterum glaber. *Folia* 1.5-6 cm. longa; rachis 3-4 cm. longus, subteres; foliola 5-8-juga, subsessilia, 0.5-1.3 cm. longae, 2-4 mm. lata, oblonga, obtusa, breviter et obtuse mucronata, supra minute nigro-lepidotis; stipulae setaceae, 4-5 cm. longae. *Inflorescentia* axillaris, racemosa, 2-6 cm. longa, 1-4-flora, folio subtendente longior vel brevior; pedunculi basin versus aculeis brevibus instructi; bractae 5 mm. longae, lanceolatae, acuminatae, deciduae; bracteolis similibus mox deciduis; pedicelli 0.5-1.2 cm. longi, glabri. *Calycis tubus* 4 mm. longus, campanulatus, glaber, dentibus circiter 1 mm. longis ovatis acutis minute ciliatis. *Vexillum* irregulariter maculatum, 1-1.3 cm. longum, 7-8 mm. latum obovatum, ungue lineari 2-4 mm. longo apice bifidiculato. *Alae* 1.3-1.4 cm. longae, 3 mm. latae, oblongae, obtusae, ungue lineari 3 mm. longo dente reflexo; carina 1.3-1.5 cm. longa, obtusa, concava, ungue 5-8 mm. longo lineari. *Tubus staminalis* 1-1.3 cm. longus basi 1.5-2 mm. latus, antheris 0.5 mm. longis, oblongis. *Ovarium* breviter stipitatum, 1.1-1.2 cm. longum, 4-alatum; stylus 3-3.5 mm. longus, arcuatus, apicem versus pubescens, stigmatibus parvis. *Fructus* 9-15 cm. longi, 1.1-1.2 cm. lati, lineares, acuminati, late 4-alati, medio pallidiores. *Semina* atrorubrobrunnea, 5 mm. longa, 2.5 mm. lata oblonga, utrinque fere truncata, glabra.

2-4 mm. long and two raised ridges above the claw. *Alae* 1.3-1.4 cm. long, 3 mm. broad, oblong, obtuse, with a linear claw 3 mm. long and a short downward projecting tooth. *Carina* 1.3-1.5 cm. long, obtuse, concave, with a linear claw 5-8 mm. long, and a projecting tooth almost at right angles to the limb. *Staminal sheath* 1-1.3 cm. long, 1.5-2 mm. broad at the base; free portion of the filaments shorter than the tube, arcuate; anthers 0.5 mm. long, oblong; free stamen bent near the base, but not swollen. *Ovary* shortly stipitate, 1.1-1.2 cm. long, four-winged; style 3-3.5 mm. long, arcuate, pubescent near the apex; stigma small. *Fruit* 9-15 cm. long, 1.1-1.2 cm. broad, linear, acuminate, broadly four-winged, with a lighter coloured band in the middle of the valves. *Seeds* dark reddish-brown, 5 mm. long, 2.5 mm. broad, oblong, almost truncate at both ends, glabrous.

TROPICAL AFRICA: Rhodesia: Victoria Falls, April, *Flanagan* 3109; North-West Rhodesia, without locality, *Rogers* 8747.

This is a very peculiar species with exactly the appearance of *S. aculeata* in regard to its vegetative and floral characters, but with fruits broadly four-winged as in *S. tetraptera*. It apparently provides a link between the two sections of the genus.



THE NATAL SPECIES OF THE SAPINDACEAE.

By E. P. PHILLIPS, M.A., D.Sc., F.L.S., Botanist in Charge of the
National Herbarium, Pretoria.'

DURING a short stay in Durban in January, 1919, I saw for the first time, in their native state, species of *Sapindaceae* growing in the Stellabush. This led me to make an examination of the material in the Natal Herbarium, and at the same time I took the opportunity of describing the species. To make the records more complete than it was possible to do from the material in the Natal Herbarium alone, all the Natal specimens in the Bolus Herbarium and Cape Government Herbarium at Capetown and the National Herbarium, Pretoria, were examined. This has added considerably to the Natal localities in which the species occur.

Twelve genera of the order *Sapindaceae* have been recorded from South Africa, and of these nine occur in Natal. Wood, in his "Handbook to the Natal Flora" (1907), cites sixteen Natal species, but he eventually dropped *Schmidelia natalensis* in his "Revised List of the Natal Flora." Of the fifteen species mentioned by Wood, only ten are represented in the Natal Herbarium, and of those unrepresented I have not seen specimens from Natal in any other herbarium.

The examination of the material has brought to light an undescribed *Bersama* which occurs in Natal, and I am of opinion that *Bersama tysoniana* also occurs there, though I have seen no Natal specimen.

KEY TO THE GENERA.

- Stamens eccentric. Fruit bladdery. A twiner..... *Cardiospermum*.
- Stamens not eccentric. Fruit membranous, fleshy, coriaceous, or winged.
 - Trees or shrubs.
 - Whole plant densely villous with stellate hairs; leaflets deeply incised..... *Melianthus*.
 - Plants most glabrous; leaves or leaflets never deeply incised.
 - Leaves broadly ovate, palmately lobed; flowers scarlet..... *Greyia*.
 - Leaves or leaflets lanceolate or elliptic; flowers greenish or whitish.
 - Leaves simple.
 - Fruit almost globose, coriaceous..... *Allophylus*.
 - Fruit flattened with membranous wings..... *Dodonaea*.
 - Leaves compound.
 - Leaflets unequal-sided; rachis with broad or narrow wings.
 - Young branches, leaf rachis, and inflorescence densely tomentose..... *Hippobromus*.
 - Plant glabrous or almost so..... *Pteroxylon*.
 - Leaflets not unequal-sided; rachis not winged, or only winged in the uppermost segment.
 - Leaves trifoliate; sepals glabrous, pedicels 1 mm. long..... *Allophylus*.
 - Leaves pinnate, if trifoliate then pedicels 1 cm. long; sepals hairy.
 - Inflorescence a panicle up to 43 cm. long.... *Sapindus*.
 - Inflorescence a raceme up to 17 cm. long.... *Bersama*.

CARDIOSPERMUM LINN.

C. halicacabum Linn.

A herbaceous climber. *Stems* deeply five-furrowed with prominent ridges, scantily pubescent. *Leaves* 3-6 cm. long, trifoliate, each leaflet of the first order digitately divided into three leaflets of the second order; petiole 1.5-1.8 cm. long, furrowed, pilose; petiole of leaflets 8-11 cm. long, similar to the petiole of the leaf; ultimate leaf-segments 1.5-2.5 cm. long, 0.5-1.3 cm. broad, ovate or ovate-lanceolate, acute, hirsute above, glabrous beneath except on the veins, with the margins deeply cut into 2-3 lobes. *Inflorescence* a compound umbel with three principal peduncles, axillary; primary peduncle 3-7.2 cm. long, four-angled, pilose, and bearing at the apex two spirally-coiled tendrils; secondary peduncles 8-1.7 cm. long, scantily pubescent. *Sepals* unequal; the smaller 2 mm. long, 2 mm. broad, ovate, rounded above, glabrous; the larger 5 mm. long, 2-5 mm. broad, obovate, obtuse. *Petals* similar to the larger sepals. *Stamens* eccentric; staminal tube 1-5 mm. long; free portion of filaments 2.5-3 mm. long, linear, glabrous, sometimes ciliated; anthers 0.75 mm. long, oblong. *Ovary* 3 mm. long, eccentric, consisting of three carpels with a broad dorsal wing and a narrow ventral wing; style none; stigmas three, ovate, concave. *Torus* between the stamens and ovary produced into two blunt lobes 5 mm. long. *Fruit* 2-3-celled bladderly capsule with a single seed in each cell; the valves not veined and pubescent with bulbous base hairs. *Seeds* black, 4 mm. long, 3.5 mm. broad, somewhat globose, glabrous. *Sonder in Harv. and Sond., Fl. Cap., 1, 237, Bot. Mag., t. 1049.*

Natal: Verulam, March, *Wood* 483, and in *Colonial Herb.* 6795, without locality, *Sanderson, Gueinzuis.*

DISTRIBUTION: Rhodesia, Pondoland, Transvaal, Bechuanaland Protectorate.

MELIANTHUS LINN.

M. villosus Bolus.

A herbaceous shrub. All parts densely covered with stellate hairs. *Branches* tomentose. *Leaves* petioled, compound, paripinnate, 5-7-jugate, 8-19 cm. long; petiole 3.2-5.5 cm. long, densely villous; leaflets 2.5-7.5 cm. long, 6-3 cm. broad, lanceolate, subacute, slightly narrowed at the base, densely villous, with the margins deeply and acutely serrate; stipules 2.2 cm. long, 4 mm. broad, ovate-lanceolate, long acuminate, acute, densely villous. *Inflorescence* a terminal raceme 28 cm. long. *Peduncle* 12 cm. long, densely villous. *Bracts* 2.3 cm. long, 1.2 cm. broad, ovate, long acuminate, pubescent. *Pedicels* 2 cm. long, terete, villous. *Sepals* unequal; lateral sepals 2.6 cm. long, 1.3 cm. broad, elliptic, obtuse, tomentose within and without; anterior sepal 1.6 cm. long, 1 cm. broad, boat-shaped, ovate in outline, long acuminate, densely tomentose. *Petals* 1.9 cm. long, 2-3 mm. broad, linear, three-lobed above, densely pilose on the upper half, with the middle lobe tongue-shaped; the lateral lobes smaller, sometimes rudimentary. *Disc* in the form of flat fleshy glands 4-5 mm. long, 2 mm. broad, truncate at the apex. *Filaments* 6-9 mm. long, subterete, pilose; anthers 4.5 mm. long, 2 mm. broad, oblong. *Ovary* 4 mm. long, pyramidal in outline, four-angled, glabrous, four-celled, each cell containing many ovules; style 5 mm. long, subterete, somewhat falcate, narrowing to a blunt point, pubescent; stigma simple.

Natal: Rocky Hill, South Downs, 4000-5000 ft., December, *Wood* 1008, 4376, and in *Natal Govt. Herb.* 5133.

DISTRIBUTION: Elands River Valley, near Mont-aux-Sources.

M. insignis, O. Kuntze. I have not seen specimens of this species, but from the short description given by Kuntze (*Rev. Gen.* III, 43) it may possibly be the same as *M. villosus* Bolus.

GREYIA HOOK. AND HARV.

G. Sutherlandii Hook. and Harv.

A moderately sized tree (*Wood*). *Branches* glabrous. *Leaves* simple, petioled, 4–10 cm. long; petiole 1–4 cm. long, somewhat compressed, glabrous, dilated and amplexicaul at the base; lamina 3–7 cm. long, 2–8 cm. broad, ovate, obtuse, somewhat cordate at the base, glabrous, 8–9-lobed with the lobes coarsely toothed, palmatinerved with distinct veins. *Inflorescence* a dense raceme, 7–11 cm. long, 3·5–5 cm. broad. *Bracts* 1 cm. long, 5 cm. broad, lanceolate, obtuse. *Pedicels* 1·5 cm. long, terete, glabrous. *Sepals* 5 mm. long, 3 mm. broad, ovate, obtuse, glabrous, connate at the base. *Petals* red, 1–1·5 cm. long, 5·5 mm. broad, oblong, concave, rounded at the apex, glabrous, shortly ciliated. *Disc* basin-shaped, with a rim 2 mm. deep and produced into ten processes 2 mm. long, each bearing a flattened anther-like structure at the apex. *Filaments* reddish, 9·2 cm. long, terete, glabrous; anthers 3 mm. long, oblong. *Ovary* 7 mm. long, 3·5 mm. broad, ovoid, subapocarpous, five-celled, each cell containing numerous ovules; style 5 mm. long, gradually narrowed above, minutely five-toothed at the apex, glabrous. *Harv. Thes. Cap. t. 1, Harv. in Harv. and Sond. Fl. Cap. II, 309; Wood, Natal Plants t. 373.*

Natal: Polela, June, *Fernando*, 83; and in *Natal Govt. Herb.* 10470; Ixopo, September, *Fannin in Natal Herb.* 16241; Inanda, 2000 ft., *Wood in Natal Govt. Herb.* 95; Van Reenens Pass, 5800 ft., November, *Wood* 4691; without precise locality, *Sutherland*.

DISTRIBUTION: Orange Free State, Tembuland, Stutterheim, Transvaal.

ALLOPHYLUS LINN.

Leaves simple..... *monophylla*.

Leaves trifoliate.

Leaflets lanceolate; petiole glabrous; stigmas linear..... *erosa*.

Leaflets mostly elliptic; petioles pubescent; stigmas ovate..... *africana*.

Leaflets obovate or oblanceolate; petioles pubescent; stigmas linear... *decipiens*.

A. monophyllus Radkl.

A large spreading shrub or small tree, 10–30 ft. high and up to 12 in. stem diameter (ex *Sim*). *Branches* with minute mealy pubescence and greyish bark. *Leaves* simple, petioled, 2·5–2·3 cm. long, 2·5–8 cm. broad, elliptic, lanceolate, or ovate, acuminate, subobtuse, slightly narrowed or more rarely subrotund at the base, glabrous above and beneath, sometimes with tufts of hairs in the angles of the lateral veins beneath, darker above, with coarsely serrated margins; petiole 1·3–5 cm. long, terete, minutely pubescent or almost glabrous. *Inflorescence* a raceme, 4–17 cm. long, sometimes the flowers in groups of two on short secondary peduncles. *Peduncles* 1·5–5 cm. long, terete, minutely pubescent or glabrous. *Pedicels* about 1·5 cm. long. *Sepals* unequal, the smaller 1·5 mm. long, 1 mm. broad, elliptic, rounded above, deeply concave, glabrous; the larger 2 mm. long, 1·75 mm. broad, subrotund, concave, glabrous. *Petals* 2 mm. long, spatulate, densely bearded within. *Filaments* 2·5 mm. long, linear, pilose; anthers 4 mm. long, oblong. *Ovary* 1 mm. long, 2 mm. broad, didymous, pubescent; style 0·75 mm. long, terete, glabrous; stigmas 0·75 mm. long, linear, divergent. *Fruit* 9 mm. long, 7 mm. in the diameter, obovate in outline, slightly narrowed at the base, glabrous. *Schmidelia monophylla*, Presl. *Fl. Cap. I, 239. Sim, Forests and Forest Floras*, 170, Pl. XXXII, Fig. 3, *S. dregeana*, Sond. in *Harv. and Sond. Fl. Cap. I, 239.*

Without precise locality, Gerrard and McKen. 525, and in *Natal Govt. Herb.* 726.

Natal: Inanda, April, *Wood*, 481, and in *Colonial Herb.*, 5678, Ifafa, August (in fruit), *Lansdell in Natal Herb.* 16126; Imont, *Wood*, 2468; edge of bush at Spring Grange, c., 800 ft., April, *Wood*, in *Natal Govt. Herb.*, 11248; without locality, *Pappe*.

Zululand: Qudeni Forest, 6000 ft., February, *Davis* 118, and in *Natal Govt. Herb.*, 8700.

DISTRIBUTION: Komgha, Kentani, East Pondoland.

A. erosus Radkl.

A shrub 4-12 ft. high (ex *Sim*). *Branches* glabrous with wrinkled greyish bark. *Leaves* petioled, trifoliate, 6-12 cm. long, petiole 2-3 cm. long, channelled above, glabrous; leaflets sessile, 3-8.5 cm. long, 9-2.6 cm. broad, lanceolate, obtuse, narrowed at the base, glabrous, minutely punctate beneath, with the margins coarsely and bluntly serrate. *Inflorescence* an axillary panicle, 4-7 cm. long, the flowers ultimately cymose in groups of 2-4; peduncle grooved, minutely pubescent or almost glabrous. *Pedicels* 1 mm. long, glabrous. *Sepals* unequal, 1 mm. long, 0.75 mm. broad, obovate, rounded above, concave, glabrous. *Petals* white, 1 mm. long, 0.75 mm. broad, obovate, subcrenate and almost truncate at the apex, densely bearded within. *Filaments* (in young flowers) 0.75 mm. long, linear, pilose; anthers 5 mm. long, oblong in outline. *Ovary* 1 mm. long, 1.5 mm. broad, obovate, in outline, pubescent; style 1 mm. long, terete, glabrous; stigmas 1 mm. long, linear, obtuse, divergent. *Fruit* 5 mm. long, 5 mm. in diameter, globose, narrowed at the base, finely and scantily pubescent, at length glabrous. *Schmidelia erosa* Arn. *Sond. and Harv. Fl. Cap. I*, 239. *Sim, Forests and Forest Flora*, 169, Pl. XXXII, Fig. 1. *S. natalensis*, *Sonder in Harv. and Sond. Fl. Cap. I*, 239.

Natal: Back Beach Bush, near Durban, 20 ft., April, *Wood, in Natal Govt. Herb.*, 11249. Durban, April, *Wood* 898, 7841, 10964, and in *Colonial Herb.*, 5693. Catos Creek, May (in fruit), *Wood in Natal Govt. Herb.*, 2469. Umgeni, August (in fruit), *Native Collector in Natal Herb.*, 16218. Port Shepstone, April, *Brown* 422; without locality, *Saunders*; Berea, *Wood*.

Sim states: "Abundant along the coast, and especially on the coast dunes from the Fish River to Natal; not found inland. This species is a shrub 4-12 ft. high and often forms a large proportion of the sea scrub. It stands sea winds where most other shrubs fail, close to the sea. Its timber is not used and is seldom heavy enough for economic purposes."

DISTRIBUTION: East London.

A. melanocarpus Radkl.

A shrub or small tree reaching 20 ft. or more in height (ex *Wood*). *Branches* minutely pubescent, at length becoming glabrous, with greyish bark. *Leaves* petioled, trifoliate (4.5), 22 cm. long; petiole 2.2-8.5 cm. long, channelled above, minutely and densely pubescent; leaflets darker above, paler beneath, shortly petioled, 2.5-12.5 cm. long, 1.2-4.5 cm. broad, elliptic, elliptic-lanceolate, ovate or ovate-lanceolate, acuminate, obtuse, rounded or more usually narrowed at the base, pubescent above and beneath, especially on the veins, sometimes almost glabrous, with a prominent mid-rib beneath and 7-11 lateral veins forming an acute angle with the mid-rib and distinct above, sometimes with distinct tufts of hairs in the angles between the mid-rib and the lateral veins on the under surface. *Inflorescence* an axillary panicle, 3.5-12 cm. long, with the flowers ultimately in cymose groups of 3-5 on short peduncles. *Peduncle* 2-4 cm. long, channelled, densely pubescent, branches of inflorescence similar to the peduncle. *Sepals* unequal, the smaller 1 mm. long, 0.75 mm. broad, more or less rotund, deeply concave, glabrous; the larger 1 mm. long, 1.5 mm. broad, ovate, rounded above, deeply concave, glabrous. *Petals* 1.5 mm. long, spatulate, bearded. *Filaments* 5 mm. long, linear, densely ciliated; anthers 5 mm. long, somewhat obovate in outline. *Ovary* 7.5 mm. long, 1.5 mm. broad, obovate in outline, pubescent, two locular, with a single ovule in each loculus; style 1 mm. long, terete; stigmas, two, reflexed, 0.75 mm. long, ovate, acuminate, acute. In male flowers filaments 1.1-5 mm. long, linear, ciliated; anthers, 0.5 mm. long, subglobose. *S. africana* D.C. *Harv. and Sond. Fl. Cap. I*, 238; *Wood, Natal Plants*, Pl. 572; *Sim, Forests and Forest Flora*, 170, Pl. XXXIII, Fig. 4. *S. melanocarpa*, Arn. in *Hook. Journ. Bot.* III, 153; *Sonder in Harv. and Sond. Fl. Cap. I*, 238. *S. leucocarpa*, Arn. in *Hook Journ. Bot.* III, 153; *Sonder in Harv. and Sond. Fl. Cap. I*, 238.

Natal: Near Phoenix Crossing, 400 ft., February, *Wood in Natal Govt. Herb.*, 1373. Berea, Durban, March, *Wood*, 11804, and in *Natal Govt. Herb.*, 13648. Woods near Durban, 150 ft., January, *Wood*, 22; Port Natal, *Drege*; *Gueinzus*, without precise locality, *Gerrard and McKen.*, 525, and in *Natal Govt. Herb.*, 94; in woods at Inanda, 1800 ft., January, *Wood*, 827, and in *Natal Govt. Herb.*, 5884. Ixopo, 4000 ft., February, *Schlechter*, 6648.

DISTRIBUTION: Rhodesia, Pondoland, Kentani, Swaziland.

A. decipiens Radkl.

Young branches pubescent, becoming glabrous with age. *Leaves* petioled, trifoliate 4-8.5 cm. long; petiole 1-3 cm. long, semiterete, pubescent; leaflets 1-6 cm. long, 0.4-2.1 cm. broad, obovate or oblanceolate, obtuse, sometimes rounded at the apex, cuneate at the base, entire or sometimes remotely crenate, glabrous above and beneath or pubescent on the mid-rib and primary lateral veins. *Inflorescence* in stalked axillary lax racemes, 4-7 cm. long, stalk 1-3 cm. long, pubescent; rachis pubescent. *Larger sepals* 1.25 mm. long, 1.75 mm. broad, more or less broadly oblong, concave; smaller sepals 1.5 mm. long, oblong, deeply concave. *Petals* 1.5 mm. long, spatulate, truncate or 2-3-lobed at the apex, with 1-2 processes on the inner face, ciliate. *Filaments* 0.5 mm. long, ciliate; anthers 0.5 mm. long, oblong in outline. *Ovary* 0.5 mm. long, didymous, pubescent; style divided to the middle; stigmas 0.75 mm. long, linear, slightly recurved. *Fruit* 6 mm. long, 6 mm. in diameter, globose, glabrous. *Schmidelia decipiens*, Arn. in *Harv. and Sond.*, *Fl. Cap. I*, 239.

Not seen from Natal; stated by Wood to have been collected there by Dregé.

Schmidelia rehmanniana, Szyszcz., collected by Rehmann (No. 9038) near Durban, I have not seen, but from the description given suspect it to be *A. melanocarpus* Radkl.

DISTRIBUTION: Graaff-Reinet, Knysna, Kingwilliamstown, Albany, Uitenhage, Stockenström, Somerset East, Swellendam.

DODONEA LINN. F.

D. viscosa Linn.

Branches glabrous, with reddish bark. *Leaves* sessile, 4-9.5 cm. long, 1-3 cm. broad, oblanceolate, rounded and obtuse or shortly acuminate and subobtuse at the apex, narrowed at the base, glabrous, gland-dotted above and beneath, with the mid-rib prominent beneath, entire. *Inflorescence* a terminal panicle, peduncle glabrous, glandular. *Pedicels* 5-9 mm. long, glandular. *Sepals* unequal, 2.5-3 mm. long, 0.75-1.5 mm. broad, lanceolate or elliptic, obtuse, glandular without, usually three-veined. *Petals* absent. *Filaments* 0.5 mm. long; anthers 1.5 mm. long, 0.75 mm. broad, oblong. *Ovary* 1.5 mm. long, 1.5 mm. in diameter, globose, densely glandular, two-locular with two ovules in each loculus; style 2 mm. long, thick, terete, glabrous, bifid at the apex. *Fruit* 1 cm. long, 1.5 cm. broad, transversely oblong, two-winged, two-celled with two seeds in each cell. *Seeds* black, 1.5 mm. long, 1.5 mm. broad, discoid, glabrous. *Sonder in Harv. and Sond.*, *Fl. Cap. I*, 242.

Natal: Shores of Natal Bay, 50 ft., September, *Wood*, 7928, and in *Natal Govt. Herb.*, 2246; near Durban, September, *Wood*, 1928, 484; and in *Colonial Herb.*, 6794.

I find that this species differs from the generic description by having a two-locular ovary.—(*E. P. P.*)

HIPPOBROMUS E. & Z.

H. alata E. & Z.

A small tree, evergreen or nearly so, 10-30 ft. high (ex *Sim*). *Branches* minutely and densely pubescent on the young parts, becoming almost glabrous on the older parts. *Leaves* petioled, paripinnate, 3-6-jugate; petiole 0.7-2 cm. long, terete or somewhat flattened above, densely tomentose; rachis winged, tomentose; leaflets 1-3.5 cm. long, 0.7-2 cm. broad, obovate, obtuse, narrowed at the base, unequal sided, giving the leaflet a falcate appearance, darker above than beneath, glabrous, usually pubescent on the

mid-rib above and beneath, coarsely serrate usually on the convex side, sometimes entire, with the mid-rib prominent beneath. *Inflorescence* an axillary raceme shorter than the leaves, sometimes grouped into a dense terminal panicle. *Pedicels* 2 mm. long, terete, densely tomentose. *Sepals* unequal, 2.5–3 mm. long, 1.75–2.5 mm. broad, elliptic or subrotund, densely tomentose without, ciliated. *Petals* 3 mm. long, 2.5 mm. broad, obovate, glabrous. *Disc* annular. *Filaments* 2.5–5 mm. long, linear, glabrous, tapering gradually from the base upwards; anthers 1.5 mm. long, elliptic, in outline. *Ovary* rudimentary, tomentose. *Fruit* 0.7 mm. long, 1 cm. in diameter, three-celled with a single seed in each cell, pilose. *Seeds* 5 mm. long, 5 mm. in diameter, almost globose, somewhat flattened or channelled on the inner face, glabrous. *Sonder in Harv. and Sond. Fl. Cap. I*, 241. *Sim, Forest and Forest Flora*, 172, *Pl. XXXIV*.

Natal: Stellabush, Durban, September, *Van der Bijl in Natal Herb.*, 16243, 16244; without precise locality or collector in *Natal Govt. Herb.*, 93; Verulam, Wood, 606.

DISTRIBUTION: Port Elizabeth, Kentani, Somerset East, Uitenhage, Albany, Stocken-stroom.

PTAEROXYLON, E. & Z.

P. utile E. & Z.

Branches glabrous, youngest with greyish bark, turning dark brown in older branches. *Leaves* petioled, pinnately compound, 6–7-jugate, 3–12 cm. long; petiole 1–2.5 cm. long, minutely pubescent, furrowed above; leaflets 1–3 cm. long, 0.5–1.5 cm. broad, oblong, obtuse or retuse at the apex, sometimes shortly mucronate, unequal at the base, entire, with the mid-rib prominent above and beneath and the lateral veins distinct, glabrous. *Inflorescence* racemose, massed at the ends of the branches, about 3 cm. long. *Peduncle* pubescent. *Flowers* often in three-flowered cymes on the raceme. *Pedicels* 4 mm. long, pubescent, articulated at the base. *Bracts* 2 mm. long, lanceolate-linear. *Calyx* 1.5 mm. long, ovate, subobtusely, pubescent. *Petals* 5 mm. long, 2 mm. broad, oblong, obtuse, glabrous. *Filaments* 3 mm. long, linear, gradually narrowing from the base upwards; anthers 1.5 mm. long, elliptic in outline, flattened, cordate at the base. In female flowers: *Ovary* 1.5 mm. long, 1.5 mm. broad, almost circular in outline, somewhat flattened, pubescent. *Style* 1 mm. long, terete, pubescent; stigma bilobed, discoid. *Fruit* 1.3 cm. long, oblong in outline, somewhat compressed, dehiscent into two valves; valves deeply keeled, glabrous.

No specimen from Natal seen.

DISTRIBUTION: Uitenhage, Somerset East, Komgha.

SAPINDUS, LINN.

S. oblongifolius Sond.

Shrub or small tree 6–10 ft. high (ex Wood). *Branches* minutely pubescent. *Leaves* petioled, paripinnate, 4–11-jugate, 18–26 cm. long; petiole 5–11 cm. long, terete, glabrous or sparsely pilose; leaflets subsessile, 3.5–12.5 cm. long, 1.5–4 cm. broad, lanceolate, rarely elliptic, obtuse, glabrous, with the mid-rib prominent beneath, entire. *Inflorescence* a panicle up to 43 cm. long. *Peduncle* pubescent, the main axis of the panicle and branches densely tomentose. *Pedicels* 3 mm. long, terete, densely tomentose. *Calyx* 6 mm. long, 4 mm. broad, elliptic in outline, deeply concave, densely tomentose without, long-ciliated. *Petals* 6 mm. long, 3.5 mm. broad, spatulate, densely ciliated, with a flat claw appendage 1 mm. long and 2.25 mm. broad and densely ciliate arising from the inner face. *Filaments* 2.5 mm. long, linear, long-ciliated; anthers 1.5 mm. long, oblong. *Disc* deeply saucer-shaped with a rim 1.5 mm. high, thin, more or less triangular in outline with rounded angles. *Ovary* 3 mm. long, 4 mm. in diameter, sub-apocarpous, three-locular with a single ovule in each loculus, densely pilose. In male flowers the filaments are slightly longer and the ovary rudimentary, otherwise similar to the perfect flowers. *Wood, Natal Plants Pl.* 89. *Sim, Forests and Forest Flora*, 173, *Pl. XXXV*. *Sond. in Harv. and Sond. Fl. Cap. I*, 240.

Natal: Near Durban, 100 ft., *Wood*, 1917, and in *Natal Govt. Herb.*, 2213; Berea, Durban, 100-200 ft., May, *Wood*, 11869, and in *Natal Govt. Herb.*, 13053; Port Natal, *Saunderson* (?).

Wood states: "In the 'Flora Capensis,' Vol. I, 240, this plant is described as diœcious. We find, however, that it is certainly polygamous, and plants which in one season bear a larger number of perfect flowers, in another season bear male ones only, or with but very few perfect ones." In the Stellabush at Durban I saw trees well over 20 ft. high.—(*E. P. P.*)

DISTRIBUTION: Komgha.

BERSAMA, FRESEN.

Mid-rib on lower surface of leaf covered with long straight hairs..... *Stayneri*.
Mid-rib quite glabrous.

Leaves usually 2-3-jugate; leaflets obovate, usually rounded at the apex, obtuse..... *lucens*.

Leaves usually 4-jugate; leaflets oblong-lanceolate, acuminate, acute.. *abyssinica*.

B. *Stayneri* Phillips.

Bark thick and rough. *Branches* glabrous, rough with longitudinal furrows. *Leaves* pinnate, petioled, 7-16 cm. long, 4-5-jugate; petiole 1-2 cm. long, very densely pilose; rachis densely pilose; leaflets 1.7-5.2 cm. long, 0.8-1.5 cm. broad, oblong-lanceolate or elliptic, acute and mucronate at the apex, rarely rounded, slightly cuneate at the base, the mid-rib deeply sunk on the under surface, which is frequently somewhat rugose, prominent beneath and with the lateral veins distinct beneath, glabrous above pilose or pubescent beneath, especially on the mid-rib. *Inflorescence* a shortly peduncled many-flowered raceme, 3.5-6.5 cm. long; peduncle 1-1.5 cm. long, very densely pilose. *Bracts* 3 mm. long, linear, tomentose. *Pedicels* 5 mm. long, terete, tomentose. *Calyx* 6 mm. long, densely tomentose, divided almost to the base. *Petals* 1.3 cm. long, 3 mm. broad, spatulate-linear, tomentose, reflexed in open flowers. *Filaments* 5 mm. long, semiterete, monadelphous and villous at the base. *Ovary* densely villous; style 3 mm. long, terete; stigma subglobose. *Fruit* 2.7-3 cm. long, dehiscing by four valves; valves woody, covered with woody protuberances. *Seeds* reddish in colour, 1.1 cm. long, 6 mm. in diameter, ellipsoid with a waxy yellow arillus at the base.

Natal: Without locality, *Stayner in Herb. Bolus*; Stinkwood Forest, Ingeli, Natal, May, *Chilvers in Herb. Forest Dept.*, 1518; Mkanzeni, Riversides, Natal, *Henkel in Herb. Forest Dept.*, 2421.

B. *lucens* Szysz.

A shrub 8-10 ft. high (ex *Wood*). *Branches* glabrous, with wrinkled greyish bark. *Leaves* petioled, compound, imparipinnate, 2-4-jugate, 10-20 cm. long; petiole 1.5-5.5 cm. long, glabrous, rarely pubescent; leaflets 1.8-9 cm. long, 1.3-4.5 cm. broad, obovate or obovate-elliptic, rarely elliptic, very rarely subacute, slightly narrowed at the base, glabrous, with the mid-rib distinct above, prominent beneath, and with thickened wavy margins. *Inflorescence* a lax axillary or terminal raceme, 6-14.5 cm. long, many-flowered; peduncle pubescent. *Pedicels* 0.4-1 cm. long, terete, densely tomentose with fine adpressed hairs. *Bracts* 0.5 mm. long, ovate, pubescent. *Calyx* gamosepalous; lobes 3-4 mm. long, 2 mm. broad, ovate or ovate-elliptic, obtuse, finely but densely pubescent without and with fine adpressed hairs within, ciliated; the two posterior lobes connate and bifid at the apex. *Petals* dull yellow, 8.5 mm. long, 2.5 mm. broad, above, oblong, obtuse, narrowed into an evident claw, pubescent. *Disc* unilateral. *Filaments* all connate at the base, 1 cm. long, terete, glabrous except at the; base the anterior filaments ciliated at the base and forming a tomentose shield 3 mm. long and 2.5 mm. broad; the posterior filaments ciliated and pubescent at the base; anthers 2.5 mm. long, 1.5 mm. broad, oblong. *Ovary* 1 mm. long, 1.5 mm. broad, subglobose, densely villous, four-locular with a single ovule in each loculus; style 7 mm. long, cylindric, pubescent on the lower half;

stigma globose, faintly lobed. *Fruit* 1.7 mm. long, 1.7 cm. in diameter, finely but densely pubescent, four-celled, each cell with a single seed, splitting into four valves when ripe. *Seeds* bright scarlet, 7 mm. long, 7 mm. in diameter, subglobose, wrinkled, flat on one side, convex on the other, with a fleshy yellow-green arillus. *Wood, Natal, Pl.* 88. *Sim, Forests and Forest Flora*, 175. *Journ. Bot. XLV*, p. 13. *Natalia lucens, Hochst., Harv. and Sond. Fl. Cap. I*, 369.

Natal: Umbilo, 20–100 ft., May, *Wood*, 12612, and in *Natal Herb.*, 14571; near Verulam, 100–200 ft., January, *Wood*, 10207, and in *Natal Govt. Herb.*, 11411; Durban Flats, near Umgeni, September, *Wood in Natal Herb.*, 2868; near Stamford Hill, *Wood*; Groenberg, February, *Wood*, 808; near Durban, 200–300 ft., March, *Wood*, 6307; without locality, *Saunders*; Palmiet, near Durban, 200 ft., February, *Wood*, 7392; Port Natal, *Gueinzius*.

DISTRIBUTION: Swaziland, Transvaal, Kentani, Pondoland, Komgha.

B. abyssinica Fresen.

A tree. *Leaves* petioled, 18–22 cm. long, pinnate; petiole 4–5 cm. long, terete, glabrous; rachis sometimes winged between the uppermost pair of leaflets; leaflets usually in four pairs, 2.5–4 cm. apart, 4–9 cm. long, 1.7–3.5 cm. broad, oblong-lanceolate or ovate-lanceolate, acuminate, acute, entire or the margins sometimes serrated in the upper half, glabrous. *Inflorescence* racemose, 10–20 cm. long on a woody peduncle 5–7 cm. long. *Calyx* campanulate, silky. *Petals* five, ligulate-spathulate, twice as long as the calyx, densely and finely silky, finally, reflexed. *Stamens* five; in the male flower nearly as long as the petals; filaments monadelphous, silky. *Capsule* about 1 cm. long, 1 cm. in diameter, subglobose, 3–4-valved, densely pubescent, without woody protuberances. *Seeds* 6 mm. long, black.

This description was drawn up partly from a fruiting specimen collected by *Schimper* in Abyssinia and partly from the description in the “*Flora of Tropical Africa*.”

CONTENTS.

I. SOUTH AFRICAN ASCOMYCETES.....	65
II. THE THORN PEARS.....	83
III. THE GENUS OCHNA.....	87
IV. THE GENUS OLINIA.....	97
V. THE GENUS CYCLOPIA, VENT.....	105

SOUTH AFRICAN ASCOMYCETES

IN THE

NATIONAL HERBARIUM.

By ETHEL M. DOIDGE.

PART II.

51. **Aphysa Rhynchosiae** (Kalch & Cke.) Theiss et Syd.

Ann. Myc. XV (1917), p. 134.

Syn. *Stigmatea Rhynchosiae* Kalch. & Cke.; Syll. Fung. I, p. 543 (on *Rhynchosia gibba*, Inanda, Natal, leg. Wood).

Parodiella Schimperi P. Henn. Bull. Herb. Boiss., I, 1893, p. 118; Syll. Fung. XI, p. 260 (on *Rhynchosia elegans* and *Vigna* sp. Eritrea).

On *Rhynchosia gibba*, Umkomaas, Natal, 25.4.19, Bottomley [11884].

On *Rhynchosia* spp., Durban, 25.5.97, Medley Wood [356]; Durban, 15.6.15, Lansdell [9015]; Kentani, 29.9.15, Pegler (Pegler No. 2357) [9116]; Inanda, Medley Wood [9478]; Kowie, 15.7.19, Doidge [12362]; Howiesons Poort, 12.7.19, Doidge [12381].

On *Dolichos gibbosus*, Van Stadens Pass, 13.11.17, Doidge [10864].

Epiphyllous, forming spots 1-2 mm. diameter, which are made up of numerous minute black points. Each point represents a hemispherical, subcuticular perithecium, 170 μ in diameter and about 60 μ high. Free mycelium wanting. Basal line brown, thin, 0.5 μ thick, lying flat on the outer wall of the epidermis; the outer membrane convex, remaining covered by the cuticle, 6-7 μ thick, parenchymatous, without any decided radial orientation except at the periphery. Asci basal, somewhat converging towards the ostiole, sessile, paraphysate, somewhat thickened round the apex, 42-48 $\times$ 12-15 μ , eight-spored. Spores distichous, ellipsoid, two-celled, not constricted, rounded at both ends, light, grey-brown, 7-10 $\times$ 4 μ .

52. **Catacauma Schotiae** n. sp.

On leaves of *Schotia speciosa*: Humewood, Port Elizabeth, 23.3.12, Doidge [2294].

Stromata epiphyllous, scattered, minute, black, convex, 0.3-0.4 mm. diameter, round or oval, unilocular. Loculus developing between the epidermis and the palisade tissue, and with its base resting on the latter which becomes concave, ellipsoid, 300-350 μ diameter, 100-120 μ high in the centre; epidermal clypeus black, opaque, 30-40 μ thick, and covered by thick cuticle which is 15-16 μ thick: the clypeus does not always extend over the whole length of the loculus, being 200-250 μ long, lateral and basal walls of loculus consisting of delicate hyphae which are laterally compressed. Asci paraphysate, oblong or ellipsoid, tapering somewhat to the rounded apex, eight-spored, sessile, 80-87 $\times$ 17-20 μ . Spores distichous, hyaline, continuous, ellipsoid, rounded at both ends, 20-24 $\times$ 6-7 μ .

Cutacauima Schotiae Doidge, nov. sp.

Stromata epiphylla, sparsa, minuta, atra, convexa, subepidermales, 0.3–0.4 mm. diametro unilocularia. Loculi ellipsoidei, 300–350 μ diametro, 100–120 μ alti; clypeo atro, opaco, 30–40 μ crasso. Asci paraphysati, cylindranei v. ellipsoidei, octospori, sessiles, 80–87 $\times$ 17–20 μ . Sporae distichae, hyalinae, continuae, ellipticae, utrinque rotundatae, 20–24 $\times$ 6–7 μ .

Hab. in foliis *Schotiae speciosae*, Port Elizabeth, 23.3.12, leg. Doidge [2294]

53. *Dictyochorella Andropogonis* n. sp.

On leaves of *Andropogon nardus*, Tug-la Valley, near Goodoo, Natal, 16.5.20, Doidge [14104].

Stromata thickly strewn on the leaf surface, small, oval to elliptic, $\frac{1}{2}$ –1 mm. long, visible on both surfaces, slightly convex, loculi mostly opening towards the upper leaf surface; often a number of stromata arising close together become confluent and form streaks up to 5 mm. long in which the separate stromata can still be distinguished. Stromata with one or few loculi. Loculi occupying the whole of thickness of leaf, 190–240 μ diameter $\times$ 160–190 μ high. Clypeus over apex of loculi black, opaque, about 30 μ thick, extending considerably beyond loculus on each side: basal clypeus 20–25 μ thick; when the loculus does not reach basal clypeus, then the intervening space is filled with brown hyphae; lateral walls (and basal where there is no clypeus) up to 20 μ thick and composed of brown compact angular cells. Stylospores brown, oval, elliptic or pyriform; with a central (?) vacuole, 8–10 $\times$ 3–4 μ . Asci paraphysate, cylindrical, rounded at apex, eight-spored, 80–90 $\times$ 12–13 μ . Spores distichous or obliquely monostichous, with three or occasionally four transverse septa, constricted at septa, tapering towards rounded at ends, two central cells broader than terminal ones and one or both with a somewhat oblique longitudinal septum, 15–17 $\times$ 6–7 μ , fuscous-olivaceous, sometimes only one central cell has a septum and it is then broader than the other.

Dictyochorella Andropogonis Doidge, n. sp.

Stromata dense disposita, minuta, elliptica v. suborbicularia $\frac{1}{2}$ –1 mm. longa, in utraque foliorum pagina visibilia. Loculi uni v. pauci in quoque stromate, 190–240 diametro, 160–190 μ alti, clypeo amphigeno atro, opaco, circ. 30 μ crasso. Asci paraphysati, cylindranei, apice rotundati, octospori, 80–90 $\times$ 12–13 μ . Sporae distichae v. oblique monstichae, transverse 3-rare, 4-septatae, ad septa constrictae, loculis 1–2 plerumque mediis septo singulo longitudinale divisi, fuscae v. olivaceae, 15–17 $\times$ 6–7 μ .

Hab. in foliis *Andropogonis nardi*, Tugela Valley, Natal, 16.5.20, leg. Doidge [14104].

54. *Phaeodothis stenostoma* (Ell. et Tr.) Theiss et Syd.

Ann. Myc. XV (1917), p. 596.

Syn. *Phyllachora stenostoma* Ell. et Tracy, Journ. of Myc., 1890, p. 76; Syll. Fung. IX, p. 1029.

On *Panicum maximum*, Bluff, Durban, 7.7.11, Doidge [1611].

Stromata oval, about 600 μ long, 300–400 μ broad, or subcircular $\frac{1}{3}$ –1 mm. diameter, very irregular in outline, usually visible on both sides of the leaf, often only on one side and surrounded by a small, pale ring, later forming elliptic groups $\frac{1}{2}$ –1 cm. long and $\frac{1}{2}$ cm. broad. A dark epidermal clypeus is developed in both leaf surfaces. Loculi, flattened-spherical, often irregular in shape, mostly 170 μ diameter, 120–130 μ high, locular wall composed of delicate, light brown hyphae, fusing at the apex with the clypeus, 12–14 μ thick. Asci cylindrical, parietal, paraphysate, briefly pedicellate, 52–60 $\times$ 7–8 μ . Spores obliquely monostichous or subdistichous, oblong, club-shaped, fuscous, rounded at both ends, straight, 12–14 $\times$ 4 $4\frac{1}{2}$ μ , somewhat constricted at the septum.

55. *Phaeodothis Tristachyae* Syd.

Ann. Myc. X (1912), p. 41.

On leaves of *Tristachya leucothrix*, Mooi River, Natal, 15.4.11, Burt-Davy [1470].

Stromata amphigenous, scattered or in short series, oblong or linear, $\frac{1}{2}$ –1 mm. long, flat, black, opaque; loculi numerous, minute, 100–150 μ diameter. Asci cylindrical, somewhat obtuse at the apex, $45\text{--}70 \times 6\text{--}12 \mu$, eight-spored. Paraphyses numerous, filiform. Spores monostichous or distichous, fusiform, 1-septate, loculi equal, not or very slightly constricted, olivaceous-brown, $12\text{--}13\frac{1}{2} \times 3\text{--}4 \mu$, each cell 1–2 guttulate.

56. *Phyllachora Crotonis* (Cke.) Sacc.

Syll. Fung. II, p. 599; Ann. Myc. XV (1917), p. 523.

Syn. *Dothidea Crotonis* Cke., Grevillea X, p. 129.

Phyllachora crotonicola Rehm., Hedwigia XXIX, 1890, p. 110; Syll. Fung. IX, p. 601.

Phyllachora Julocrotonis Bres., Hedwigia, XXXV, 1896, p. 300; Syll. Fung. XIV, p. 668.

On leaves of *Croton sylvaticum*, Inanda, Natal, January, 1881, Medley Wood (Wood No. 406) [9489].

Stromata visible on both sides of the leaf, $\frac{1}{3}$ – $\frac{1}{2}$ mm. diameter, round, convex on both surfaces, on the upper surface smooth, shiny, often becoming more or less coalescent, and then up to 1 mm. in diameter and more or less irregular in outline. Single stromata unilocular, scattered irregularly over leaf surface. Loculi with an upper and lower epidermal clypeus which is no broader than the loculus. The lateral walls are not so well developed and are light brown. Loculi 200–300 μ diameter, spherical. Asci cylindrical to somewhat club-shaped, paraphysate, briefly pedicellate, $60\text{--}70 \times 10\text{--}14 \mu$. Spores distichous, one-celled colourless, ellipsoid, rounded at both ends, $14\text{--}16 \times 7\text{--}9 \mu$.

57. *Phyllachora Cynodontis* (Sacc.) Niessl.

Not. Pyren., p. 54; Syll. Fung. II, p. 602; Ann. Myc. XV (1917), p. 447.

Syn. *Phyalospora Cynodontis* Del., Bull. Soc. Myc. Fr., 1890, p. 183; Syll. Fung. IX, p. 599.

Phyllachora graminis var. *Cynodontis*, Sacc. Rabh. F. Eur., 2441.

On *Cynodon dactylon*, Pretoria, 12.7.03, Burt-Davy [145 and 170]; Groenkloof, Pretoria, 3.5.10, Burt-Davy [917]; Cedara, Natal, 18.1.11, Fisher [1172]; Vereeniging, 8.6.10, Burt-Davy [935]; Bloemfontein, 6.4.17, Potts [11311].

On *Cynodon incompletus*, Pretoria, 19.5.06, Pole Evans [17]; 19.6.11 [1587].

Stromata round-angular, scattered, 0.3–0.5 mm. diameter, sometimes coalescing, slightly convex, black. Clypeus epidermal, opaque, only on upper side of the stroma, and usually epiphyllous; on both sides of the leaf, but then the basal clypeus is shorter and not so well developed. Loculi small, spherical, 160–180 μ diameter, usually not penetrating right through the leaf, with a thin delicate wall, scattered or in small groups. Asci paraphysate, cylindrical, club-shaped, pedicellate, $50\text{--}60 \times 12\text{--}15 \mu$. Spores hyaline, monostichous or distichous, oval-ellipsoid, $10\text{--}12 \times 5\frac{1}{2}\text{--}6 \mu$.

58. *Phyllachora Evansii* Syd.

Ann. Myc. X (1912), p. 40; XIII (1915), p. 459.

On *Setaria sulcata*, Barberton, March, 1911, Pole Evans [1280]; Durban, 21.3.19, Bottomley [12231]; Isipingo, Natal, 27.3.18, Bottomley [11659].

On *Setaria nigrirostris*, Vereeniging, Transvaal, 26.2.17, Burt-Davy [10021].

On *Setaria aurea*, Barberton, 20.1.12, Doidge [2012]; Garstfontein, Pretoria District, 26.3.11, Pienaar [1265].

Stromata visible on both sides of the leaf, scattered or in rows, on yellowish leaf spots, ellipsoid, $\frac{1}{2}$ –1 $\times$ $\frac{3}{4}$ – $\frac{1}{2}$ mm., dull black; clypeus in both upper and lower epidermis, 20 μ thick, opaque. Loculi few, ellipsoid, 100–150 μ high, 260–230 μ diameter, with thinner, delicate, brown lateral walls. Asci cylindrical, club-shaped, 70–120 $\times$ 8–12 μ ; paraphysis numerous. Spores mono- or distichous, ellipsoid to oblong, rounded at both ends, straight, colourless, 14–18 $\times$ 6–8 μ . Conidia filiform or sickle-shaped, 15.25 $\times$ 1 μ .

59. *Phyllachora lucens* (Cke.) Sacc.

Syll. Fung. II, p. 600; Ann. Myc. XV (1917), p. 562.

Syn. *Dothidea lucens* Cke., *Grevillea* X, p. 128.

On leaves, Inanda, Natal, 1881, Medley Wood (Wood No. 581) [9487 and 10443].

Stromata unilocular, punctiform, only visible on the upper leaf surface, convex, black, shining, thickly crowded on yellow leaf spots, 6–10 mm. in diameter. Loculi flattened spherical, 220–260 μ diameter, 130–160 μ high; the base does not reach as far as the lower epidermis: the locular wall is 8–10 μ thick, consisting of brown cells, but weakly developed, and here and there not very distinct: at the apex it unites with the short black epidermal clypeus. Asci cylindrical, paraphysate, eight-spored, briefly pedicellate, 60–70 $\times$ 8–12 μ . Spores distichous or obliquely monostichous, oblong, colourless, one-celled, straight, with blunt ends, 13–15 $\times$ 4 $\frac{1}{2}$ –5 $\frac{1}{2}$ μ .

60. *Phyllachora Penniseti* Syd.

Ann. Myc. XIII (1915), p. 39 (on leaves *Pennisetum Benthami*, Congo).

On leaves of *Pennisetum sphacelatum*, Tugela Valley, near Goodoo, Natal, 16.5.20, Doidge [14102].

Stromata conspicuous on both leaf surfaces, scattered or crowded and arranged in rows, round or oblong, $\frac{1}{2}$ –1 mm. long, or forming larger stromata by confluence, black, opaque, plurilocular. Loculi irregular, large, 280–400 μ long, 200–300 μ high. Clypeus in both upper and lower epidermis, 15–25 μ thick, lateral wall of loculus 8–10 μ broad, brown: stylospores straight or flexuous, continuous, filiform, hyaline, 18–35 $\times$ 1 μ . Asci clavate, rounded at the apex, 70–95 $\times$ 20–26 μ , eight-spored, paraphysate; spores distichous or obliquely monostichous, broadly rounded at both ends, continuous, hyaline, 17–24 $\times$ 11–18 μ .

61. *Phyllachora heterospora* P. Henn.

Syll. Fung. XXII, p. 425; Ann. Myc. XV (1917), p. 453.

On leaves of *Panicum maximum*, Bluff, Durban, 7.7.11, Doidge [1611]; Kentani, 1.5.14, Pegler [7773]; Kentani, May, 1915, Pegler (Pegler No. 1953) [9095].

On *Panicum* sp., Barberton, 22.8.12, V. d. Bijl [5137].

Both leaf surfaces are comparatively thickly beset with stromata, which are round or elliptic, $\frac{1}{2}$ –1 mm. diameter, convex, somewhat shiny, not usually coalescing, and not situated on discoloured leaf areas, the whole leaf becomes yellow. Clypeus epidermal, 20 μ thick, on one or both leaf surfaces. Loculi about 4–10 in each stroma; when the stroma is only on one side of the leaf, occupying about two-thirds of its thickness, closely crowded, 180–230 μ diameter, 160 μ high. When two stromata occur at the same point on opposite sides of the leaf the loculi only reach the centre. Wall brown, about 10 μ thick. Asci cylindrical, paraphysate, 60–80 $\times$ 10–16 μ , eight-spored; spores mostly distichous, ellipsoid, tapering somewhat to both ends but rounded, one-celled, hyaline, 13–17 $\times$ 7–8 μ .

62. *Phyllachora Tecleae* n. sp.

On leaves of *Teclea natalensis*, 5.6.15, Kentani, Pegler (Pegler No. 2324) [9079].

Stromata in small groups on yellow leaf spots, developing on both sides of the leaf, with a preference for the upper surface, visible on both leaf surfaces, oval, ellipsoid or almost round, 0.5 to 2.5 mm. long, black, somewhat shiny, raised on both surfaces, but not

convex, groups of stromata are often oval or elliptic in outline between the lateral veins. The normal thickness of the leaf is about $240\ \mu$, the thickness of the stroma is $300\text{--}400\ \mu$. Loculi occupying the whole thickness of the leaf, flattened spherical, $300\text{--}400\ \mu$ diameter; epidermal clypeus on both leaf surfaces very thick, black, opaque, on the upper surface $40\text{--}50\ \mu$ thick; basal clypeus thinner, $23\text{--}25\ \mu$; lateral wall consisting of thin, delicate, laterally compressed hyphae; leaf tissue between the loculi permeated in every direction by brown hyphal threads. Asci richly paraphysate, cylindrical, eight-spored, with a short foot, $80\text{--}90 \times 10\text{--}13\ \mu$. Spores obliquely monostichous, one-celled, hyaline, ellipsoid or subfusoid, tapering to each end, but rounded not pointed at ends, $13\text{--}17 \times 5\text{--}6\ \mu$.

Phyllachora Tecleae Doidge, nov. sp.

Stromata amphigena plerumque epiphylla, maculis flavidulis insidentia, elliptica v. suborbicularia, $0\cdot5\text{--}2\cdot5$ mm. diametro atra, parum nitidula. Loculi suborbiculares, $300\text{--}400\ \mu$ diametro, clypeo epidermale amphigeno, atro, opaco, $40\text{--}50\ \mu$ crasso. Asci copiose paraphysati, cylindracei, octospori, $80\text{--}90 \times 10\text{--}13\ \mu$. Sporae oblique monostichae, continuae, hyalinae, ellipsoidae v. sub-fusoidae, utrinque attenuatae sed rotundatae, $13\text{--}17 \times 5\text{--}6\cdot5\ \mu$.

Hab. in foliis *Tecleae natalensis*, Kentani, 5.6.15, leg. Pegler [9079].

63. *Phyllachora Tricholaenae* P. Henn.

Engl. Bot. Jahrb. XXIII, p. 541; Syll. Fung. XIV, p. 672; Ann. Myc. XV (1917), p. 461.

On leaves of *Tricholaena rosea*, Winklespruit, Natal, 14.2.13, Pole Evans [5641].

Stromata oval, black, shining, slightly convex, scattered or crowded, $\frac{1}{2}\text{--}1$ mm. long, clypeus formed in epidermis of both leaf surfaces, but is more extensive on the side of the apex of the loculus than on the basal side. Loculi delimited at base and apex by the opaque clypeus, lateral walls thinner and more delicate. Loculi spherical, about $240\ \mu$ diameter; asci club-shaped, pedicellate, $55\text{--}60 \times 15\text{--}18\ \mu$, paraphysate. Spores distichous, elliptic, abruptly rounded at ends, $12\text{--}13 \times 7\text{--}8\ \mu$.

64. *Telimena Arundinariae* n. sp.

On leaves of *Arundinaria tessellata*, Goodoo Bush, Natal, 20.5.20, Doidge [14103.]

Stromata on yellow leaf spots, scattered or more rarely in groups, minute, oval to ellipsoid, $0\cdot5\text{--}0\cdot7$ mm. long, dull black, convex on upper leaf surface, penetrating through leaf and becoming visible as minute, dull-black specks on the lower surface. Loculi usually one, sometimes two in each stroma, $120\text{--}180\ \mu$ diameter, $100\text{--}120\ \mu$ high, occupying from three-quarters to whole of thickness of leaf, with a dark epidermal clypeus on the upper side, $15\text{--}20\ \mu$ thick; when the loculus occupies whole thickness of leaf there is also a clypeus developed in the lower epidermis; remainder of locular wall composed of delicate, laterally compressed hyphae. Asci cylindrical, paraphysate, rounded at apex, eight-spored, briefly stipitate, $100\text{--}120 \times 15\text{--}18\cdot5\ \mu$.

Spores obliquely monostichous or distichous, cylindrical, fusiform, tapering somewhat to both ends, four-celled, cells subequal in length, or middle cells somewhat larger than the terminal ones, hyaline or subhyaline, $27\text{--}33 \times 8\text{--}9\cdot5\ \mu$.

Telimena Arundinariae Doidge, nov. sp.

Stromata epiphylla, maculis flavidulis insidentia, sparsa v. gregaria, minuta, elliptica, atra, $0\cdot5\text{--}0\cdot7$ mm. longa, loculos $1\text{--}2$ includentia, in hypophylla etiam conspicua. Loculi $120\text{--}180\ \mu$ diametro, $100\text{--}120\ \mu$ alti, clypeo epiphylo v. amphigeno, atro, $15\text{--}20\ \mu$ crasso. Asci cylindracei, paraphysati, apice rotundati, octospori, breviter pedicellati, $100\text{--}120 \times 15\text{--}18\cdot5\ \mu$. Sporae oblique monostichae v. distichae, cylindraceae v. fusiformae, parum utrinque attenuatae, 3-septatae, hyalinae v. sub-hyalinae, $27\text{--}33 \times 8\text{--}8\cdot5\ \mu$.

Hab. in foliis *Arundinariae tessellatae*, Natal, 20.5.20, leg. Doidge [14103].

65. *Telimena corticicola* n. sp.

On woody stems of *Helichrysum* sp., Alice, C.P., 18.11.17, Doidge [10975].

Stromata scattered irregularly on the greyish bark, minute, dull black, oval to narrow ellipsoid, 0.5 to 0.8 mm. long, somewhat convex, unilocular, or with few loculi sunken in the cortex, not erumpent. Loculi oval to flattened spherical, opening by an apical pore, 300–340 μ diameter, 200–240 μ high, fusing at the apex with the black, opaque, clypeus which is 18–20 μ thick and extends beyond the apex of the loculus a distance of 300–400 μ . The space between the clypeus and the upper part of the loculus is filled with a black opaque hyphal web similar to that forming the clypeus, and from this a few dark hyphae run out into the substratum. The lower part of the locular wall consists of delicate, almost colourless hyphae, which are strongly compressed laterally. Asci paraphysate, cylindrical, rounded at the apex, very briefly pedicellate, 80–110 $\times$ 20–23 μ , rather thick walled. Spores distichous, colourless, cylindrical, rounded at both ends, four-celled, primary cross-wall in the centre, each of two primary cells being later divided in the centre by a secondary wall. Thick walled, wall about 2 μ thick, outer edge smooth, inner curving inward to meet transverse walls and giving the spore the appearance of being constricted, 24–27 $\times$ 10–13 μ .
Telimena corticicola Doidge, nov. sp.

Stromata sparsa, corticicola, atra, elliptica, 0.5–0.8 mm. longa, leniter convexa, loculos paucos includentia. Loculi ovati v. suborbiculares, 300–400 μ diametro, 200–240 μ alti, clypeo opaco, 18–20 μ crasso. Asci paraphysati, cylindranei, apice rotundatae, brevissime pedicellati, 80–110 $\times$ 20–23 μ . Sporae distichae, hyalinae, cylindraceae, utrinque rotundatae, 3-septatae, 24–27 $\times$ 10–13 μ .

Hab. in cortice *Helichrysi* sp., Alice, C.P., 18.11.17, leg. Doidge [10975].

66 *Systremma Pterocarp* n. sp.

On leaves and twigs of *Pterocarpus ceriseus*, Khami Ruins, Rhodesia, 14.7.20, Bottomley [14101].

Leaf stromata epiphyllous, rarely hypophyllous, scattered, situated on yellowish leaf spots, dull black, round or ellipsoid, 1–1.5 mm. diameter, 450–600 μ high, usually with a concentric zone of secondary stromata with a radius of about 1 mm. from centre of primary stroma. Stromata on twigs and midribs of leaves similar, but usually oval to ellipsoid, somewhat larger and without secondary stromata.

The stromata arise between the epidermis and the palisade cells, the cuticle becomes arched, and a dark-brown epidermal clypeus is formed. The base is also composed of small dark cells, and rests on the palisade cells, but the central part of the stroma is formed of paler, more delicate tissue in which are produced one large and a number of smaller conidia! loculi more or less spherical but irregular in size. The conidia are round, oval, pyriform, somewhat angular, brown, and 5–7 μ diameter. The conidial stroma attains a height of 150–250 μ , and does not rupture the cuticle. The ascostroma arises from beneath the conidial stroma, and is similar to it in structure, the latter is pushed up and the cuticle ruptured, its dark base forming the covering layer of the ascostroma. Loculi few, spherical to ellipsoid, or pyriform, 120–200 μ diameter. Asci numerous aparaphysate, eight-spored, thin-walled, evanescent, oblong-cylindrical, about 70 $\times$ 10 μ . Spores mono-half distichous, two-celled, brown, cylindrical, with blunt ends, slightly constricted, 10–13.3 $\times$ 5–6.3 μ , upper loculus 6–8.6 μ , lower 3.3–5 μ long.

Systremma Pterocarp Doidge, nov. sp.

Stromata epiphylla, rarius hypophylla, maculis flavidulis insidentia, atra, orbicularia v. elliptica, 1–1.5 mm. diametro, 450–600 μ alta, subepidermalia. Loculi pauci, orbiculares, elliptici v. pyriformi, 120–200 μ diametro. Asci numerosi, aparaphysati, octospori, oblongi v. cylindranei, 70 $\times$ 10 μ . Sporae monostichae v. subdistichae, 1-septatae, brunneae, cylindranei, utrinque subtruncatae, leniter constrictae, 10–13.3 $\times$ 5–6.3 μ ; loculo supero majore 6–8.6 μ longo, infero 3.3–5 μ longo.

Hab. in foliis caulisque *Pterocarp* *cerisei*, Khami Ruins, Rhodesia, 14.7.20, leg. Bottomley [14101].

67. *Teratosphaeria fibrillosa* Syd.

Ann. Myc. X (1912), p. 39; XV (1917), p. 631.

On leaves of *Protea grandiflora*. Bains Kloof, near Wellington, 26.2.11, Doidge [1217]; 21.2.12, Doidge [2059]; 5.4.12, Stoneman [2232].

Loculi ellipsoid or pyriform, perithecium-like, 150–220 μ diameter, sunken under the epidermis, arranged in centrifugal, irregularly branched, radiating lines, to a diameter of 1 cm. Locular wall consisting of concentric layers of delicate stromatic hyphae. Apex of loculus penetrating the epidermis but not protruding. No true ostiole. Vegetative stroma meagre, penetrating into the intercellular spaces of the mesophyll below the epidermis in the form of perpendicular hyphal strands. At the apex the hyphae also penetrate the adjacent epidermal cells, but without forming a clypeus. Asci basal, quite sessile, club-shaped, thickened round apex, 70–110 $\times$ 25–35 μ , paraphysate, eight-spored. Spores distichous or tristichous cylindrical, straight, medially slightly constricted at the septum, light-brown, rounded at both ends, 42–52 $\times$ 8–9 μ .



Fig. 1

68. *Meliola capensis* (K. & Cke.) Th.

Ann. Myc. X (1912), p. 19; Trans. Roy. Soc. S. Afr., V, p. 731.

Syn. *Asterina capensis* K. & Cke., Grevillea 9, p. 32.

This fungus occurs commonly on *Hippobromus alatus*. Its mycelial setae were described as being straight, simple, opaque, 200–400 $\times$ 5–10 μ , pellucid near the acute apex. Examination of a large number of collections shows that in the majority at least 50 per cent. of the setae are forked at the apex. The apex of the setae should therefore be described as simple, acuminate, or with two to three acute teeth. [Fig. 2.]

69. *Meliola Carissae* n. sp.

Amphigenous, forming irregular spots 5–6 mm. diameter or larger areas by confluence. Mycelium forming an irregular network, branches opposite, anastomosing. Hyphae slightly undulate, 6–7 μ thick, not infrequently constricted at septa, cells 20–30 μ long; capitate hyphopodia alternate, 20–30 $\times$ 10–13 μ , terminal cell variable in form, sometimes club-shaped or sublobed, but most frequently truncate. Mucronate hyphopodia not very numerous, unilateral or alternate, with a short thick neck, 18–20 $\times$ 6–7 μ . Mycelial setae fairly numerous, simple, straight, 550–700 μ long, 9–10 μ thick at base, and tapering very gradually to the blunt apex, black and opaque at base, but becoming brown and slightly translucent near apex. Perithecia not very numerous, 180–200 μ diameter, spherical, black, surface cells slightly convex. Asci two-spored, evanescent, spores four-septate, oblong, slightly constricted, rounded at both ends, brown, 43–47 $\times$ 16.5–20 μ . [Fig. 1.]

On leaves of *Carissa arduina*, Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12296]; Van Stadens, C.P., 13.11.17, Doidge [10870]; Bathurst, C.P., 14.7.19, Doidge [12341]; Kovie, C.P., 15.7.19, Doidge [12361].

Meliola Carissae Doidge, nov. sp.

Amphigena, maculas irregulares, 5–6 mm. diametro formans; hyphis subundulatis, 6–7 μ crassis, nonnunquam ad septa constrictis, cellulis, 20–30 μ long, ramis oppositis, anastomosantibus; hyphopodiis capitatis alternis, 20–30 $\times$ 10–13 μ , cellula superiore




Fig 2.

variabile, aut clavata v. sublobata aut truncata; hyphopodiis mucronatis paucis, unilateralibus v. alternis, 18–20 $\times$ 6–7 μ ; setis mycelicis satis numerosis, rectis, simplicibus, 550–700 μ longi, basi opaco, 9–10 μ crassis, apice obtuse sensim attenuatis; peritheciis 180–200 μ diametro, ascis bisporis; sporis 4-septatis, oblongis, leniter constrictis, utrinque rotundatis, brunneis, 43–47 $\times$ 16.5–20 μ .

Hab. in foliis *Carissae arduinae*, Pirie Forest, 8.7.19, leg. Doidge [12296].

70. *Meliola dracoenicola* Pat. et Har.

Bull. Soc. Myc. Fr. XXIV (1908), p. 14, Sacc.; Syll. Fung. XXII, p. 56.

On leaves of *Behnia reticulata*, Louis Trichardt, 8.5.19, Putterill [11848].

This fungus was originally described from the Congo on *Dracoena* sp.; it has not been previously recorded from South Africa.

Amphigenous, black, forming spots 2–5 mm. diameter, easily detachable from the leaf surface. Hyphae brown, straight, 7–9 μ thick; cells 17–20 μ long.; branches opposite, numerous, anastomosing; capitate hyphopodia alternate or unilateral, stipitate, 17–20 $\times$ 10–13 μ , stipe cell short, cylindrical, 5–7 μ long, head cell globular; mucronate hyphopodia numerous, alternate or opposite, interspersed with the capitate hyphopodia, pyriform or

ampulliform with a straight or curved neck, $15-20 \times 6-7 \mu$. Mycelial setae very numerous, erect, abruptly geniculate at the base or taking a broad curve upwards, opaque, $250-350 \mu$ long, $10-11.5 \mu$ thick, dividing near the apex into two or three short spreading branches, $10-17 \mu$ long, each of these being divided at the apex into two or three small tapering branches, $6-10 \mu$ long, occasionally the latter are themselves bidentate. Perithecia not numerous, black, carbonaceous, globose, $250-300 \mu$ diameter. Spores straight, brown, cylindrical, four-septate, constricted obtuse, $43-46 \times 16 \mu$. [Fig. 3.]

71. *Meliola oleicola* n. sp.

Sub. *M. amphitricha*, Proc. Roy. Soc., South Africa, V, p. 733.

Amphigenous, forming round, dull-black spots, 2-4 mm. diameter, or greater areas by confluence. Mycelium olivaceous, very freely branched and interwoven; branches



Fig. 3.

opposite, anastomosing: capitate hyphopodia alternate, stipitate, $26-30 \mu$ long, terminal cell ovate or club-shaped, straight or somewhat curved, $18-20 \times 10 \mu$, stipe cell short, closely appressed to or forming an acute angle with the branch. Mucronate hyphopodia on separate branches, opposite, alternate or unilateral, slender ampulliform, base not much wider than curved neck, $20-23 \times 6-7 \mu$. Mycelial setae rather numerous, scattered, simple, straight or slightly flexuous, $240-300 \mu$ long, $9-10 \mu$ thick at base, opaque up to acute apex. Perithecia not very numerous, globose, $220-250 \mu$ diameter, verrucose: surface cells convex. Asci two to four spored, early evanescent. Spores oblong, four-septate, very slightly constricted, broadly rounded at both ends, chestnut-brown when mature, $45-50 \times 20-23 \mu$. [Fig. 4.]

On leaves of *Olea laurifolia*, Buccleuch, Natal, 17.7.18, Doidge [11557]; Lemana, Zoutpansberg District, 7.8.11, Doidge [1835]; Tabankulu, Pondoland, 4.4.15 [9103]; Woodville Forest, George, 11.11.17, Doidge [10937].

On *Olea Pegleri*, Kentani, 29.8.14, Pegler (Pegler 1948) [8382-8785].

On *Oleaceae* (?), Woodbush, Zoutpansberg District, 4.8.11, Doidge [1767].

Meliola oleicola Doidge, nov. sp.

Amphigena, maculas atrae orbiculares 2-4 mm. diametro formans. Mycelium ramosum, intertextum, ramis oppositis anastomosantibus, hyphopodiis capitatis alternis, stipitatis, 26-30 μ long, cellula superiore ovata v. clavata, recta v. leniter curvata, 18-20 $\times$ 10 μ ; cellula inferiore breve, hyphopodiis mucronatis oppositis, alternis v. unilateralibus, ampullaceis, 20-23 $\times$ 6-7 μ ; setis mycelicis satis numerosis, sparsis, opacis, rectis v. leniter flexuosis, simplicibus, 240-300 μ long, basi 9-10 μ crassis, apice acutis. Perithecia globosa, 222-250 μ diametro, verrucosa. Asci 2-4 spori, mox evanescentes. Sporae oblongae 4-septatae, leniter constrictae, utrinque rotundatae, 45-50 $\times$ 20-23 μ .

Hab. in foliis *Oleae laurifoliae*, Buccleuch, Natal, 17.7.18, leg. Doidge [11557].



Fig. 4

72. *Meliola Panici* Earle.

Muhlenbergia, 1901, p. 12, Sacc.; Syll. Fung. XVII, p. 550.

On leaves of *Stipa dregeana* var. *elongata*, Langholm Estates, Bathurst District, 14.7.19, Doidge [12358].

The type specimen was collected in Porto Rico, and this species has not previously been recorded from South Africa. The original description is as follows:—

Forming minute spots 1-3 mm. diameter, which often become confluent. Hyphae 6-8 μ thick, sparingly branched. Capitate hyphopodia usually alternate and forming a broad angle with the hyphae, 16-20 $\times$ 6-8 μ , terminal cell broader than long, irregular, usually three to four lobed. Mucronate hyphopodia numerous, opposite, inserted at right angles, broadly ampulliform, paler, 12 μ long. Mycelial setae numerous, straight, simple, obtuse, obscurely septate, opaque, 400-600 $\times$ 8-10 μ , tapering gradually towards the apex. Perithecia scattered, black, three to four spored, evanescent. Spores cylindrical, compressed, four-septate, slightly constricted, rounded at both ends, 40 $\times$ 12 $\times$ 8 μ .

The South African specimen differs slightly from the above description in that the mycelium is rather dense and copiously branched and anastomosing; the cells of the hyphae are rather short, 13–16 μ long. [Fig. 5.]

73. *Irene Zeyheri* n. sp.

Amphigenous, forming thin black spots 2–4 mm. diameter, or larger areas by confluence; mycelium not dense, forming an irregular network; branches opposite or irregular, anastomosing. Hyphae straight or somewhat sinuous, 6–6.5 μ thick: cells 20–30 μ long; capitate hyphopodia alternate, 18–23 μ long, terminal cell very variable in form, lobed in various ways, 6–14 μ broad at apex; mucronate hyphopodia opposite or unilateral, interspersed with the capitate hyphopodia; ampulliform 16–20 $\times$ 6–7 μ . Mycelial setae none. Perithecia scattered, spherical, black, carbonaceous, 190–220 μ diameter, surface cells convex. Asci two-spored, evanescent, spores oblong, brown, four-septate, slightly constricted, rounded at both ends, 39–44 $\times$ 16–18 μ .



Fig 5

On leaves of *Eugenia Zeyheri*, Howiesons Poort, 12.7.17, Doidge [12388]; Pirie Forest, 8.7.19, Doidge [12272]; Van Stadens Pass, 13.11.17, Doidge [10878].

Irene Zeyheri Doidge, n. sp.

Amphigena, maculas, tenues, atrae formans. Mycelium effusum, hyphis rectis v. leniter sinuosis, 6–6.5 μ crassis, ramis oppositis v. irregularibus, anastomosantibus; hyphopodiis capitatis alternis. 18–23 μ longis, cellula superiore varia, lobata, apice 6–14 μ lata; hyphopodiis mucronatis oppositis v. unilateralibus, ampullaceis 16–20 $\times$ 6–7 μ ; setis mycelicis nullis. Perithecia sparsa, globosa, atra, 190–220 μ diametro. Asci bispori. Sporae oblongae, brunneae, 4-septatae, leniter constrictae, utrinque rotundatae, 39–44 $\times$ 16–18 μ .

74. *Calothyrium Psychotriae* n. sp.

Hypophyllous, forming thin, rusty-brown patches 10–12 mm. diameter. Mycelium fuscous, branching and anastomosing irregularly, hyphae about $5\ \mu$ thick, without hyphopodia. Thyriothechia in groups, very pale, flattened hemispherical, or irregular in shape on account of lateral pressure of adjacent thyriothechia, $400\text{--}430\ \mu$ diameter; composed of radiating hyphae $4\text{--}5\ \mu$ thick, margin entire, but more or less undulate. Asci fairly numerous, paraphysate, eight-spored, spherical or somewhat pyriform, $50\text{--}53\ \mu$ diameter, or $60\text{--}64 \times 30\ \mu$. Spores conglobate, hyaline thick-walled, two-celled, slightly constricted, oblong, rounded at both ends, cells subequal, $26\text{--}29 \times 11\cdot5\text{--}13\ \mu$.

On leaves of *Psychotria capensis* (same leaves as *Meliola amphitricha*), 10.7.11, Amanzimtoti, Natal, Doidge [1575].

Calothyrium Psychotriae Doidge, nov. sp.

Hypophyllis, maculas tenues, 10–12 mm. diametro formans. Mycelium fuscum, ramosum, hyphis circ. $5\ \mu$ crassis, hyphopodiis nullis. Thyriothechia gregaria rotundata v.



Fig. 6.

mutua pressione angulata v. compressa, $400\text{--}430\ \mu$ diametro, radiatim ex hyphis $4\text{--}5\ \mu$ crassis contexta, ad ambitum haud fimbriata. Asci paraphysati, octospori, globosi v. subpyriformi, $50\text{--}53\ \mu$ diametro vel $60\text{--}64 \times 30\ \mu$. Sporae conglobatae, hyalinae, 1-septatae, leniter constrictae, oblongae, utrinque rotundatae, $26\text{--}29 \times 11\cdot5\text{--}13\ \mu$.

Hab. in foliis *Psychotriae capensis*, Amanzimtoti, Natal, 10.7.11, leg. Doidge [1575].

75. *Asterina rotonicola* n. sp.

On leaves of *Croton rivularis*, East London, 19.7.19, Doidge [12427]; Langholm Estates, Bathurst, 14.7.19, Doidge [12355].

Epiphyllous, forming thin black, carbonaceous spots, 2–3 mm. diameter. Mycelium pale fuscous, not dense, branching and anastomosing irregularly. Hyphae undulating, $3\text{--}3\cdot5\ \mu$ thick. Cells $18\text{--}25\ \mu$ long. Hyphopodia unilateral or alternate, minute, continuous,

pyriform, 6-7 μ high $\times$ 3.5 μ . Thyriothecia rather numerous, scattered, flattened hemispherical, 120-140 μ diameter, formed of radiating hyphae about 3 μ diameter, margin not fimbriate, dehiscing by a stellate fissure, no basal membrane. Asci aparaphysate, eight-spored, subspherical or ovate, sessile, 36-40 $\times$ 23-33 μ . Spores conglobate, two-celled, brown, slightly constricted, cells subequal, epispore rough at maturity, 17-20 $\times$ 10 μ . Pycnidia similar in form to thyriothecia, but somewhat smaller, about 100 μ diameter; conidia ovate, brown, continuous, without a hyaline band, 17-20 $\times$ 12-13 μ .




Fig. 7

Asterina crotonicola Doidge, nov. sp.

Epiphylla, maculas tenues atras, 2-3 mm. diametro formans. Mycelium fuscum ramosum, hyphis undulatis 3-3.5 μ crassis, cellulis 18-25 μ longis; hyphopodiis unilateralibus v. alternis, minutis, continuis, pyriformis, 6-7 μ altis, 2.5 μ latis. Thyriothecia satis numerosa, sparsa, orbicularia, 120-140 μ diametro, radiatim ex hyphis circ. 3 μ crassis contexta, stellatim dehiscencia ad ambitum haud fimbriata. Asci aparaphysati, octospori, subglobosi v. ovati, sessiles, 36-40 $\times$ 23-33 μ . Sporae conglobatae, 1-septatae, brunneae, in maturitate verrucosae, 17-20 $\times$ 10 μ . [Fig. 7.]

Hab. in foliis *Crotonis rivularis*, East London, 19.7.19, leg. Doidge [12427].




Fig. 8

76. *Asterina Scolopiae* n. sp.

On leaves of *Scolopia Zeyheri*, East London, 19.7.19, Doidge [12414].

On *Scolopia* sp., Mossel Bay, 22.7.15, Pole Evans [9067].

Hypophyllous, forming very thin spots about 5 mm. diameter. Mycelium delicate, diffuse. Hyphae straight or somewhat undulate, branching and anastomosing irregularly, 3-3.3 μ thick; hyphopodia unilateral or alternate, briefly stipitate, often at right angles to hypha, cylindrical, straight or slightly curved, 10-13.3 $\times$ 3-3.5 μ . Thyriothecia scattered,

flattened-hemispherical, 150–170 μ diameter, composed of radiating hyphae about 3 μ thick; at margin a number of these continue to extend towards and anastomose with vegetative hyphae, dehiscing by a stellate fissure. No basal membrane. Asci subspherical paraphysate, eight-spored, sessile, 33–40 μ diameter. Spores conglobate, two-celled, slightly constricted, ellipsoid, cells subequal, 20–23.5 $\times$ 9–11 μ . Pycnidia similar in form to thyriothecia, but smaller, 80–90 μ diameter. Conidia brown, continuous without a hyaline band, angular kite-shaped, 18–25 $\times$ 10–13.5 μ . [Fig. 8.]

Asterina Scolopiae Doidge, nov. sp.

Hypophylla, maculas tenues, circ. 5 mm. latas, formans. Mycelium laxum, hyphis rectis v. subundulatis, 3–3.3 μ crassis; hyphopodiis unilateralibus v. alternis, breviter stipitatis, cylindraceutis, rectis v. leniter curvatis, 10–13.3 $\times$ 3–3.5 μ . Thyriothecia sparsa, orbicularia, 150–170 μ diametro, radiatim ex hyphis circ. 3 μ crassis contexta, ambitu plus minus fimbriata, stellatim dehiscencia. Asci subglobosi, paraphysati, octospori, sessiles, 33–40 μ diametro. Sporae conglobatae, 1-septatae, leniter constrictae, ellipsoidae, cellulis subaequalibus, 20–23.5 $\times$ 9–11 μ .

Hab. in foliis *Scolopiae Zeyheri*, East London, 19.7.19, leg. Doidge [12414].

77. *Lembosia piriensis* n. sp.

On leaves of *Trichocladus ellipticus*, Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12301].

Epiphyllous, on leaf spots up to 1 cm. diameter, sometimes numerous and confluent, covering greater part of leaf surface. Hyphae slender, fuscous, 3–3.5 μ thick, branching and anastomosing irregularly. Hyphopodia sessile, about 4.5 μ diameter, hemispherical or sublobed, developing at irregular intervals. Thyriothecia scattered, ellipsoid, straight, simple or occasionally forked, 300–400 $\times$ 190–240 μ , composed of radiating hyphae about 3 μ thick, margin fimbriate. Asci paraphysate, eight-spored, ovate, somewhat thickened round the apex, sessile, 37–40 $\times$ 20–24 μ . Paraphyses hyaline, filiform, about 2 μ thick, equalling or slightly exceeding the asci. Spores two-celled, slightly constricted, ellipsoid, upper loculus broader, 17–20 $\times$ 6–7 μ , brown when mature. Pycnidia flattened-hemispherical, 100–120 μ diameter. Conidia minute, hyaline, bacilloid.

Lembosia piriensis Doidge, nov. sp.

Epiphylla, maculas usque 1 cm. latas insidentia; hyphis tenuibus, fuscis, 3–3.5 μ crassis, ramosis; hyphopodiis sessilibus, circ. 4–5 μ diametro hemisphaericis, v. sublobatis, dispersis. Thyriothecia sparsa, elliptica, recta, simplices v. nonnunquam bifurcata, 300–400 $\times$ 190–240 μ , radiatim ex hyphis 3 μ crassis contexta, ambitu fimbriata. Asci paraphysati, octospori, ovati, sessiles, 37–40 $\times$ 20–24 μ . Paraphyses hyalinae circ. 2 μ crassae. Sporae 1-septatae, leniter constrictae, ellipsoidae, 17–20 $\times$ 6–7 μ , in maturitate brunneae, loculo superiore latiore.

Hab. in foliis *Trichoclaedi elliptici*, Pirie Forest, 8.7.19, leg. Doidge [12301].

78. *Trichopeltula Carissae* n. sp.

On leaves of *Carissa arduina*, East London, 24.11.17, Doidge [10914]; 19.7.19 [12424].

Epiphyllous, when thinly sown each thallus is plainly visible, consisting of a number of narrow, dull-black, irregularly branched radiating fibrillae; frequently the thalli are so numerous that they coalesce, and the whole leaf surface becomes encrusted with the dull-black growth; thallus irregular in breadth, 95–275 μ broad, formed of narrow, rectangular, radiating-prosenchymatous cells. Conidial pycnothecia, 90–115 μ diameter, containing bacilloform, hyaline, non-septate conidia, 6.8 μ $\times$ 1–1.5 μ . Pycnothecia containing asci somewhat larger, 120–140 μ diameter: asci paraphysate, eight-spored, obovate or obclavate, broadly rounded at the apex, constricted at the base, with a very short foot, 24–30 $\times$ 13–16 μ . Spores tristicous, hyaline, oblong, slightly curved, four to five celled, usually four-septate when mature, 10–14 $\times$ 3–4 μ .

Trichopeltula Carissae Doidge n. sp.

Epiphylla, thallus vittaeformis, repens, ramosus; vittae brunneo-atrae, irregulares, 95–275 μ latae, radiato-prosenchymaticae, cellulis anguste rectangularibus. Pycnothecia in thallo formata, alia conidiofera, 90–115 μ , alia ascofera 120–140 μ diametro, applanato hemisphaerica, demum apice perforata. Conidia bacillaria, hyalina, continua, 6–8 $\times$ 1–1½ μ . Asci paraphysati, octospori, obovati v. obclavati, supra late rotundati infra in nodum brevissimum coarctati, 24–30 $\times$ 13–16 μ . Sporae tristichae, hyalinae, oblongae, subcurvatae, quadrisepatae, 10–14 $\times$ 3–4 μ .

Hab. in foliis *Carissae arduinae*, East London, 24.11.17, leg. Doidge [10914].

79. *Trichopeltula kentaniensis* n. sp.

On *Xymalos monospora*, Kentani, 3.3.15, Pegler (Pegler No. 1999) [8866]; Zwartkop, near Maritzburg, Natal, 19.7.18, Doidge [11598].

On *Schmidelia monophylla*, Kentani, 3.3.15, Pegler (Pegler No. 2300) [8865].

This fungus is very similar in appearance to the species on *Carissa*, from which it differs chiefly in the size and form on the asci and spores; the latter are three to four septate, but usually four-septate when mature.

Trichopeltula kentaniensis n. sp.

Epiphylla, thallus vittaeformis, repens, ramosus; vittae brunneo-atrae, irregulares, 64–200 μ latae, radiato-prosenchymaticae, cellulis anguste rectangularibus. Pycnothecia in thallo formata, alia conidiofera 90–110 μ , alia ascofera 160–180 μ diametro, applanato hemisphaerica, demum apice perforata. Conidia bacillaria, hyalina, continua, 4–6 $\times$ 1–1½ μ . Asci paraphysati, octospori, ellipsoidei v. subclavati, apedicellati, 30–36 $\times$ 13–16 μ . Sporae tristichae, hyalinae, subclavatae, rectae, quadrisepatae, 15–17 $\times$ 3.5–5 μ .

Hab. in foliis *Xymalos monosporae*, Kentani, 3.3.15, leg. Pegler [8866].

Fungi belonging to this group appear to be very common in many localities, particularly in the neighbourhood of East London: but, although material has been collected at different seasons of the year, only the two species described above have been found with mature asci. Immature fungi belonging to the *Trichopeltaceae* have been collected near East London and elsewhere on the following hosts: *Acokanthera spectabilis*, *Behnia reticulata*, *Buxus MacOwani*, *Chaetacme aristata*, *Elaeodendron capense*, *Elaeodendron croceum*, *Eugenia* sp., *Putterlickia pyracantha*, *Rhoicissus rhomboidea*, *Toddalia natalensis*, *Trichocladus ellipticus*, *Umtisa Listeriana*.

80. *Scoleopeltis Cassipoureae* n. sp.

Ascomata hypophyllous, not situated on leaf spots, scattered, superficial, easily becoming detached, opaque, dark-green in the centre, translucent, paler at edges, 550–650 μ diameter, composed of a dense web of fine hyphae 1–2 μ thick.

Ostiole distinct, round 23–36 μ diameter. Asci paraphysate, ellipsoid, sessile, 120–170 $\times$ 20–26 μ , four to eight spored. Spores parallel or twisted, filiform, 100–120 μ long, six to seven broad at centre, tapering somewhat towards rounded ends, ten to fifteen septate, constricted, easily separating even in ascus into very uneven portions 6–15 μ long.

On leaves of *Cassipourea verticillata*, Buccleuch, Natal, 17.7.18, Doidge [11586].

Scoleopeltis Cassipoureae Doidge, nov. sp.

Peritheciis hypophyllis, sine maculis, sparsis, superficialibus, facile secedentibus, opae atro-coeruleis, lenticulari-scutatis margine alatis, 550–650 μ diametro marginem versus coerule pellucidi, ibique ex hyphis 1–2 μ latis tenuibus. maeandrice denseque conjunctis contextis, praeterea margine ca. 30–60 μ latoferre hyalino ex hyphis 1–2 μ latis anastomosantibus composito cinctis, ostiolo distincto, rotundato, 33–36 μ lato: ascis paraphysatis, ellipsoideis, sessilibus, 120–170 $\times$ 20–26 μ . 2–8 sporis, sporidiis parallelis v. tortis, filiformis 100–120 μ $\times$ 6–7 μ , utrinque leniter attenuatis, 10–15 septatis, constrictis, facile secedentibus.

Hab. in foliis *Cassipoureae verticillatae*, Buccleuch, Natal, 17.7.18, leg. Doidge [11586].

81. *Asterinella Mimusopsidis* n. sp.

On leaves of *Mimusops obovata*, Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12267]; East London, 19.7.19, Doidge [12409].

Hypophyllous, forming thin black spots up to 5 mm. diameter; mycelium reticulate, alternate or opposite, branching, composed of fuscous-brown hyphae, more or less torulose, and undulate, varying in thickness from 3–6 μ , articulation more or less irregular; hyphopodia none; thyriothechia crowded in centre, or scattered, flattened-hemispherical, 240–320 μ diameter, dehiscing by collapse of central cells or by radial fissure, composed of straight radiating hyphae 3–3.5 μ thick, margin entire or with numerous rather delicate hyphae about 3 μ thick running out into the mycelium. Asci paraphysate, ovate or narrow ovate, sessile, eight-spored, thickened round apex, 60–70 $\times$ 25–35 μ . Spores conglobate, two-celled, brown, constricted at septum, 28–33 $\times$ 13–16.5 μ , slightly rough at maturity, upper loculus broader, broadly rounded at both ends.

Asterinella Mimusopsidis Doidge, nov. sp.

Hypophylla, maculas tenues usque 5 mm. latas formans, mycelium ex hyphis fusciculis, subtorulosis, undulatis 3–6 μ crassis formatum. Hyphopodia desunt. Thyriothechia conferta v. sparsa, orbicularia, 240–320 μ diametro, radiatim ex hyphis rectis 3–3.5 μ crassis contexta, peripherice hyphis longiusculis, flexuosis, 3 μ crassis fimbriata. Asci paraphysati, ovati, sessiles, octospori, ad apicem incrassati, 60–70 $\times$ 25–35 μ . Sporae conglobatae, 1-septatae, brunneae, constrictae, 28–33 $\times$ 13–16.5 μ , in maturitate minute verrucosae, loculo superiore latiore.

Hab. in foliis *Mimusopsidis obovatae*, Pirie Forest, 8.9.19, leg. Doidge [12267].

82. *Meliola gemellipoda* n. sp.

On leaves of *Jasminum angulare*, Langholm Estates, Bathurst District, 14.7.19, Doidge [12352].

Epiphyllous, forming round black velvety spots 1–5 mm. diameter. Mycelium compact, fairly dense, formed of straight or slightly wavy hyphae 10 μ thick; cells 20–30 μ long; branching opposite or irregular; capitate hyphopodia opposite, usually one pair to each cell of hypha, 16–23 $\times$ 9–10 μ , stipe cell short, 3–5 μ , terminal cell ovate to cylindrical, rounded at the apex. Murconate hyphopodia rare, interspersed with capitate hyphopodia, ampulliform 20–30 μ long $\times$ 10 μ thick at base. Mycelial setae fairly numerous, simple, straight, black, opaque except near the somewhat blunt apex, 9–10 μ thick at base, tapering gradually towards apex, 360–600 μ long. Perithecia 200–250 μ diameter, black, globose, smooth. Asci two to three spored, evanescent. Spores four-septate, cylindrical or tapering slightly towards both ends, rounded at both ends, scarcely constricted, 53–60 $\times$ 16.5–20 μ . Closely related to *M. Toddaliae*.

Meliola gemellipoda Doidge, nov. sp.

Epiphylla, maculas atrae, orbiculares, 1–5 mm. diametro efformans; mycelio ex hyphis ramosis, septatis, 10 μ crassis composito; hyphopodiis capitatis oppositis 16–23 $\times$ 9–10 μ , cellula superiore ovata v. cylindrica; hyphopodiis mucronatis raris, ampulliformibus usque ad 30 μ longis; setis myceliis numerosis, simplicibus, rectis, opacis, 360–600 μ longis, obtusis; peritheciis 200–250 μ diametro, atris, laevis, globosis; ascis, 2–3 sporis, evanescentibus; sporidiis 4-septatis, ad septa vix constrictis, 53–60 $\times$ 16.5–20 μ .

Hab. in foliis *Jasminii angularis*, Bathurst District, Cape Province, 14.7.19, leg. Doidge [12352].

83. *Irene Rinorea* n. sp.

On leaves of *Rinorea ardesiaeflora*, Karkloof, Natal, 14.7.21, Doidge [14961].

Amphigenous, forming small, round, dull-black spots 1–2 mm. diameter: these are occasionally rather numerous and become confluent, but usually they are scattered and only one to two spots occur on a leaf.

Mycelium reticulate: hyphae dark-brown, semi-opaque, 6-7 μ thick, cells mostly 18-20 μ long, branches opposite, anastomosing, capitate hyphopodia alternate, usually one to each cell, stipitate, 17-26 μ high, terminal cell variously lobed with rounded or truncate lobes 17-20 μ broad. Mucronate hyphopodia rather numerous, interspersed with the capitate hyphopodia, opposite, alternate or unilateral: they are rather peculiar in form, reminding one of penguins at rest, 17-20 μ long and 5-6 μ thick at the base; mycelial setae none.

Perithecia spherical, black, carbonaceous, 140-200 μ diameter, surface cells conical or rounded, projecting to a height of 16-18 μ . (These could hardly be termed appendages; and the species is accordingly assigned to the genus *Irene*.) Asci fugaceous, not seen. Spores four-septate, cylindrical, rounded at both ends, constricted, $40-46 \times 16.5-21.5 \mu$.

Irene Rinoreae Doidge, nov. sp.

Amphigena, plagulas minutae, atrae, 1-2 mm. diametro efformans. Mycelio reticulato. Hyphis brunneis, subopacis, 6-7 μ crassis, cellulis plerumque 18-20 μ longis, hyphopodiis capitatis alternis, stipitatis 17-26 μ altis, cellula superiore varie lobata, lobis rotundatis v. truncatis. Hyphopodiis mucronatis numerosis, oppositis, alternis v. unilateralibus, 17-20 μ longis, basi 5-6 μ crassis.

Setis myceliis nullis. Peritheciis globosis atris, tuberculatis, 140-200 μ diametro. Ascis non visis. Sporidiis 4-septatis, cylindricis, utrinque rotundatis, ad septa constrictis, $40-46 \times 16.5-21.5 \mu$.

Hab. in foliis *Rinoreae ardesiaeeflorae*, Karkloof, Natal, 14.7.21, leg. Doidge [14961].

84. *Phyllachora myrsinicola* n. sp.

On leaves of *Myrsine melanophleos*, Duncairn, near Maritzburg, 13.7.21, Doidge [15015].

Stromata epiphyllous, not visible on lower surface of leaf, and not surrounded by discoloured leaf tissue, scattered, $\frac{1}{2}$ -1 mm. diameter, black, somewhat shiny, flat, very slightly raised above the epidermis.

Several loculi in each stroma, occupying about half the thickness of the leaf, which is here swollen to about twice its normal thickness. Loculi covered by a common epidermal clypeus which is black, opaque, and variable in thickness. Loculi not closely crowded, subspherical, 180-240 μ diameter, lateral and basal walls well developed, consisting of laterally compressed hyphae, usually hyaline, but sometimes brown at the base and where the lateral walls join the clypeus. Asci clavate $80-100 \times 20-27 \mu$, thick-walled, rounded at the apex, eight-spored, paraphysate, briefly pedicellate. Spores distichous, one-celled, hyaline, ellipsoid or rhomboid-ellipsoid, $16.5-23.5 \times 9-10 \mu$.

Phyllachora myrsinicola Doidge, nov. sp.

Stromata epiphylla, atra, in inferiore foliorum pagina non visibilia, sparsa, $\frac{1}{2}$ -1 mm. diametro parum nitidula, loculi pauci in quoque stromate, suborbiculares, 180-240 μ diametro, clypeo epidermale atro, opaco. Asci paraphysati, clavati, octospori, breviter pedicellati, $80-100 \times 20-27 \mu$. Sporae distichae, continuae, hyalinae, ellipsoidae v. subrhomboidae, $17.5-23.5 \times 9-10 \mu$.

Hab. in foliis *Myrsine melanophleos*, Duncairn, near Maritzburg, Natal, 13.7.21, leg. Doidge [15015].

85. *Clypeosphaeria natalensis* n. sp.

On leaves of *Viscum nervosum* growing on *Rapanea melanophleos*, Glen Burnie, Highflats, Ixopo District, Natal, 20.8.21, J. W. Bews [14867].

Perithecia and pycnidia epiphyllous, in scattered groups, similar in form and size, immersed, lenticular, spherical, convex at the apex with a black, opaque epidermal clypeus; 160-200 μ in diameter, very slightly papillate in the centre, where it becomes perforated by a pore; wall black, carbonaceous, 16-20 μ thick.

Asci paraphysate, subpyriform, broadly rounded at the base, narrower at the apex, thick-walled around the apex, which is traversed by a pore, eight-spored, evanescent, $90-120 \times 20-27 \mu$. Spores fuscous, four to five septate, cylindrical or subclavate, parallel or obliquely distichous, $42-54 \times 7.5-10 \mu$.

Conidia hyaline, cylindrical or clavate, rounded at both ends, transversely six to eight septate, $50-62 \times 10-11 \mu$.

Clypeosphaeria natalensis Doidge, nov. sp.

Peritheciis et pycnidiis epiphyllis, gregariis, immersis, lenticulariter sphaeroideis, vertice applanato, convexulo, centro leniter papillulatis demum poro perforatis, atris, $160-200 \mu$ diametro; ascis octosporis, paraphysatis, subpyriformibus, basi late rotundatis, sessilibus, apice incrassatis poro pertusis, evanescentibus, $90-120 \times 20-27 \mu$. Sporidiis fuscis oblique; distichis v. parallelibus, 4-5 septatis, cylindricis, v. subclavatis, $42-54 \times 7.5-10 \mu$.

Conidiis hyalinis, cylindricis v. clavatis, utrinque rotundatis, 6-8 septatis, $50-62 \times 10-11 \mu$.

Hab. in foliis *Visci nervosi*, Ixopo District, Natal, 20.8.21, leg. Bews [14867].

EXPLANATION OF ILLUSTRATIONS.

- Fig. 1. *Meliola Carissae*. (a) Mycelium with capitate hyphopodia. (b) Spores. (c) Mycelial setae.
 Fig. 2. *Meliola capensis*. Tips of mycelial setae.
 Fig. 3. *Meliola dracoenicola*. (a) Mycelium. (b) Spores. (c) Tips of mycelial setae.
 Fig. 4. *Meliola Oleicola*. (a) Mycelium. (b) Spores. (c) Tips of setae.
 Fig. 5. *Meliola Panici*. (a) Mycelium. (b) Spores. (c) Tips of setae.
 Fig. 6. *Irene Zeyheri*. (a) Mycelium. (b) Spores.
 Fig. 7. Mycelium and spores of *Asterina crotonicola*.
 Fig. 8. Mycelium of *Asterina Scolopiae*.

THE THORN PEARS

(*Scolopia* spp.).

By E. P. PHILLIPS, M.A., D.Sc., F.L.S., Division of Botany, Pretoria.

THE genus *Scolopia* was founded by Schreber in 1789, and the following year Loudon established the genus *Phoberos* for the same group of plants. It is under the latter name that the species are described in the "Flora Capensis." The genus is included by Bentham and Hooker under the natural order *Bixineae*, but Engler only includes the genus *Bixa* in this family and places *Scolopia* in the family *Flacourtiaceae*. Gilg in Engler's "Botanische Jahrbüch," vol. 40, p. 481, gives a list of the African species, and describes a new species from the Transvaal, viz. *Scolopia Engleri*. Sim ("Forests and Forest Flora," p. 126) recognized three South African species only, viz. *S. Mundii*, *S. Zeyheri*, and *S. Flanaganii*, and regards *S. Ecklonii* and *E. Gerrardii* as co-specific with *S. Zeyheri*. My examination of herbarium material has led me to recognize *S. Zeyheri* and *S. Ecklonii* as specifically distinct, and to regard *S. Gerrardii* and *S. Engleri* as varieties of *S. Ecklonii*. A very distinct species of the genus, collected at Barberton by Mr. G. Thorncroft, is described here.

The presence or absence of thorns is evidently not a constant character as Sim notes, as specimens sent in by Mr. A. O'Connor from the Woodbush Forest, Transvaal, are stated to be sometimes thorny and sometimes thornless. Though this may be the case, yet my examination of the genus does not confirm Sim's suggestion that *S. Zeyheri* and *S. Ecklonii* are the same species. Both the character of the leaves and the presence or absence of hairs on the inflorescence can be used to distinguish them. In herbarium material thorns were present on some specimens of *S. Zeyheri*, and it is only from information supplied by Mr. O'Connor that we can be certain of the presence of thorns in *S. Ecklonii*, as this character was not observed in any of the herbarium specimens. Mr. O. B. Miller, commenting on specimens collected by him in the Amanzamnyama and Buffalo Neck Forests, Mount Frere District, writes: "Yesterday I found a very large old 'red pear' with several sucker shoots growing from the stem at a height of 3 ft. 6 in. from ground-level. So far as I could see the rest of the tree was quite unarmed. It is the first time I have ever observed a thorned tree of this species, except in the seedling stage, when it is always armed. I have since seen another 'red pear' with armed suckers, and the native guard says they are fairly common in this district."

I have not seen authentic specimens of *S. Engleri*, but have examined Transvaal plants which agree with Gilg's description of this species. In some respects it appears to be an intermediate form between *S. Zeyheri* and *S. Ecklonii*, having the pubescent inflorescence of the former and the leaves of the latter, but the specimens are so similar to *S. Ecklonii* and the distribution limited to the bushveld that I prefer keeping it as a variety of this species. I have also decided to place *S. Gerrardii* as a variety of *S. Ecklonii*, as though while the specimens hitherto placed under the former species have a distinctive appearance owing to the size of the leaves, yet the examination of a large series of specimens of *S. Ecklonii* indicated that this species merges into *S. Gerrardii*.

I have to thank the Chief Conservator of Forests, Dr. Schönland, the Directors of the Transvaal and South African Museums, for the loan of the material in their respective herbaria.

KEY TO SPECIES.

Inflorescence a dense cylindric spike about 1 cm. in diameter.....	<i>Thorncroftii</i> .
Inflorescence a lax raceme.	
Leaves small, usually about 2 cm. long and 1 cm. broad; fruit minutely pubescent	<i>Flanaganii</i> .
Leaves usually over 2 cm. long and 1 cm. broad; fruit glabrous.	
Inflorescence pubescent.	
Leaves usually ovate in outline.....	<i>Zeyheri</i> .
Leaves usually obovate or oblanceolate in outline (Transvaal species)	<i>Ecklonii</i> , var. <i>Engleri</i> .
Inflorescence glabrous.	
Leaves usually abruptly acuminate and distinctly serrated.....	<i>Mundii</i> .
Leaves not abruptly acuminate, entire or sometimes remotely serrated.....	<i>Ecklonii</i> .

S. Thorncroftii, Phillips.

Arbor 6–7 mm. altus. *Rami* glabri. *Folia* 5·5–9·5 cm. longa, lanceolata vel obovata, aliquando subacuminata, apice obtusa, glabra. *Inflorescentia* densa, cylindrica, circa 1 cm. lata, pubescentia. *Petala* 1·25 mm. longa, linearia, apice acuta. *Filamenta* 6 mm. longa. *Ovarium* 1·5 mm. latum, globosum; stylus 2 mm. longus, apice paullo bifidus.

TRANSVAAL.—Barborton Div.: Barborton, 2900 ft., flowers white, *Thorncroft* 811, and in *Natal Herbarium* 14245.

Tree 20 ft. high. *Branches* glabrous, with the epidermis peeling off in membranous flakes. *Leaves* petioled; petioles 1–1·5 cm. long; lamina 4·5–8 cm. long, lanceolate or obovate, sometimes subacuminate, obtuse, entire, sometimes with wavy margins and with the veining distinct above and beneath, glabrous. *Inflorescence* in dense cylindric axillary spikes about 1 cm. in diameter. *Axis* pubescent. *Flowers* subsessile. *Sepals* 1·5 mm. long, ovate, obtuse, ciliate, pubescent. *Petals* 1·25 mm. long, linear, acute. *Glands* 0·5 mm. long, arranged in a continuous row on the receptacle. *Stamens* numerous; filaments 6 mm. long, surrounded at the base with long hairs; anthers ovate in outline, somewhat apiculate. *Ovary* 1·5 mm. in diameter, globose; style 2 mm. long, subterete, faintly bilobed at the apex.

This is a very distinct species and easily recognized by the dense cylindric inflorescence.

S. Flanaganii (Sim, "Forest and Forest Flora," p. 127, Pl. III.)

A small tree. *Branches* glabrous with raised lenticels. *Leaves* subsessile, 1·1–3 cm. long, 0·5–1·5 cm. broad, elliptic, rarely obovate, obtuse, sparsely serrated on the upper half, with distinct veining, glabrous. *Flowers* not seen. *Fruit* 0·5 mm. in diameter, subglobose, finely pubescent.—*Xylosma Flanaganii*, Bolus, in "Journ. Bot.," Jan., 1896¹.

CAPE PROVINCE.—Kingwilliamstown Div.: Kingwilliamstown, *Sim* 2571, and in *Natal Govt. Herb.* 8616.

This species is easily distinguished from the rest of the genus by the sessile leaves and the pubescent fruits.

S. Zeyheri [Szyszy in Pl. Rehmann (1887), 19].

A thorny shrub 8–10 ft. high (ex *Galpin*). *Youngest branches* minutely pubescent or glabrous, sometimes armed with spines 1·5–17 cm. long. *Leaves* petioled; petioles 0·5–1·5 cm. long; lamina 1·3–7 cm. long (usually 2·5–4 cm. long), 1–5 cm. broad (usually 2–3 cm. broad), ovate, ovate-elliptic, elliptic, elliptic-lanceolate, or obovate, more rarely subrotund or orbiculate, obtuse or rounded at the apex, more rarely subacuminate or retuse, rounded or cuneate at the base, with entire margins, more rarely subcrenate, with the veining distinct beneath, glabrous. *Inflorescence* an axillary raceme, 2·8–6·5 cm. long, usually lax, but sometimes more or less dense, though never broadly cylindric. *Axis* of inflorescence and pedicels finely but densely pubescent. *Sepals* 0·75–2 mm. long, 0·75–1·5 mm. broad, ovate, oblong, elliptic or suborbicular, obtuse or acute

sometimes subacuminate, usually shortly ciliate, pubescent without. *Petals* when present 1 mm. long, 0·75 mm. broad, ovate. *Glands* 0·25–0·5 mm. long, arranged in a continuous row on the receptacle. *Stamens* numerous; filaments 1·5–3 mm. long, surrounded at the base by long hairs; anthers 0·5–1 mm. long, ovate, or suborbicular in outline. *Ovary* 0·5–1·25 mm. in diameter, globose or ellipsoid, glabrous; style 1·5–2 mm. long, subterete, faintly bilobed at the apex (sometimes each lobe again faintly lobed). *Fruit* (mature?) 5 mm. in diameter, globose, glabrous, tipped with the persistent style.—*Sim*, "*Forest and Forest Flora*," p. 126, *pl. ii*; *Gilg in Engl. "Bot. Jahrb."*, 40, p. 481; *Phoberos Zeyheri*, *Arn.*, "*Flora Capensis*," I, 68.

CAPE PROVINCE.—Bathurst Div.: Port Alfred, *Schlechter* 2733. Stutterheim Div. Katberg, *MacOwan in Natal Govt. Herb.* 1113. Komgha Div.: In woods near Komgha, 2000 ft., *Flanagan*, 863. East London Div.: East London, *Wood in Herb. Galpin* 2828. Somerset East Div.: Zuurborg, *Holland* 279; banks of Klein Visch River, near Somerset East, *MacOwan* 808; woods on slopes of Boschberg, 3000 ft., *MacOwan* 1324. Queenstown Div.: Ridges at Finchams Nek, 4000 ft., *Galpin* 1822.

EAST GRICUALAND.—Maclear Div.: In wooded kloofs, Pot River Berg, 5500 ft., *Galpin* 6830.

NATAL.—Near Sinkwasi River, 500 ft., wood, Lundies Hill, *Wood* 10170, and *in Natal Govt. Herb.* 10780.

S. Mundii (Warb. in Engl. and Prantl. *Pflanzenfam.* iii, 6a, 29).

Branches glabrous. *Leaves* petioled; petioles 0·5–1 cm. long; lamina 1·5–7·5 cm. long, 0·7–4 cm. broad (usually 4–5 cm. long and about 2 cm. broad), abruptly acuminate, rarely not acuminate, acute or subacute, more rarely rounded or obtuse, cuneate at the base, with sharply serrated margins, veining distinct beneath, glabrous. *Inflorescence* a lax axillary raceme, 2–3 cm. long. *Axis* of inflorescence and pedicels quite glabrous. *Sepals* 1·5–2·5 mm. long, 1·5–2·5 mm. broad, ovate or subrotund, sometimes subacuminate, acute, obtuse or sometimes rounded at the apex, glabrous. *Glands* 0·5–1 mm. long, usually quadrate in outline, somewhat fleshy and arranged in a continuous row on the receptacle. *Stamens* numerous; filaments 3·5–5 mm. long, surrounded at the base with long hairs; anthers 0·75–1 mm. long, oblong, elliptic or orbicular in outline, usually apiculate at the apex. *Ovary* 1–3 mm. in diameter, globose or ellipsoid, glabrous; style 2·5–3 mm. long, subterete, divided at the apex into two short lobes, sometimes each lobe again faintly lobed. *Fruit* about 1 cm. in diameter, globose, glabrous.—*Sim*, "*Forests and Forest Flora*," p. 126, *pl. i*; *Gilg in Engl. "Bot. Jahrb."*, 40, p. 481; *Phoberos Mundtii*, *Arn.*, "*Flora Capensis*," I, 68.

CAPE PROVINCE.—Cape Div.: Kirstenbosch, in wood, Table Mountain, *Zeyher*. Knysna Div.: Grootkops River, Concordia, *Keet in Herb. Forest Dept.* 2373. Port Elizabeth Div.: Van Staadens, *Zeyher* 3785. Albany Div.: Woods about Grahamstown, 2000 ft., *MacOwan* 957, and *in Natal Govt. Herb.* 226. Komgha Div.: In woods at Prospect Farm Komgha, 2100 ft., *Flanagan* 62. Queenstown Div.: Lesseyton Nek, near Queenstown, 4000 ft., *Galpin* 2146; mountain top, Gwatyn Farm, 5400 ft., *Galpin* 8295.

TRANSKEI.—Tembuland, towards reservoir, near Cala, *Royffe* 204; Pondoland, Engeobo Div.: Manina Forest, *Zahn in Herb. Forest Dept.* 2040; Amamzamyama Forest, Mount Frere Dist., *Miller in Herb. Forest Dept.*, 3208, 3223.

NATAL.—Polela Div.: Polela, Drakensbergen, 6000–7000 ft., *Evans* 624, and *in Natal Govt. Herb.* 7407.

ZULULAND.—Qudeni Forest, 6000 ft., *Davis* 122, and *in Natal Govt. Herb.* 8807.

TRANSVAAL.—Eimelo Div.: Nooitgedacht, *Pott* 5097, and *in Herb. Transvaal Mus.* 14932.

BASUTOLAND.—Thaba-Bosin Div. : Banks of Mohale River, *Jacottet in Herb. Dieterlen* 1062. Maseru Div. : Matukeng, slopes of Qeme Mountain, *Dieterlen* 852

A tree 15–20 ft. high (*Evans*). A tree 10 ft. high ; drupe yellow (*Galpin*) ; Sesuto name “Moopa-Koma.”

S. Ecklonii [Szyszy, Pl. Rehmann (1887)].

Branches glabrous with raised lenticels. *Leaves* petioled ; petioles 0·5–1 cm. long ; lamina 2·5–11 cm. long, 0·7–3·6 cm. broad, mostly elliptic-lanceolate, more rarely lanceolate, oblanceolate or obovate, rounded or obtuse at the apex, more rarely subacute, cuneate at the base, with entire margins, rarely with a few teeth, with distinct veining, glabrous. *Inflorescence* a lax axillary raceme. *Axis* and pedicels quite glabrous. *Sepals* 1–2 mm. long, 0·75–1 mm. broad, ovate or oblong, obtuse, acute, or subacute, labrous. *Petals* when present 1–1·5 mm. long, linear or oblong. *Glands* forming a rim on the receptacle. *Stamens* numerous ; filaments 2–5 mm. long, surrounded at the base with long hairs ; anthers oblong or ovate in outline. *Ovary* 1·5 mm. in diameter, subglobose ; style 1·5–3 mm. long, faintly or distinctly 2–4-lobed at the apex. *Fruit* 4·5 mm. in diameter, globose.—*Sim.* “*Forests and Forest Flora*,” p. 127 ; *Gilg in Engl.* “*Bot. Jahrb.*,” 40, p. 481 ; *Phoberos Ecklonii*, Arn., “*Flora Capensis*,” I, 68.

CAPE PROVINCE.—Uitenhage Div. : Van Staadens Mountains, *Zeyher* 782. Albany Div. : In woods near Grahamstown, *MacOwan* 1324.

NATAL.—Near Tugela River, 500 ft., *Wood* 3951, and in *Natal Govt. Herb.* 4578.

ZULULAND.—Qudeni Forest, 6000 ft., *Davis* 99, and in *Natal Govt. Herb.* 8806 ; swamp, Umhlatuzi Forest, *Ballenden in Herb. Forest Dept.* 2559.

TRANSVAAL.—Pietersburg Div. : Potatobosch, Woodbush Forest, 4900 ft., *Botha in Herb. Forest Dept.* 2132 ; *O'Connor in Herb. Forest Dept.* 2009.

Var. **Gerrardii**, Phillips.

Leaves up to 8 cm. long and 5 cm. broad, with usually undulate margins and sometimes remotely serrated.—*Scolopia Gerrardii*, *Harv.*, “*Flora Capensis*,” II, 548.

NATAL.—Westville, near Durban, 4000–5000 ft., *Wood* 8597, and in *Natal Govt. Herb.* 9320 ; Sydenham, 300–400 ft., *Wood* 11402.

PONDOLAND.—Engcobo Div. : Manina Forest, *Zahn in Herb. Forest Dept.* 2041.

Var. **Engleri**, Phillips.

Agrees with the type, but the inflorescence is slightly pubescent.—*S. Engleri*, *Gilg in Engl.* “*Bot. Jahrb.*,” 40, 481.

TRANSVAAL.—Derdepoort, *Miss Leendertz* 190. Rustenburg Div. : Rustenburg, *Miss Leendertz* 9773. Pretoria Div. : Betfontein, *Theiler in Herb. Transvaal Museum* 12365 ; Kaalplaats, *Mogg in Govt. Herb.* 16289.

THE GENUS OCHNA.

By E. P. PHILLIPS, M.A., D.Sc., F.L.S.

THE National Herbarium has on many occasions received specimens of *Ochna* from the Forest Department for identification, but found that the naming of the species in herbaria was unsatisfactory. Forest officers have suggested that more species are found in the South African forests than those generally accepted as occurring within the Union, and as some of the species are of economic value it was pointed out by the Chief Conservator of Forests that the genus should be critically examined. With the permission of the Chief of the Division of Botany the writer undertook this work. Without the cordial co-operation of the Forest Department this would not have been possible, and my appreciation is here expressed of the willing assistance received from Mr. J. J. Kotze, B.A., B.Sc., the acting Research Officer at the time. To the various forest officers who have supplied specimens and information the writer is also deeply indebted, especially to Mr. A. J. O'Connor, the District Forest Officer at Doornboom, northern Transvaal, who first pointed out the existence of two distinct species of *Ochna* which occur in the Woodbush Forest, and which had previously been regarded as either *O. arborea* or *O. atropurpurea*, var. *natalitia*. My thanks are also due to the Director of the South African Museum, who allowed me facilities for working at the Cape Government Herbarium, and to Dr. S. Schönland, Mrs. F. Bolus, B.A., Mrs. R. Potts, and Dr. P. J. van der Bijl for lending me the material in their respective herbaria, and to the Director of Kew for comparing some specimens with the Kew material.

At first sight the division of the genus into species appeared an easy matter, but as more material was examined the difficulties of limiting the species increased. The writer has critically examined nearly 200 herbarium sheets, but failed to find any constant characters, either floral or vegetative, which could be used as diagnostic characters for a key to the species, and has had to rely on the general "facies" of the specimens in most cases and has found it difficult to express the differences in words. That the specimens have been grouped correctly may be assumed, as handling a large number of specimens makes one familiar with their "look," but whether a particular specimen can be readily placed in its correct species with the aid of the key given here is more doubtful. However, this difficulty will only apply to a few species related to *O. atropurpurea*; the others are fairly distinct.

In the "Flora Capensis" (Vol. I, 448), three species of *Ochna* are described, but my investigations show that there are nine distinct species in South Africa with the possibility of a tenth. This latter is represented by two sheets illustrating a plant with a prostrate habit, but as Harvey, in the "Flora Capensis" under *O. arborea*, mentions that "shoots from the roots occasionally produce flowers," it is just possible that the specimens may represent such shoots, but no information on this point is available. In all the herbarium specimens examined only *O. pulchra* was correctly distinguished, while the limits of *O. arborea* and *O. atropurpurea* were confused. *O. pulchra* is a typical Transvaal species, but also occurs in Rhodesia, and as far as our records go is absent from the Cape Province and Natal. It is readily distinguished from all the other South African species by the large racemose inflorescence. Other distinguishing characters are the entire leaves (or they usually appear entire to the naked eye) and the long stigmas, which are one-third to a quarter the length of the style. The other species which has entire (or apparently so) leaves is *O. arborea*, but this does not occur in the Transvaal. It has been recorded from the George,

Kynsna, Uitenhage, Albany, Stockenstrom, East London, Kingwilliamstown, Komgha, and Kentani Divisions, and then appears in Natal and extends to Lourenco Marques. The inflorescence in this species is also racemose (a very abbreviated raceme), but specimens have passed through my hands in which the flowers are either solitary or 2-4-nate at the apex of short shoots; this is, however, very rare. The third species, *O. atropurpurea*, as I have limited it, is composed of plants of a shrubby habit, 2-3 ft. high, and can usually be recognised at a glance by the branches having a distinct lepidote appearance due to the presence of numerous prominent lenticels; the leaves are usually elliptic, sharply serrated, and the flowers solitary. It has a similar distribution to *O. arborea*, but has also been recorded from Zululand and Barberton. The variety *natalitia* has been kept as a distinct species; it differs from *O. atropurpurea* in habit, and the branches have not the lepidote appearance; the flowers are rarely solitary, and the leaves longer and lanceolate. Though Harvey recognized this variety, he has nevertheless confused it with the species, as in the Cape Government Herbarium there is a sheet (*E. and Z.* 925) labelled on the ticket *O. atropurpurea* and written up as such in Harvey's handwriting, which is undoubtedly *O. natalitia*. This species is not so common in the south-eastern districts as *O. atropurpurea*, but is recorded from several localities in Natal, and then passes through Swaziland to Barberton. Ecklon and Zeyher record it from the Uitenhage Division. The correct limitation of this species has proved the most troublesome, and at first I was inclined to recognize three species. A further examination proved that the first grouping of the specimens was very artificial and that this species should be regarded as a variable one with a wide range of distribution. *O. Chilversii* appears to be confined to the eastern forests, it resembles *O. natalitia* in general habit, but is readily distinguished from this species by the larger number of carpels in the ovary, and the flowers are usually much larger. The two species, *O. O'Connorii* and *O. Holstii*, which occur in the Woodbush Forest are very distinct from all the other South African species. Both are deciduous trees of timber size, and may perhaps be confused in herbaria if leaves are only present, but they differ considerably. *O. Holstii* has a rough bark and ellipsoid fruits, while *O. O'Connorii* has a smooth bark with kidney-shaped fruits. The remaining species, *O. leptoclada*, is found in the Barberton and Lydenberg Districts and also in Rhodesia. Specimens in mature foliage are easily recognized by the oblanceolate leaves, which are long attenuated at the base.

The African species of *Ochna* were monographed by Diels in 1904 (*Engl., "Bot. Jahrb."* Vol. XXXIII, p. 232), to which account the reader is referred for a key to all the known African species.

The writer was granted facilities by Mr. Kotze for extracting information about the species from notes sent to the Forest Department by various forest officers.

O. Chilversii. This species is known in the Cwebe Forest by the natives as "sibomyu." The Conservator of Forests, Transkeian Conservancy, remarks: "I was struck with the nice appearance of the wood. It is a soft (hardwood), easy to work. . . . It is plentiful in the Cwebe and other coast forests and approximates *arborea* in size, and could easily be exploited. Possibly it might make an excellent hammer handle." Mr C. C. Robertson remarks on the difference between this species and *O. arborea*. He reported that the "wood, which seems to me far from soft, appears to have the same general character as boxwood, and would probably be suitable for similar purposes," and he suggested that it might be tried for whatever purpose *O. arborea* is used for. The forester at Cwebe, who has handled the wood, describes it as "easy to work and soft."

In the Notinsela Forest the natives know the tree as "umbomvane umncince," and according to Mr. F. M. Fegen the tree occurs both in the Notinsela and Mpimbo Reserves, and appears to be plentiful in the latter locality, and no doubt also in the surrounding forests.

Forester Chilvers mentions that he has only seen one specimen in the Ingeli Forest Reserve. He describes it as a "small tree, stem 5 inches in diameter; bark not smooth as in *O. arborea*; leaves not deciduous."

O. Holstii. This species occurs in the Ingeli Forest Reserve and is also found in the Woodbush Forest in the northern Transvaal. The specimens in the Ingeli Reserve, Forester Chilvers describes as "trees of timber size with deciduous leaves; stem attaining to a diameter of 2 ft. to 2 ft. 6 in. with a rough bark." The species is quite plentiful at Ingeli and is known to the sawyers as the "real red pear" as distinguished from *Scolopia*. Forester Tustin describes this species as "a branched tree 40-50 ft. high, stem 18 in. in diameter, with a bole 15-20 ft. Bark grey on the outside and deep red inside. Flowers in September and October and fruits ripen in December. I have only found the tree growing in denser parts of the forest. It is found throughout my patrols (Impetynne and Ingeli) and is sparingly dotted through the forests. Saplings are numerous compared with old trees. I have not seen this tree used for any purpose."

District Forest Officer A. O'Connor, in submitting specimens from the Woodbush Forest, states it to be "a fairly large deciduous tree. The bark is comparatively smooth, though not with the hard smoothness of *O. O'Connorii*, being more flaky and 'chippy.' It is beet-red under the surface."

Mr. O. Miller (the acting District Forest Officer at Doornboom at the time) states that the species is fairly abundant in the lower, but scarce in the upper, slopes of the Woodbush Forest. It attains a height of 45 ft., with a trunk diameter, at breast high, of 30 in. The species flowers in November and sets its fruits in January.

O. O'Connorii. This tree has only been recorded from the Woodbush Forest, and was collected by Mr. A. J. O'Connor. He noted the differences between this and *O. arborea*, and describes it as a "small tree with smooth bark, very much like the ordinary Cape plane." The local name is "rooihout," "rooi izerhout."

O. arborea. This species may attain the size of a timber tree or may be only 8-10 ft. high. In the Horseshoe Forest, Kingwilliamstown, Forester Whithal records this species as a "tree 30 ft. high, with a diameter of 26 in. at breast high, and is found on a dolerite formation." At Knysna, Mr. J. D. Keet, the District Forest Officer, records it "as common in all parts of the district, being a small tree which forms part of the under storey of the forest, and rarely reaches a height of 1 ft. at breast high." Locally known as the "rooihout." The wood is very strong and of great durability, and is generally used for making handles of tools and axes. The seeds are said to be attacked by caterpillars. Forester Watton states it to be a plant growing in scrub at Knysna, and is known by the name of "rooihout." Forester Strauch sends the following notes with his specimens: "Very plentiful in all the reserves in this (Alexandria) forestal district, and is found throughout the forest. Average height, 10 ft.; average girth, 4 in. In the kloofs it grows much larger and it is found that the bark is much lighter than in trees growing on ridges or exposed positions. The timber is used for making axe and hammer handles. The tree appears to grow more or less plentifully over the forests in the Alexandria District, which are all on sandy soil."

Sim* is in error when he states that this species occurs in the Transvaal.

O. natalitia. This species, which has been regarded as a variety of *O. atropurpurea*, rightly deserves specific rank. Forester F. M. Fegen reports it as a "small tree about 6 ft. high growing at Notinsela Forest. It is very free in flowering and has scented flowers. I have seen several specimens of this tree in the Mpimbo Forest, but it does not appear to grow higher than 6 or 8 ft. Many young plants of *O. atropurpurea* are growing close to this." Forester Fegen further remarks that both species (*O. natalitia* and *O. atropurpurea*) grow on the edge of the forest and appear to be gregarious, growing thickly together. I have not found them well in the forest. Many stunted plants are in "dobo" grass, and have been scorched by fire and have thrown out branches from the base. This species should make a handsome garden plant, and fruiting specimens are particularly beautiful. The Pondos call it "mbomvane ncinci," but do not appear to use it for any purpose.

* Sim, "Forest Flora," p. 164.

O. atropurpurea. Forester Fegen records this as a "small shrub growing to edge of Maseko Forest and which is very plentiful. It has flowered and fruited very sparingly this year (1915) and fruiting specimens are scarce. The leaves, although identical in shape, are smaller on every specimen than those of *O. natalitia*. It is also found at Notinsela. Galls are nearly always present, and the plants, like *O. natalitia*, are confined to the edge of the forest."^{*}

O. pulchra. A small or large tree, 6-20 ft. high. The trunk is smooth and usually white, which makes the recognition of this tree in the veld easy. It is usually in flower late in September, and at the end of October the sepals take on a red colour which makes the species so conspicuous. The young fruits are then forming, and they mature in December. The flowers are sweetly scented; at first the sepals are bright green, then change to olive-brown and red. The seeds are said to be poisonous.

KEY TO SPECIES.

Inflorescence in definite long or short racemes, sometimes paniced, very rarely solitary or 2-3-nate at the apices of short shoots, but if so then the leaves entire or appearing entire to the naked eye.

- | | |
|--|---------------------|
| Ovary of 10-13 carpels..... | <i>Chilversii</i> . |
| Ovary of less than 10 carpels..... | |
| Leaves entire or appearing entire to the naked eye (rarely serrated, but if so, then never with prominent lenticels), not acuminate; bark smooth. | |
| Racemes 2-12 cm. long..... | <i>pulchra</i> . |
| Racemes much abbreviated..... | <i>arborea</i> . |
| Leaves serrated, long acuminate, branches not densely covered with prominent white lenticels; fruit ellipsoid..... | <i>Holstii</i> . |
| Leaves serrated, shortly acuminate or cuneate at the apex, rarely rounded; branches sometimes covered with prominent white lenticels; fruits ellipsoid or kidney-shaped. | |
| Fruits kidney-shaped; lenticels not numerous and prominent; a tall tree. | |
| Leaves obtuse; Transvaal species..... | <i>O'Connorii</i> . |
| Leaves acute; Eastern species..... | <i>arborea</i> . |
| Fruits ellipsoid, lenticels numerous and prominent; a large shrub 6-8 ft. high..... | <i>natalitia</i> . |

Inflorescence solitary or 2-4-nate arising from the apex of abbreviated shoots.

Small trees or erect shrubs.

- | | |
|--|-----------------------|
| Mature leaves up to 9 cm. long, oblanceolate, usually long-attenuate at the base..... | <i>leptoclada</i> . |
| Mature leaves very rarely up to 8 cm. long, rounded or narrowed at the base, never long-attenuate. | |
| Flowers 2-5-nate on much abbreviated shoots from the axils of the leaves..... | <i>natalitia</i> . |
| Flowers solitary, very rarely 2-5-nate, but then arising from the apex of short shoots and not in the axils of the leaves. | |
| Lenticels usually dense and prominent, giving the branches a lepidote appearance to the naked eye; teeth on the margins of the leaves usually triangular in shape, erect, and acute..... | <i>atropurpurea</i> . |
| Lenticels usually present, but branches not appearing lepidote to the naked eye; teeth on margins of leaves usually incurved. | |
| Epidermis flaking from the branches in membranous strips; leaves up to 4 cm. long.. | <i>pretoriensis</i> . |
| Epidermis rarely evidently flaking in membranous strips; leaves up to 8 cm. long.. | <i>natalitia</i> . |

O. Chilversii Phillips. *Arbor parva*. *Cortex* sulcata. *Folia*, 3-8 cm. longa, 6-2.5 cm. lata, lanceolata, apice obtusa, basi cuneata, glabra, marginibus serratis. *Inflorescentia* in racemo brevissime disposita. *Ovarium*, 10-13 carpellis.

^{*} Sim, "Forest Flora," p. 164.

A small tree with a trunk 5 in. in diameter. *Bark* rough. *Branchlets* with reddish-brown to almost black bark, pustulate or not, not flaking. *Leaves* 3-8 cm. long, '6-2.5 cm. broad, lanceolate, rarely more or less oblanceolate, obtuse, cuneate at the base and passing into a short petiole, rarely rounded, with the midrib distinct above and beneath and the lateral veins evident or not, with serrated margins, glabrous. *Inflorescence* on short shoots in abbreviated racemes, up to 7-flowered. *Pedicels* 1.7-2.2 cm. long, terete, articulated 1-3 mm. from the base, glabrous. *Sepals* 1.4 cm. long, '7 cm. broad, ovate, obtuse, and rounded at the apex; two of the sepals cucullate at the apex, glabrous (in dried specimens the folding of the sepals gives them the appearance of being acuminate). *Petals* 1.7 cm. long, 1.4 cm. broad, obovate, rounded at the apex, narrowed at the base into a short broad claw. *Filaments* 5.5 mm. long; anthers 3 mm. long, linear-oblong in outline, dehiscent by two apical slits. *Ovary* of 10-13 carpels, about 1 mm. long; style 1 cm. long, somewhat compressed; stigmas either capitate or saucer-shaped, with the stigmatic surfaces arranged round the rim. *Fruit* unknown.

EASTERN REGION.—Cwebe Forest, July, *Pretorius in Herb. Forest Dept.* 1625; Notinsela Forest, *Fegen in Herb. Forest Dept.* 1817; Ingeli Forest Reserve, December, *Chilvers in Herb. Forest Dept.* 1945; at mouth of Umkwani River, October, *Tyson*, 2619.

O. pulchra (Hook. fil., Ic. Pl. t. 588).

A small or large tree, 6-20 ft. high. *Branches* usually white or light grey to brown in colour, quite smooth or rough, with the epidermis peeling off in membranous flakes. *Leaves* subsessile, 3.5-11 cm. long, 1.5-5 cm. broad, usually elliptic, sometimes elliptic-lanceolate, rarely oblanceolate, rounded or obtuse at the apex, sometimes mucronate, usually narrowed at the base, more rarely rounded or subcordate; margins entire or appearing entire to the naked eye, more rarely with subulate teeth adpressed to the margins, with the mid-rib prominent beneath and the lateral veins distinct, glabrous. *Inflorescence* a many-flowered raceme, 2-12 cm. long. *Pedicels* 1-2.5 cm. long, glabrous, articulated at the base. *Sepals* 6-8 mm. long, 5 mm. broad, elliptic or suborbicular, obtuse, sometimes with cartilaginous teeth on the margins. *Petals* 7.5-9 mm. long, 4.5-5.5 mm. broad, obovate, narrowed at the base. *Filaments* 2 mm. long; anthers 1.5-2 mm. long, oblong or oblong-linear in outline. *Ovary* of 5-7 carpels; style 2 mm. long; stigmas as many as the carpels, about one-third to a quarter as long as the style, recurved, with faintly bilobed apices. *Fruit* 1.1-1.3 cm. long, 8 mm. in diameter, kidney-shaped, and attached to the torus at the middle.—" *Flora Capensis*," I, 449.

TRANSVAAL.—Northern Transvaal without precise locality, *Becker in Herb. Albany Mus.*; Pretoria District: Premier Mine, *Rogers* 19846; Magaliesberg, October, *Burke* 302, *Schlechter* 3623; Meintjes Kop, September, *Robertson in Herb. Forest Dept.* 1662; kopjes round Pretoria, September, *Miss Leendertz* 300, 381, May, *Burt-Davy* 2449; Koedoespoort, September, *Miss Stent in Goot Herb.* 10527; Wonderboom, October, *G.A.Z. in Herb. Forest Dept.* 1686, *Mogg in Herb. Forest Dept.* 1673 and in *Goot Herb.* 11509, *Galpin* 6977, *Burt-Davy* 2678 and in *Herb. Forest Dept.* 545, *Miss Leendertz* 431. Rustenburg Dist.: Plains at foot of Magaliesberg, *Miss Pegler* 1032; Wolluterskop, *Howlett in Goot Herb.* 16176, *Pole Evans in Goot Herb.* 15843; Silikats Nek, De Kroon, *Putterill in Goot Herb.* 16176; Rustenburg, *Miss Leendertz* 3369. Waterberg Dist.: Warmbaths, *Burt-Davy* 2626; Leeuwpoort Tin Mines, *Rogers* 19071, 21874; Pietpotgietersrust, *Burt-Davy* 2289, *Miss Leendertz* 5635, 2392, *Collector (?) in Herb. Bolus* 10a; Chunies Poort, *Pole Evans* 19460.

RHODESIA.—Veld, Victoria Falls, August-October, *Miss Gibbs* 305; sandy soil, Victoria Falls, 2000 ft., *Rogers*, 5301, 5302, 5423; in open forest, Victoria Falls, 2900 ft., *Galpin* 7050; Livingstone, 3000 ft., October, *Rogers*, 7450; Victoria, *Monro* 1594, 2108.

This species is usually in flower late in September, and at the end of October the sepals take on a red colour which makes the plant so conspicuous. The young fruits are then forming and they mature in December. The flowers are sweetly scented; at first the sepals are bright green, then change to olive-brown and red. The seeds are said to be poisonous.

O. arborea (Burch. ex D.C. Prodr. 1, 736).

A shrub 8 ft. to a tree 30–40 ft. high. *Bark* smooth. *Branches* with light or dark grey bark, with prominent lenticels and with the epidermis often peeling in membranous flakes. *Leaves* subsessile, 1·5–6·5 cm. long, 0·8–3 cm. broad, mostly elliptic more rarely lanceolate-elliptic or obovate, obtuse or rounded at the apex, rarely acute, narrowed or rounded at the base, quite entire or appearing so to the naked eye, very rarely distinctly serrated; teeth when present more or less subulate and adpressed to the margin of the leaf (except *Wood* 5878, in which the serrations are distinct). *Inflorescence* less than 1 cm. long, in a much abbreviated raceme of 3–7 flowers, very rarely solitary or 2–3-nate at the apex of short shoots. *Pedicels* 0·7–1·2 cm. long, articulated about 1 mm. from the base. *Sepals* 6·5 mm. long, 3·5–5 mm. broad, elliptic or ovate-elliptic, rounded at the apex. *Petals* 7·5–1·3 cm. long, 3–5·5 mm. broad, oblong or oblong-obovate, cuneate at the base. *Filaments* 2·5–3 mm. long; anthers 1·5 mm. long, oblong in outline. *Ovary* of 4–6 carpels; style 5·5–6 mm. long; stigmas as many as the carpels, or sometimes stigma subglobose. *Fruit* ellipsoid or almost kidney-shaped, attached to the torus at the middle.—“*Flora Capensis*,” I, 449; *Sim*, “*Forest and Forest Flora*,” 163 *pl. xxix*.

George Dist.: George, August, *Rogers* 4039; Woodville Forest, George, *Sim* in *Herb. Forest Dept.* 3116, 3117. Knysna Dist.: Knysna Commonage, *Keet* 498; Christies Kloof, Gouna, near Knysna, *Collector (?)* in *Herb. Forest Dept.* 1342; Gouna River, September, *Keet* in *Herb. Forest Dept.* 2529; Kaffirkop Forest, c. 1500 ft., January, *Watson* in *Herb. Forest Dept.* 2943; De Eiland, *Rabbets* in *National Herb.* 1420, 1421. Uitenhage Dist.: Oliphants Hoek and Addo, September–October, *E. and Z.* 924; Zuurberg, *Drège*. Alexandria Dist.: Forests near Alexandria, *Strauch* in *Herb. Forest Dept.* 3012, 3013, 3005. Albany Dist.: Blaauw Krantz Bridge, October, *Galpin* 265: without precise locality, *E. and Z.*; Bothas Hill, December, *Salisbury* 357; Salentown, *Zeyher*; near Grahamstown, *MacOwan* 110. Stockenstrom Dist.: Katberg *Mrs. Hobson* 1446. East London Dist.: Coast near East London, April, *Sim* 2153; East London, July, *Miss Gane* 260; near Nahoon River, August, *Galpin* 3291. Kingwilliamstown Dist.: Along the Buffalo River, near Kingwilliamstown, November, *Flanagan* 359; Horseshoe Forest, December, *Whithal* in *Herb. Forest Dept.* 2941; Mkobisa Forest, December, *Van der Merwe* in *Herb. Forest Dept.* 2939; Western Pirie Forest, January, *Kopke* in *Herb. Forest Dept.* 2940. Victoria East Dist.: Auckland Forest, January, *Ackermann* in *Herb. Forest Dept.* 2942. Komgha Dist.: Woods near Prospect Farm, November, *Flanagan* 359. Kentani Dist.: Forest near Kentani, July, *Miss Pegler* 1216. Pondoland, Mpingo Forest, *Fegen* in *Herb. Forest Dept.* 3033, 3089.

Natal: Port Natal, *Gueinzins*; Inanda, August, *Wood* 687, and in *Natal Govt. Herb.* 5878; without precise locality, *Gerrard and McKen* 834, and in *Natal Govt. Herb.* 73; Victoria County, *Wood*; Shafton, Howick, *Mrs. Hutton* 153.

Delagoa Bay: *E. and Z.* 926; *Madame Borl'* 176, 16; *Rogers* 21373.

A large shrub (*Wood*). A forest tree, 15 ft. high, with masses of yellow flowers (*Miss Pegler*). A small tree throughout the whole of the Knysna Forest; generally used for making handles of tools and axes; wood very strong and heavy and of great durability; seeds attacked by caterpillars (*Herb. Forest Dept.* 1342).

O. O'Connorii, Phillips. *Arbor. Cortex* plana. *Folia* 5–10 cm. longa, 1·3–5 cm. lata, lanceolata, raro elliptica vel ovato-lanceolata, apice paullo angustata, obtusa, *Inflorescentia* in racemo brevissime disposita; flores solitarii vel 2–3-nati. *Fructus* 1·5 cm. long, 8 cm. latus, reniformis.

A tree. *Bark* comparatively smooth, somewhat flaky and “chippy,” beet-red beneath the surface (*O'Connor*). *Branchlets* somewhat pustulate, reddish-brown to dark-brown in colour, not flaking. *Leaves* 5–10 cm. long (usually 6–8 cm. long), 1·3–5 cm. broad, lanceolate, rarely elliptic or ovate-lanceolate, slightly tapering at the apex, rarely acuminate or rounded and retuse, obtuse, sometimes slightly tapering to the base and rounded, with the mid-rib

prominent above and beneath and the lateral veins distinct above; glabrous. *Inflorescence* racemose with the flowers either solitary or 2-3-nate in the raceme. *Pedicels* 1-2·2 cm. long, articulated 1-4 mm. from the base. *Sepals* 7 mm. long, 5 mm. broad, broadly elliptic, rounded at the apex. *Petals* 1·1-1·2 cm. long, 5-6·5 mm. broad, obovate, rounded above, narrowed into a short broad claw at the base. *Filaments* 2·5 mm. long, anthers 2·5 mm. long, linear-oblong in outline. *Ovary* of 5-6 carpels; style 5·5 mm. long; stigmas as many as the carpels. *Fruit* 1·5 cm. long, 0·8 cm. broad, kidney-shaped, attached to the torus at the concave side.

TRANSVAAL.—Barberton Dist.: Van der Merwes Bush, Pilgrims Rest, 5-5750 ft., January, *Burt-Davy* 1409; Barberton, February, *Orange in Colonial Herb.* 1691; *Legat in Colonial Herb.* 1316. Pietersburg Dist.: Woodbush Forest, October, *Grenfell* 3, and in *Colonial Herb.* 1017; *Dryer*; December-January, *Botha in Herb. Forest Dept.* 2929, 2934; *Legat in Herb. Transvaal Mus.* 9711; by stream at Potatobush, c. 4750 ft., January, *Burt-Davy* 1193; *Eastwood in Colonial Herb.* 1286; Woodbush, 4900 ft. October, *O'Connor in Herb. Forest Dept.* 2198; *Lane-Poole in Herb. Forest Dept.* 547; *Grenfell in Colonial Herb.* 1112; without collector in *Herb. Forest Dept.* 542.

Known as the Cape Plane, Rooihout, Machailo, Rooi Yzerhout, Morelle.

O. Holstii (Engl. in Engl. Pflanzenw. Ostafri. c. 273).

A tree of timber size. *Bark* rough. *Branchlets* varying in colour from light to dark brown, scarcely pustulate, not flaking. *Leaves* deciduous, 5-8 cm. long, 0·8-2 cm. broad, lanceolate, acuminate, obtuse at the apex, narrowed at the base into a short petiole, mid-rib distinct above and beneath, lateral veins not evident, with the margins serrated and with the serrations very erect, glabrous. *Inflorescence* on short shoots in abbreviated racemes, up to 7-flowered. *Pedicels* 1·3-2·7 cm. long, terete, articulated 1-3 mm. from the base. *Sepals* 1 cm. long, lengthening in the fruit, 0·5 cm. broad, elliptic, rounded at the apex, glabrous (in dried specimens the folding of the sepals gives them the appearance of being acuminate). *Petals* 8 mm. long, 5 mm. broad, obovate, rounded at the apex, narrowed at the base into a short broad claw. *Filaments* 5 mm. long; anthers 2 mm. long, oblong in outline, dehiscing by apical slits. *Ovary* of 4-5 carpels about 1 mm. long; style 6 mm. long; stigma globose, faintly lobed. *Fruit* ellipsoid, attached to the torus at one end.

EASTERN REGION.—Tembuland, Engcobo Mtn., *Flanagan* 2696. Natal: Ngomi, *Tustin in Herb. Forest Dept.* 3115; Ingeli Forest Reserve, December-January, *Chilvers in Herb. Forest Dept.* 1944, 1946; *Tustin in Herb. Forest Dept.* 2907, 2984.

TRANSVAAL.—Pietersburg Div.: Woodbush, December, *Botha in Herb. Forest Dept.* 2935; 4900 ft., November, *O'Connor in Herb. Forest Dept.* 2197; without collector in *National Herb.* 1242. Lydenburg Div.: Pilgrims Rest, Van der Merwes Bush, 5400-5750 ft., January, *Burt-Davy* 1408; Drakensberg at Pilgrims Rest, *Collector (?) in Herb. Forest Dept.* 549. Barberton Div.: Barberton, *Rogers* 14996.

This tree flowers in November and sets its fruits in January. The sawyers call it the "real red pear" as distinct from *Scolopia*. In the Woodbush it is known as "rooi yzerhout."

O. natalitia (Engl. et Gilg. in Engl. "Bot. Jahrb.," 33, 236; name only).

Large shrub or small tree. *Branches* usually with prominent lenticels, often numerous but not appearing lepidote to the naked eye. *Leaves* sessile, 2-9 cm. long, 0·5-2·5 cm. broad, usually lanceolate, more rarely lanceolate-elliptic or lanceolate-linear, cuneate or slightly narrowed at the base, very rarely rounded at the base, obtuse or rounded at the apex, with serrated, very rarely entire margins, with the teeth usually incurved; the mid-rib distinct above and beneath and the lateral vein evident. *Inflorescence* either solitary or 2-nate from the apex of short shoots, or 2-5-nate arising in the axils of the leaves, very rarely in abbreviated racemes. *Pedicels* 0·4-1·5 cm. long, articulated at the base or

1 mm. from the base. *Sepals* 7–8 mm. long, 4·5–5·5 mm. broad, elliptic or rounded at the apex. *Petals* 0·9–1·2 cm. long, 6–7 mm. broad, obovate, narrowed at the base. *Filaments* 1·5–2 mm. long; anthers 1·5–2·5 mm. long, oblong in outline. *Ovary* of 5–9 carpels: style 4·5–6·5 mm. long; stigma either discoid or stigmas as many as the carpels. *Fruit* 0·8–1·4 cm. long, 5–9 mm. in diameter, ellipsoid, more rarely subglobose. *O. atropurpurea*, var. *natalia*, “*Flora Capensis*,” I, 448; *Sim*, “*Forest and Forest Flora*,” p. 163.

COAST REGION.—Uitenhage Div.: Uitenhage, May, *E. and Z.* 935; Motinsele Forest, December, *Fegen in Herb. Forest Dept.* 1816, 1818.

EASTERN REGION.—Natal: Berea, 200 ft., August, *Wood* 10060; *Wood in Natal Govt. Herb.* 10062, 10064; without precise locality, *Wood* 6418; near Durban, *Wood* 72; Benvie, Karkloof, 4000–500 ft. *Wylie in Natal Govt. Herb.* 10826; (*Wood* 10034); Umzim-yati, 200–500 ft., October, *Wood* 11458; without precise locality, *Cooper* 1252.

ORANGE FREE STATE: Elandshoek, September, *Rogers in Herb. Transvaal Museum* 2587.

SWAZILAND.—Komassan Valley, Havelock Concession, July, *Galpin* 983.

TRANSCAAL.—Without precise locality, *Miss Stent, in Herb. Transvaal Mus.* 8779. Pietersburg Div.: The Downs, *Rogers* 22133; Tzaneen, *Rogers* 12540. Barberton Div.: Unvoti Creek, near Barberton, 3000 ft., November, *Galpin* 660; *Rogers* 20267; Nelspruit, October, *Breyer in Herb. Transvaal Mus.* 17701; between White River and Nelspruit, *Burt-Davy* 1510; Carolina, September, *Roberts in Herb. Transvaal Mus.* 15874.

O. leptoclada (Oliv., “*Fl. Trop. Afr.*” I, 318).

A bush, 2–3 ft. high (*Galpin*). *Branches* somewhat pustulate with ash-grey bark and with the epidermis peeling off in membranous flakes. *Young leaves* 2·5–6·5 cm. long, 0·6–1 cm. broad, lanceolate, subacute, tapering at the base, with serrated margins, sometimes subentire. *Mature leaves* 5–11·5 cm. long, 1·4–3·4 cm. broad, mostly oblanceolate, sometimes lanceolate-elliptic, rounded at the apex, usually long-attenuate at the base, more rarely only narrowed, with the mid-rib prominent above and beneath, and the lateral veins distinct, with serrated margins, rarely subentire. *Inflorescence* mostly unbellate on abbreviated shoots, 3–4-flowered, sometimes flowers solitary. *Pedicels* 1–2·5 cm. long, articulated at the very base. *Sepals* 7 mm. long, 5 mm. broad, obovate or elliptic, rounded at the apex. *Petals* 1 cm. long, 5 mm. broad, obovate, rounded at the apex, narrowed into a claw at the base. *Filaments* 3·5 mm. long; anthers 1·5 mm. long, oblong-linear. *Ovary* of 5 carpels; style 4·5 mm. long; stigma capitate. *Fruit* about 6 mm. in diameter, subglobose.

TRANSCAAL.—Barberton Div.: Barberton, November, *Rogers* 18264, and *in Herb. Transvaal Mus.* 15976; mountain sides, Moodies, near Barberton, September–October, 3500 ft. *Galpin* 590; mountains near Barberton, *Bolus* 7693. Lydenburg Div.: Pilgrims Rest, c. 3500 ft., October, *Rogers*, 23068, and *in Herb. Transvaal Mus.* 20541.

RHODESIA.—North-West Rhodesia: Pemba, 3500 ft., October, *Rogers* 8579; Ndola, October, *Allen*, 304; Chilanga, 3500 ft. *Rogers* 8455. Untali Div.: Odanzi River Valley, *Teague* 399, 497.

BELGIAN CONGO: Elisabethville, 4800 ft., *Rogers* 10194.

O. atropurpurea [L.C. in “*Ann. Mus. Par.*,” XVII (1811), 412].

Small shrub, 3–5 ft. high. *Branches* densely covered with prominent lenticels giving them a lepidote appearance to the naked eye. *Leaves* subsessile, 0·8–5 cm. long, 0·4–1·9 cm. broad, usually elliptic, rarely elliptic-ovate, elliptic-oblong, or elliptic-lanceolate, more rarely lanceolate, obtuse, rounded or slightly narrowed at the base, with serrated margins and the teeth usually triangular erect and acute, more rarely incurved, with the mid-rib

prominent above, and lateral veins evident. *Flowers* solitary at the apex of short shoots, very rarely 2-nate. *Pedicels* 1-1.5 cm. long, articulated at the base or about 1 mm. from the base. *Sepals* 6-7 mm. long, 3-6 mm. broad, elliptic, elliptic-ovate or elliptic-oblong, rounded at the apex. *Petals* 0.9-1.3 cm. long, obovate, cuneate at the base. *Filaments* 2-2.75 mm. long; anthers 2-3.5 mm. long, oblong or linear in outline. *Ovary* of 5-6 carpels; style 5-7 mm. long; stigmas as many as the carpels. *Fruit* 7 mm. long, about 5 mm. in diameter, subglobose.—“*Flora Capensis*,” I, 448; *Sim*, “*Forests and Forest Flora*,” p. 163, pl. xix, fig. 2.

CAPE PROVINCE.—Caledon Div.: Genadendal, 2500 ft., April, *Schlechter* 7749; Bedford, June, *Miss Nicol* 49. Knysna Div.: Woods at Knysna, *Neudigate in Herb. Albany Museum*; *Watton in National Herb.* 1419. Port Elizabeth Div.: Van Stadens, September, *Mrs. Patterson* 737, *MacOwan in Cape Govt. Herb.* Albany Div.: Howisons Poort, near Grahamstown, *MacOwan in Herb. Austro-Afric.* 1766; Rockliffe, near Sidbury, November, *Miss Daly* 771; Grahamstown Commonage, 2000 ft., September, *Schönland* 570; rocky hills near Grahamstown, 2000-2500 ft., August, *Galpin* 48; near Grahamstown, September, *MacOwan* 111; and in Natal Govt. Herb. 850; January, *Breyer in Herb. Transvaal Mus.* 16866; Signal Hill, near Grahamstown, 2200 ft., September, *Bolus* 1920. Queenstown Div.: Among rocks on hillside, Junction Farm, Queenstown, 2500 ft., *Miss Wiggill in Herb. Galpin* 8257. East London Div.: East London, December, *Breyer in Herb. Transvaal Mus.* 16544. Komgha Div.: Wood near Prospect Farm, November, *Flanagan* 322. Stockenström Div.: In bush at Willisdale, September, *Scully* 77; Kat River, *Bartels*; Katherg Forest, January, *Stapels in Herb. Forest Dept.* 2937, 2938. Stutterheim Div.: Stutterheim, December, *Rogers* 12726; Transkei, Umtata, *Convent School Herb.* 194; Kentani, 1200 ft., October, *Miss Pegler* 194; rocky spots near Bazya, 2000 ft., October, *Baur* 258; Maseko Forest, Negeleni, December, *Fegen in Herb. Forest Dept.* 1819; Pondoland, Mgezama River, near McAllister's Drift, Port St. Johns, *Miller in Herb. Forest Dept.* 3024.

NATAL.—Near Umkomaas River, October, *Tyson* 2659; Natal Bay, *Gueinzins*; without locality, *Wood* 190.

ZULULAND.—Qudeni Forest, 6000 ft., January, *Davis* 108.

TRANSVAAL.—Barberton Div.: Near Barberton, September, *Bolus*. A small stunted shrub. Plentiful on edge of forest, but does not appear to seed freely (*Fegen*).

Var. *angustifolia*, Phillips. Differs from the type in having smaller, narrower lanceolate leaves.

TRANSVAAL.—Barberton Div.: Margin of wood, upper Moodies, 4500 ft., June, *Galpin* 963, and in Natal Government Herb. 8761. Ermelo Div.: Nooitgedacht, December, *Mrs. Pott* 5096, and in Herb. Transvaal Mus. 14933. Lydenburg Div.: Between Pilgrims Rest and Sabi, *Rogers* 23410.

SWAZILAND.—Havelock Concession, *Saltmarsh in Herb. Galpin*. A shrub 3 ft. high (*Galpin*).

O. pretoriensis, Phillips (“*Flowering Plant S. Afr.*,” vol. ii).

Branchlets with light to dark-brown bark, peeling in membranous strips, not distinctly pustulate. *Leaves* 1.5-3.8 cm. long, 0.5-1.7 cm. broad, lanceolate, obovate, elliptic, or oblanceolate (mostly oblanceolate), obtuse at the apex, cuneate, more rarely rounded at the base, with serrated margins and with the mid-rib distinct and lateral veins evident. *Flowers* solitary, very rarely 2-nate, arising at the apex of abbreviated shoots. *Pedicels* 1-1.5 cm. long, articulated at the base or 1-2 mm. above the base. *Sepals* 7-8 mm. long, 4-5 mm. broad, ovate or elliptic, rounded at the apex, sometimes 2-3-lobed at the apex, enlarging in fruit. *Petals* 1.5 cm. long, 8.5 mm. broad, obovate, rounded at the apex, narrowed at

the base into a claw. *Filaments* 4·5 mm. long; anthers 2 mm. long, oblong in outline. *Ovary* of 4–5 carpels; style 5 mm. long; stigmas as many as the carpels. *Fruit* 7–8 mm. long, 5–6 mm. broad, more or less ellipsoid.

TRANSVAAL.—Pretoria Div.: Below River Redoubt, Daspoort Rand, October (fruits), *Pole Evans* 157; Pretoria, December (fruits), *Nelson in Herb. Transvaal Mus.* 11709; February (fruits), *Miss Leendertz* 65, and in *Herb. Transvaal Mus.* 4199; November (flowers) *Miss Leendertz* 301; Meintjes Kop, March (fruits), *Pole Evans* 353; Kopje near Gezina, *Phillips in National Herb.* 1422; *Fouche in National Herb.* 1491; Moorddrift, October (fruits), *Leendertz* 2131, and in *Herb. Transvaal Mus.* 7333; Rikatla, October (fruits), *Junod* 147, and in *Herb. Transvaal Mus.* 20146; near Messina, c. 1800 ft., November (fruits), *O'Connor in Herb. Forest Dept.* 1936; Messina, October (fruits), *Rogers* 19398, 20029, 19244, and in *Herb. Transvaal Mus.* 17762.

THE GENUS OLINIA.

By JOAN HOFMEYR and E. P. PHILLIPS, M.A., D.Sc., F.L.S.,
Division of Botany, Pretoria.

THE genus *Olinia* was founded by Thunberg in 1799 ("Roem. Arch.," II, i, 4, 1799), and belongs to the natural order *Lythraeae*. The genus contains important forest trees, and was first dealt with in this connection by Sim in his "Forests and Forest Flora," p. 227. Sim only recognized one species, viz. *O. cymosa*, but divided this into three varieties, as follows:—

Var. *latifolia*, mostly in the Western Province.

Var. *intermedia*, found in the Midland Conservancy and sparingly elsewhere.

Var. *acuminata*, the common form in the Eastern and Transkeian Conservancies.

The examination of all the material in the South African herbaria has led us to regard var. *latifolia* and var. *intermedia* as belonging to the species *O. cymosa*, and var. *acuminata* as a distinct species which Klotzsch first named *O. acuminata*. The leaf characters which Sim gives for the first two varieties are not constant in the areas in which they are stated to occur, and we have failed to find even constant varietal characters. The var. *acuminata*, which has again been raised to specific rank can easily be distinguished from *O. cymosa* by the shape of the petals and, to a lesser degree, by the shape of the leaves. In the former the petals are "linear-spathulate" (Pl. II, figs. 2, 3), while in the latter (*O. cymosa*) the petals are "obovate-spathulate" (Pl. I, figs. 2, 3). These characters, in conjunction with its distinct distribution, all point to the tree being quite distinct from *O. cymosa*. It is the plant which Sim has figured on Plate LXXII in his work cited above.

In the "Index Kewensis" two other specific names are mentioned, viz. *O. ternata*, Gilg and *O. micrantha*, Decr. As no other descriptions of these were available in South Africa, the Director of Kew was asked for assistance. In a letter from Kew received in January, 1921, it is stated that "the name *O. ternata*, Gilg is apparently not yet published but Gilg determined a specimen himself collected by Mrs. H. Hutton at Howison's Poort near Grahamstown as this species." There can be no doubt that this is *O. cymosa*. From the description of *O. ternata*, Decne., kindly sent to us by the Director of Kew, we also have no hesitation in confirming this to be *O. cymosa*. Decaisne founded his description on Burchell 3592, which was not represented in any of the South African herbaria. Another species from Zululand and East Pondoland was named by Dr. Stapf as *O. radiata* but the name has not been published. As this name is now known to foresters, we do not propose to change it.

The Curator of the Botanical Department of the British Museum was good enough to furnish a scrap of *Welwitsch* 991 collected at Huilla and stated to be *O. cymosa*, but an examination of this makes it very doubtful whether *O. cymosa* does occur in tropical Africa. This has now been confirmed by Kew, as the Director writes: "*Welwitsch* 991 is not represented at Kew, but reference to the British Museum material leads us to agree that it is distinct from *O. cymosa*. It very closely resembles *O. usambarensis* (Holst 9115)."

Through the courtesy of the Forest Department, we have been able to give some notes on two of the species, *O. cymosa* and *O. radiata*.

Mr. J. D. Keet, District Forest Officer, Knysna, writes regarding *O. cymosa*: "Found in all forests in Knysna and Zitzikamma Districts, mostly in forests on the coastal plateau at elevations of 700 to 900 ft. approximately, less frequent in the forests on the foothills and mountain ravines and scarce in the coastal scrub. Its total range is from 300–400 to 2000 ft. approximately. It is comparatively intolerant of shade and superfluous soil moisture, and generally avoids cool slopes, ravines, low-lying ground, and streams. It is often a forerunner of the forest-like 'Keur' (*Virgilia capensis*), 'Beech' (*Myrsine melanophlebos*), etc., and like the Sneezewood (*Ptaeroxylon utile*), it prefers the crests of ridges, rocks, and stony ground. Other conditions being suitable, it grows equally well on soils originating from Table Mountain Sandstone, Bokkeveld, Conglomerate, and old sand-dunes (?). The result is that it is found mostly on ridges on the north and west slopes near the crest of a ridge, the edge of the forest, and frequently in small groups or as isolated trees in the 'fijnbos.' It is a medium-sized tree in height and usually in bole. The thickest I have measured is 15 ft. in girth, the largest bole 32 ft., and the greatest height 70 ft. (estimated). The thicker trees are often somewhat gnarled, fluted, and buttressed, and with unsound heartwood. Where exposed to the full light it forms full-foliaged and much-branched symmetrical crowns on low stems. This is generally the case in older trees which have been a forerunner of the forest. The bark is thin, usually dark, and decorticating in scales on the lower part of stems of older trees; sometimes reddish or brownish pigment shows in the cracks and under the scales. Bark on the smaller branches smooth and grey. At a glance in the forest the bark may sometimes be confused with that of 'Quar' (*Plectronia obovata*). The wood is used in wagon work (such as felloes, spokes, long-wagon poles), fencing and telegraph poles, railway sleepers, furniture, etc. The plant coppices freely, and coppice shoots are vigorous on small and medium sized trees, but usually too dense, in which case all may die back at a height of 6–8 ft., when fewer will reach pole size in twenty to twenty-five years. On stools of trees felled in 1917 at Harkerville the coppice is now 6–8 ft. high. Seedlings occur sparingly in open space in worked-out forests and in the 'fijnbos' edge, but are absent in dense forest. Several attempts to germinate the seeds in nurseries have either failed completely or given very poor results. Seedlings and saplings in the forest are fairly fast growing. In the forest a rough and ready test by which the tree may be recognized is the strong prussic acid or almond smell of the leaves, twigs, freshly-cut bark, and sapwood. The name 'mountain hard pear' is applied to this species."

O. radiata. This is said to be the largest tree growing in the Ingeli Forest, and is over 12 ft. girth at breast high. The District Forest Officer, Transkeian Conservancy, reports "that the bark has a strong almond-like smell, and when placed in water turns it a blue-black colour. The wood is used for felloes." Forester Fegen, in submitting specimens of the tree, states: "This tree grows to a large size and is very plentiful in the Izinja, Ngododo, Qakama, Maseko, and neighbouring forests, and is not found elsewhere in the district. The young wood is quadrangular, usually red in colour, as are the leaves. Flowers white with a tinge of purple in small dense clusters. Fruit plum-shaped, reddish purple in colour when ripe, is eaten by birds, and I hear by natives. The tree flowers about November and the fruit is ripe about July. The wood is reported to be hard and durable, but nothing is really known about the tree here. Young trees and saplings are used by the natives." Forester Leigh gives the native name of the tree as "umpanzi," and states that sawyers confuse this with "umpanzite" (*Phyllanthus amapodensis*) as the wood is the same. It is fairly abundant in the Ntsubane Forest. Forester Leigh also states that he only knows of it growing between Unsikaba and Umzimhlava Rivers, and that it is a large semi-deciduous tree 60–70 ft. high with a girth of 10–12 ft. Old trees grow out in ribs near the ground. Used for wagon work (naves, felloes, etc.), and believed to make a good charcoal. It is plentiful in parts of the Ntsubane Forest, but on the whole scarce.

O. acuminata. We have not been able to obtain any special information about this species, but in the Woodbush Forest, Pietersburg, Transvaal, it is known as "roodebosje."

The distribution of the three species found in South Africa is fairly well defined.



Stella Gower.

Olinia cymosa, Thunb.

Olinia cymosa extends from the Cape Peninsula along the coastal belt as far as the Kei mouth, beyond which we have no record of its occurrence. MacOwan collected the species at Somerset East, which is the only record of its inland occurrence.

Olinia radiata. Is found in the Pondoland forests and extends as far as Ingeli in Natal.

Olinia acuminata. This is an inland species. The most southern record is at Queenstown, and from there it extends to the slopes of the Drakensberg in East Griqualand to the mountainous districts of the eastern Transvaal (Barberton), and occurs again in the Woodbush Forest in the Pietersburg District. An outlier of the species is recorded from the Magaliesberg in the Transvaal.

The flowers are largely infected by a Hemiptera larva which appears to arrest their development, which is reflected in the length of the calyx-tube. In some specimens the calyx-tube is much longer than the petals, in other it equals the petals, while specimens are found in which the calyx-tube is shorter than the petals. All gradations are found even on the same specimen.

Two specimens of *O. cymosa* differ slightly from the other specimens of the species examined in that the petals are rounded at the apex and not bluntly pointed (East London, Rattray 840, Somerset East, Bolus 1772), but in specimens collected by Flanagan near the Kei mouth both forms of petals occur.

KEY TO SPECIES

Petals obovate-spathulate.....	<i>cymosa</i> .
Petals linear-spathulate.	
Inflorescence lax, about half as long as the subtending leaf.....	<i>acuminata</i> .
Inflorescence compact very much shorter than the subtending leaf.....	<i>radiata</i> .

Olinia cymosa, Thunb., "Flora Capensis," 194. (Plate I.)

Branches glabrous. *Leaves* 3·7 cm. long, 1·3·5 cm. broad, lanceolate, obovate, elliptic, acuminate, obtuse, sometimes rounded at the apex, narrowed at the base into a distinct petiole, margins entire, sometimes wavy, glabrous, green and shiny above, paler beneath. *Inflorescence* axillary, a dense trichotomous cyme, very much shorter than the subtending leaf. *Bracts* opposite, same length as the flowers, often deciduous. *Penduncles* minutely pubescent. *Calyx-tube* 1·6 mm. long, glabrous or minutely hairy with five minute teeth. *Petals* five, inserted at the throat of the tube, about one-third the length of the calyx, rarely equalling it, spathulate, acute or obtuse at the apex, with a cluster of hairs at the base. *Scales* five incurved, alternating with the sepals, minute, obovate, densely hairy. *Stamens* five, adnate to calyx below the scales; filaments very short. *Ovary* inferior, five-celled, with three pendulous ovules from a central placentum in each cell; style subulate; stigma obtuse. *Fruit* globose, 1 cm. in diameter.—*Flora Capensis*, II, 520, Excl. var. *acuminata*; Sim, "Forest and Forest Flora," 227, Excl. var. *acuminata*, pl. LXXII, fig. 14.

CAPE.—Slang Kuil, Table Mountain, October, MacOwan in *Herbarium Austro-africanum* 927; Kirstenbosch, Table Mountain, November, December, Harcey, July, Zeyher 179 and 244; August, L. Kensis in *Herb. Bolus Austro-africana* 10748; Table Mountain, August, M. Page in *Herb. Afric. Bolusianum* 1633; Devil's Peak, September, Zeyher; Paarl, August, October, Drège; Hottentot Hollands Mts., July, Zeyher; Swellendam, Voormansbosch, October, Zeyher; Grootvadersbosch, October, Pappe 130, Zeyher, 2434; Knysna, Commonage forests, August, J. D. Keet in *Herb. Forestry Dept.* 2336; May, Scott-Elliot in *Forest Dept. Herb.* 1349; Collector (?) *Forest Dept. Herb.* 1350. Port Elizabeth: September, I. L. Drège in *Herb. Albany Museum* 541; August, F. Paterson in *Herb. Africanum Bolusianum* 2320, 2319, 1155; August, Dom. Honsley in *Herb. Albany Museum*. Humansdorp: August, F. A. Rogers in *Herb. Albany Museum* 3016. Van Stadens: September, T. V. Paterson in *Herb. Albany Museum* 230. Bethelsdorp: September, T. V.



Stella Gower.

Olinia acuminata, Klotz.

Paterson in Herb. Albany Museum 2151. Somerset East : November, *H. Bolus* 1772. Grahams' own : "Amos's Vil," *MacOwan* 349 ; August, September, *E. E. Galpin in Herb. Albany Museum* 88. Howisons Poort : September, *Schönland* 818 ; *Zeyher* 2465. East London : October, *Geo. Rattray in Herb. Albany Museum* 840 ; Kei Mouth, July, *Flanagan* 2341.

Olinia acuminata, Klotz in "Otto and Dietr. Allg. Gartenz," IV (1836), 27 (Pl. II).

Branches glabrous. *Leaves* 2-3 cm. long, 0.5-1.5 cm. broad, lanceolate, elliptic, acuminate, narrowed at the base into a more or less distinct petiole, with entire margins, glabrous, green and shiny above, paler beneath. *Inflorescence* an axillary trichotomous cyme, a little shorter than the subtending leaf. *Bracts* deciduous. *Calyx-tube* 2-9 mm. long, glabrous or minutely pubescent, with five minute teeth. *Petals* five, inserted at the throat of the calyx-tube, about one-third the length of the calyx, linear, rounded at the apex, with a small tuft of hairs at the base. *Scales* five, incurved, alternating with the petals, minute, obovate, densely hairy. *Stamens* five, adnate to calyx below the scales ; filaments very short. *Ovary* inferior, five-celled, with three pendulous ovules from a central placentum in each cell ; style subulate ; stigma obtuse. *Fruit* globose, 3 mm. in diameter.—"*Flora Capensis*," II, 520, *O. cymosa*, var. *acuminata* ; *Sim*, "Forests and Forest Flora," 227.

CAPE.—Queenstown, Rockwood, Bongolo, February, *E. E. Galpin* 2500. Transkei, Kaffraria, January, *T. R. Sim* 2026 ; June, 2099.

NATAL.—Maclear, Pot. River Berg, March, *E. E. Galpin* 6628, 6629 ; Drakensberg, May, *J. Medley Wood* 4957 ; Giant's Castle, June, *R. E. Simons in Herb. Trans. Museum* 15965.

TRANSVAAL.—Barberton, Upper Moodies, January, *E. E. Galpin* 1275 ; Rustenburg, January, *Olive Nation in Herb. Bolusianum* 1906 ; Magalisherg, *Zeyher* 494 ; Woodbush, northern Transvaal, January, *O. J. O'Connor in Forest Dept. Herbarium* 1458 ; Dulstroom, January, *F. O. Noome in Herb. of Trans. Museum* 20812 ; Elandspruitberg, December, *R. Schlechter* 3864.

Olinia radiata, sp. nov. (Plate III). *Rami* glabri. *Folia* 5-8 cm. longa, 1.8-3.8 cm. lata, lanceolata, obovata, lanceolato-elliptica vel elliptica, subacuminata, apice obtusa vel aliquando subrotundata, basi angustata. *Inflorescentia* axillaria, compacta. *Pedunculus* minute pubescens. *Calyci tubus* 1.4 mm. longus, glaber vel minute pubescens. *Petala* 1-2 mm. longa, linearia, spathulata, apice acuta. *Squamae* minutae, obovatae, pilosae.

Branches glabrous. *Leaves* 5-9 cm. long, 1.8-3.8 cm. broad, lanceolate, obovate, lanceolate-elliptic, or elliptic, mostly subacuminate, obtuse, more rarely rounded at the apex, narrowed at the base into a more or less distinct petiole, margins entire and usually wavy, glabrous. *Inflorescence* axillary, a dense trichotomous cyme, very much shorter than the subtending leaf. *Bracts* opposite, smaller than the flowers, deciduous. *Peduncles* minutely pubescent. *Calyx-tube* 1-4 mm. long, glabrous, or minute hairy ; teeth very minute. *Petals* five, inserted at the throat of the calyx-tube, 1-2 mm. long, linear-spathulate, acute, with a small tuft of hairs at the base. *Scales* incurved, alternating with the petals, minute, obovate, densely hairy. *Stamens* five, adnate to the calyx below the scales ; filaments very short. *Ovary* inferior, five-celled, with three pendulous ovules from a central placentum in each cell ; style subulate, stigma obtuse. *Mature fruit* globose, 1.8 cm. in diameter.

TRANSKEI : Ngadodo Forest, Ngqeleni, November, *Fegen in Herb. Forest Dept.* 2286, and in *Government Herb.* 18361 ; September, *Fegen in Herb. Forest Dept.* 2490. PONDOLAND : Ngadu Forest Station, May, *Van der Vyck in Herb. Forest Dept.* 2472. Lusikisiki Dist. : Ntsubani Forest, *A. Leigh in Herb. Forest Dept.* 2051, 1553 ; Mpanzi, February, *A. Leigh in Herb. Forest Dept.* 1839. NATAL : Ingeli Forest Station, November, *C. W. Chilceers in Herb. Forest Dept.* 1942 ; Impetyne, February, *J. S. Henkel in Herb. Forest Dept.* 2382.



Stella Gower.

Otinia radiata, Phill. and Hofmeyr.

EXPLANATION OF PLATES.

Plate I.—*Olinia cymosa*, Thunb.

Fig. 1. Portion of branch.

Fig. 2. Flower.

Fig. 3. Longitudinal section of flowers showing stamens, scales, ovary, and style.

Fig. 4. Scale.

Fig. 5. Cross-section of ovary.

Fig. 6. Stamen.

Figs. A, B, C. Types of leaves.

Plate II.—*Olinia acuminata*, Klotz.

Fig. 1. Portion of branch.

Fig. 2. Flower.

Fig. 3. Longitudinal section of flower.

Fig. 4. Scale.

Fig. 5. Stamen.

Fig. 6. Cross-section of ovary.

Plate III.—*Olinia radiata*, Phill. and Hofmeyr.

Fig. 1. Portion of branch.

Fig. 2. Flower.

Fig. 3. Longitudinal section of flower.

Fig. 4. Scale.

Fig. 5. Stamen.

Fig. 6. Stamen and scale.

Fig. 7. Cross-section of ovary.



THE GENUS CYCLOPIA, VENT.

By JOAN HOFMEYR and E. P. PHILLIPS, M.A., D.Sc., F.L.S.,
Division of Botany, Pretoria.

THE genus *Cyclopia* was founded by Ventenat in 1808. Two years later a plant, now known as *Cyclopia genistoides*, was figured in the *Botanical Magazine*, t. 1259, and described as the type of a new genus, viz. *Ibbetsonia*. In 1825 the genus *Cyclopia* was dealt with in De Candolle's "Prodromus" (Vol. II, p. 101), and three species described, and it was not until 1861 that Harvey revised the genus in the "Flora Capensis" (Vol. I, p. 6). Since Harvey's account nothing has been written on the genus, but in the interval much material has accumulated in local herbaria which extends our knowledge of the species.

Several species of *Cyclopia* have become of some commercial value locally as the source of "bush tea," and it was with the object of clearly defining the known species as a foundation on which any future experimental work might be based that this revision of the genus was undertaken.

Harvey in the "Flora Capensis" recognized nine species, but our examination of all the available local material has led us to subdivide the genus into twelve very distinct species. The species are all shrubs with ternate, very often revolute leaves, and yellow flowers. They are mostly confined to the Western Province and the south-eastern regions, but one species extends as far as Port Elizabeth. According to observations made by Forest Officers, most of the species grew well in all situations and soils, except on stiff clayey soils. The cultivation of the species used as "bush teas" should be experimented with, and should not prove difficult, though the young seedlings may require some shelter until firmly established.

The method at present used for preparing "bush teas" for local consumption is to heat the leaves in an oven for about an hour to produce "sweating," after which they are dried in the sun. When thoroughly dry the leaves and smaller twigs are removed from the stem and larger branches, packed, and sold under various names. If the demand for "bush tea" is sufficiently great, cultivation trials will be justified, as it may then be possible to put a superior article on the local market.

Our thanks are due to the Curators of the Albany Museum Herbarium, the Cape Government Herbarium, the Transvaal Museum Herbarium, and the Forest Department for allowing us to examine the material in their respective herbaria.

C. genistoides. The District Forest Officer, Elgin, Caledon District, reports: "This species grows on the slopes of mountains and hills, and is found in fairly large quantities on the Greenland Range, near Elgin. It grows to approximately 3 ft. in height. The leaves and flowers are sun-dried." Known as "honey-tea."

C. tenuifolia. The District Forest Officers, Elgin, Caledon District, reports: "This species is not unlike the 'honey tea' (*C. genistoides*); the flowers are similar in colour and appearance, but the leaves are finer and longer. It is sun-dried before being used, and has a stronger flavour." Known as "vlei tea."

C. brachypoda. The District Forest Officer, Elgin, Caledon District, reports: "The leaves are broader and longer than the other two species (*C. genistoides* and *C. tenuifolia*). It also flowers later. It is found in large quantities throughout the Caledon and Swellendam Districts, and shares popularity with 'honey tea' (*C. genistoides*) as a beverage amongst the poorer classes. It is prepared in a similar manner to 'honey and vlei tea,' known as 'Heidelberg tea.'"

C. subternata. The District Forest Officer at Grootvadersbosch, Swellendam District, reports: "This variety is known as the 'common bush tea,' and thrives abundantly along the slopes of the Langeberg Mountain Range in the Divisions of Swellendam and Riversdale, and large quantities are available each year. The flowering season is generally in the month of August. Large quantities are collected on private ground, the bulk of which is used by poor whites and coloured people, very little being sold owing to the small amount (1½d. per lb.) offered by local buyers after preparation." A form of this species is known as "Hottentot's tea," and according to the District Forest Officer at Grootvadersbosch, "only small quantities of this tea can be found growing, especially in the heart of the Langeberg Mountain Range. Flowering season in August. Very little tea is collected annually."

C. Ashtonii. The District Forest Officer at Grootvadersbosch, Swellendam District, states: "This tea, like *C. subternata*, thrives well along the mountain slopes of the Langeberg, but prefers the damper places. Consequently, the only large patches are found on the mountain side. The flowering season is about November, which is practically the only time that this tea is collected, and then only in small quantities." Locally known as "vlei tea."

C. Burtonii. The Conservator of Forests, Knysna, reports: "It is known in the Zwartberg Mountains as 'heuning or sugar tea.' The shrub grows to a height of 1½–18 in., and is only found in a few places on the Zwartberg Mountains. Apparently it grows at a higher altitude than the ordinary 'bush tea,' and appears able to stand the frost better."

CYCLOPIA, VENT.

Erect branched shrubs. *Leaves* sessile, palmately trifoliate; leaflets narrow-linear, lanceolate or very rarely ovate, glabrous or pubescent, frequently with revolute margins; stipules none. *Peduncles* axillary, one-flowered, with two, rarely three, bracts at the base. *Flowers* bright yellow. *Calyx* subequally 5-cleft, with an intruse base. *Vexillum* roundish, plaited at the base and with a short claw; alae oblong, with a transverse fold; carina incurved, obtusely rostrate, with a small triangular pocket. *Stamens* separate or slightly connate at the base; filaments dilated. *Ovary* glabrous, several ovuled. *Legume* oblong, compressed, with coriaceous valves.

KEY TO SPECIES.

Plants glabrous or nearly so.

Leaflets flat or with slightly recurved margins.

- | | |
|---|-------------------------|
| Bracts three..... | 1. <i>Burtonii</i> . |
| Bracts two. | |
| Leaflets ovate, cordate or truncate at the base..... | 2. <i>latifolia</i> . |
| Leaflets elliptic, oblong or sub lanceolate-linear, obtuse or acute. | |
| Pedicels exceeding the small obtuse or acute bracts; lobes of the calyx equalling the tube or rarely shorter..... | 3. <i>subternata</i> . |
| Pedicels the same length as the bracts; the lobes of the calyx exceeding the tube..... | 4. <i>brachypoda</i> . |
| Leaflets linear or filiform with strongly revolute margins. | |
| Bracts very broad, obtuse..... | 5. <i>tenuifolia</i> . |
| Bracts boatshaped, acute..... | 6. <i>genistoides</i> . |

Plants villous or pubescent at least on the younger parts.

- | | |
|---|---------------------------|
| Bracts ribbed, recurved at the apex..... | 7. <i>pubescens</i> . |
| Bracts smooth, straight. | |
| Flowers pedicellate. | |
| Calyx glabrous. | |
| Leaflets linear, strongly revolute..... | 8. <i>galioides</i> . |
| Leaflets narrowly ovate, cordate at the base, slightly revolute.. | 9. <i>Ashtonii</i> . |
| Calyx softly hairy..... | 10. <i>Bolusii</i> . |
| Flowers subsessile. | |
| Lobes of the calyx falcate..... | 11. <i>sessiliflora</i> . |
| Lobes of the calyx oblong, not falcate..... | 12. <i>montana</i> . |

1. *C. Burtonii*, sp. nov.

Rami glabri. *Foliola* subsessilia, 0·6–1·2 cm. longa, 2 mm. lata, lanceolato-elliptica, apice acuta, plana, marginibus leviter reflexis. *Bractea* 3, 1–2 mm. longae, lanceolatae, apice acutae. *Calyx* 3 mm. longa; lobi ovato-acuminati. *Vexillum* 8·5 mm. longum, 6 mm. latum, apice acutum; ala oblonga; carina rostrata.

Oudtshoorn Division: Zwartberg Mountains, November, *Burtonia* in Herb. Forest Dept. 2914.

A much-branched glabrous shrub, 30–37 cm. high. *Leaflets* subsessile, 0·6–1·2 cm. long, 2 mm. broad, lanceolate-elliptic, acute at both ends, flat, with slightly reflexed margins. *Bracts* three, 1–2 mm. long, lanceolate, acute, equalling the pedicel; one adnate to the pedicel and smaller than the other two. *Calyx* 3 mm. long; lobes ovate, acuminate, longer than the tube. *Vexillum* 8·5 mm. long, 6 mm. broad, acute; alae narrow-oblong; carina very obtusely rostrate.

2. *C. latifolia* (D.C., Prodr., 2, p. 101).

Plant glabrous. *Stem* strongly ribbed. *Leaflets* sessile, 0·5–2·3 cm. long, 0·3–1·5 cm. broad, ovate, mucronate, cordate or truncate at the base, with entire slightly recurved margins; the middle leaflet the longest. *Bracts* two, 0·5 cm. long, sharp pointed, as long as or almost as long as the pedicel, persistent. *Pedicel* 0·8–1 cm. long. *Calyx* 0·5 cm. long; segments acute, slightly falcate, as long as the tube; the lowest segment longer and minutely toothed at the base.—“*Flora Capensis*,” II, 6.

Cape Division: Table Mountain, October, *Bodkin* in Herb. *Bolus* 8038.

3. *C. subternata* (Vog. Linn. 10, p. 595).

A branched glabrous shrub. *Stems* ribbed. *Leaflets* 1·25–3·75 cm. long, 2–5 mm. broad, elliptic, oblong or lanceolate, acute or obtuse, flat, with reflexed margins. *Bracts* two, 2 mm. long, rarely 4 mm. long, broadly ovate, mucronate, shorter than the pedicels. *Pedicel* 4–11 mm. long. *Calyx* 4–5 mm. long; lobes deltoid, obtuse or subacute, as long as or shorter than the tube.—*C. Vogelli*, Harv., var. *subternata*, Harv., “*Flora Capensis*,” II, 6.

George Division: Without precise locality, September, *District Forest Officer* in Herb. Forest Dept. 3112. Port Elizabeth Division: Port Elizabeth, *Patterson* 863; Witte Els Bosch, *Zitzikama*, September, *Galpin* 3899. Riversdale Division: Muis Kraal, near Garcias Pass, October, *Galpin* 3900; *Phillips* in Herb. *Musei Austro-Afric.* 1991; Toll House. 1700 ft., October, *Bolus* 11249. Swellendam Division: Grootvadersbosch, *District Forest Officer* in Herb. Forest Dept. 2844, 2845. Cape Division: Capetown, October, *Tyson*. Paarl Division: French Hoek, September, *Bolus*. Tulbagh Division: Witsenberg Vlakke, November, *Zeyher*. Knysna Division: Knysna, September, *Keet* in Herb. Forest Dept. 2539; Plettenberg Bay, *Zeyher*. Ceres Division: Koude Bokkeveld, September, *Schlechter* 8922, *Burchell* 5549. Oudtshoorn Division: Zwartbergen, 3000 ft., October, *Schlechter* 5583. Humansdorp Division: Mountains near Humansdorp, Herb. *Bolus* 1056; near French Hoek, c. 2400 ft., November, *Bolus* 5151 (in this specimen the base of the calyx is intruse and thus the lobes appear to be longer than the tube).

Specimens collected by Gatherstone at Cango and by Kennedy at Humansdorp compare well with *C. subternata*, except that they have a split fleshy calyx.

4. *C. brachypoda*, comb. nov.

A glabrous shrub. *Stems* ribbed. *Leaflets* 1.25-2.5 cm. long, 2-3 mm. broad, lanceolate or narrow-linear, acute or obtuse, flat, with reflexed slightly wavy margins. *Bracts* two, 4-6 mm. long, ovate-lanceolate, acuminate, very acute, as long as the pedicel. *Calyx* 5-7 mm. long; lobes longer than the tube, falcate, acute.—(*C. Vogelii*, Harv., var. *brachypoda*, Harv., "*Flora Capensis*," II, 6.

Paarl Division: French Hoek, October, *Phillips* 1097. Tulbagh Division: Witsenberg, December, *Ecklon* and *Zeyher* 1147; September, *Zeyher* 354.

Var. *intermedia*. *Bracts* and *pedicels* 1 mm. long. *Leaves* 6-8 mm. long, 2 mm. broad, very obtuse.—*C. Vogelii*, Harv., var. *intermedia*, "*Flora Capensis*," II, 7; *C. intermedia*, E. Mey Com., p. 3, excl. lit c.

Caledon Division: Without precise locality, *District Officer in Herb. Forest Dept.* 2841. Riversdale Division: Langeberg, Riversdale, August, *Muir* 843.

A specimen collected at Elgin in the Caledon Division (*Herb. Forest Dept.* 2891) compares well with *C. brachypoda*, but the material is insufficient to be quite certain of the species.

5. *C. tenuifolia* (Lehm. Linn. 5, p. 373).

Plant glabrous. *Stem* ribbed. *Leaflets* 1.25-5 cm. long, linear-filiform, with revolute margins. *Bracts* two, 3 mm. long, 5 mm. broad, folded longitudinally and overlapping each other, obtuse or truncate, strongly keeled, glabrous without, sparsely hairy within. *Pedicel* 1.25-1.75 cm. long, elongating in fruit. *Calyx* 5 mm. long; lobes as long as the tube, acute, with slightly woolly margins; the lowest lobe longer and broader than the others, minutely toothed at the base. *Legume* oblong, with a recurved mucro at the apex; valves leathery.—"*Flora Capensis*," II, 7.

Caledon Division: Elgin, October, September, *District Forest Officer in Herb. Forest Dept.* 2840, 3114. Swellendam Division: River Zonder Einde, *Zeyher* 2257. Riversdale Division: Corento River Farm, Riversdale, November, *Muir* 50, and in *Herb. Galpin* 5072.

6. *C. genistoides* (Br. Hort. Kew., ed. 2, vol. 3, p. 5).

Stem ribbed, glabrous or nearly so. *Leaflets* 1.25-3.75 cm. long, narrow-linear, with strongly revolute margins. *Flowers* in a dense mass near the ends of the branches. *Bracts* 5 mm. long, acute and mucronate at the apex, boat-shaped, folded longitudinally, glabrous without, with a few scattered hairs on the inner surface. *Pedicel* 6-8 mm. long. *Calyx* 8-11 mm. long; lobes longer than the tube, broadly falcate, with a mucronate apex, margins woolly; lowest segments longer than the others and toothed at the base.—"*Flora Capensis*," II, 7.

Caledon Division: Caledon, August, *Rogers* 11017; October, *Bolus*; *District Forest Officer in Herb. Forest Dept.* 2842. Cape Division: Tokai Plantation, September, *Herb. Forest Dept.* 1052; Cecelia Plantation, Ross in *Herb. Forest Dept.* 3113; near Capetown, October, *Bolus* 7272; Devil's Peak, November, *MacOwan* 1113, and in *Herb. Musei Austro-Afric.* 6546; Constantia, September, *Schlechter* 1461. Stellenbosch Division: Stellenbosch, October, *Mrs. De Jongh in Herb. Galpin* 4390. Swellendam Division: Swellendam, *Ecklon* and *Zeyher*. Clanwilliam Division: Clanwilliam, *Mader in Herb. Austro-Afric.*

7. *C. pubescens* (E. and Z. Enum. No. 1146).

Stem pubescent. *Leaflets* 1.25-3.75 cm. long, about 1 mm. broad, narrow-linear, with revolute margins, pubescent, at length becoming almost glabrous. *Flowers* densely massed. *Bracts* 4-6 mm. long, almost as broad, ribbed, acuminate, recurved at the apex, pubescent, shorter than the pedicel. *Pedicel* 1.25-1.75 cm. long, villous. *Calyx* 8 mm. long, pubescent; lobes taper-pointed from a broad base, twice as long as the tube.—"*Flora Capensis*," II, 8.

Port Elizabeth Division : Port Elizabeth, September, *Drège in Herb. Albany Museum* 167.

8. *C. galioides* (D.C., Prodr. 2, p. 101).

Young branches ribbed, densely villous, at length becoming almost glabrous. *Leaflets* arising from a thick flattened node, 1.25–2.5 cm. long, up to 4 mm. broad, linear, with strongly revolute margins, with a recurved mucro at the apex, truncate at the base, pubescent when young. *Bracts* 4–6 mm. long, ovate-lanceolate, acuminate, longer than the pedicel. *Pedicel* 2–4 mm. long. *Calyx* 6–13 mm. long, glabrous; lobes broadly falcate, acuminate, longer than the tube.—“*Flora Capensis*,” II, 8.

Cape Division : Fish Hoek, May, *Bolus in Herb. Norm. Austro-Afric.* 67; Muizenberg, Zeyher; Simonstown, February, *Compton*.

9. *C. Ashtonii*, sp. nov.

Rami sericei. *Foliola* 1.75 cm. longa, 4 mm. lata, basi cordata, marginibus revolutis. *Bractae* 4 mm. longae, ovatae, ciliatae, intus villosae. *Calyx* 8 mm. longus; lobi falcati.

Swellendam Division : Langeberg Mountains, *Ashton in Herb. Forest Dept.* 2843.

Young branches silky. *Leaflets* from a thick flattened node, up to 1.75 cm. long, 4 mm. broad, thick, cordate at the base, with revolute margins, paler beneath. *Bracts* 4 mm. long and slightly broader, ovate, ciliate, villous on the inner surface. *Calyx* 8 mm. long; lobes broadly falcate.

10. *C. Bolusii*, sp. nov.

Rami villosi. *Foliola* 1.2 cm. longa, apice mucronata, marginibus revolutis, villosa. *Bractae* 4 mm. longae, ovatae, acuminate, villosae. *Pedicellus* 7 mm. longus. *Calyx* 4 mm. longus; lobi falcati, villosi.

Oudtshoorn Division : In stony places at summit of the Zwartberg Pass, December, *Bolus* 11465.

A small-branched villous shrub. *Leaflets* 1.2 cm. long, mucronate at the apex, with strongly revolute margins, villous. *Bracts* 6 mm. long, broadly ovate, acuminate, villous. *Pedicels* 7 mm. long. *Calyx* 4 mm. long; lobes broadly falcate.

11. *C. sessiliflora* (E. Mey. Com., p. 4).

Young branches and leaves densely villous, at length becoming glabrous. *Leaflets* 1.25 cm. long, linear, with strongly revolute margins, subsessile. *Bracts* 6 mm. long, broadly ovate, acuminate. *Flowers* ciliate. *Calyx* softly villous; lobes falcate.—“*Flora Capensis*,” II, 8.

Caledon Division : Genadendal, December, *Schlechter* 9826.

12. *C. montana*, sp. nov.

Rami juniores sericei. *Foliola* 1.25–2.5 cm. longa, 2 mm. lata, linearia, basi cordata, marginibus revolutis. *Bractae* 4–6 mm. longae, ovatae, glabrae. *Calyx* sericeous, 8 mm. longus; lobus inferius ceteris major.

Tulbagh Division : Sneeuwgat Valley, Great Winterhoek, November, *Phillips* 1693.

Var. *glabra*. Calycis glabris difert.

Ceres Division : Matroosberg, December, *Phillips* 1943.

Young branches and leaves silky, at length becoming glabrous. *Leaflets* 1.25–2.5 cm. long, 2 mm. broad, linear, cordate at the base, with revolute margins. *Flowers* subsessile. *Bracts* 4–6 mm. long, broadly ovate, glabrous, longer than the pedicel. *Calyx* 8 mm. long, densely silky; the lower segment largest and broadest; the middle segments shortest, subacute, broadly oblong ciliate.

Var. *glabra* differs in having a glabrous calyx.

CONTENTS.

	PAGE
REVISIÓN OF SOUTH AFRICAN SPECIES OF RHYNCHOSIA.....	113
THE GENERA ALOE AND MESEMBRYANTHEMUM.....	139
SOUTH AFRICAN GRAMINEAE.....	170
A FUNGUS OF ECONOMIC IMPORTANCE ON THE AVOCADO.....	179
SPECIES OF ELEPHANTORRHIZA IN THE SOUTH AFRICAN HERBARIA	187

REVISION OF SOUTH AFRICAN SPECIES OF RHYNCHOSIA.

By EDMUND G. BAKER, F.L.S., Assistant Keeper, Department of Botany, Natural History Museum, South Kensington.

INTRODUCTION.

At the request of Dr. I. B. Pole-Evans, C.M.G., and Dr. E. P. Phillips, I have examined the collection of the genus *Rhynchosia* from the National Herbarium at Pretoria, and I have also had an opportunity of seeing the plants of this genus from the Herbaria at Capetown and Durban, thanks to the courtesy of the Curators of these collections. I am also much indebted to the Director of the Botanical Museum at Upsala for allowing me to have the loan of Thunberg's types, and to Dr. Schinz for kindly sending me portions of types of many of his species.

Since the publication of the *Leguminosae* by Harvey in the "Flora Capensis," in 1861-62, a considerable amount of additional information has been acquired, and I am glad to take the opportunity offered to me of placing this on record. It is, perhaps, unnecessary to discuss in detail publications anterior to the "Flora Capensis." De Candolle, in the second volume of his "Prodromus," in 1825, described several species of *Rhynchosia* from South Africa, among these being *R. angustifolia* founded on *Glycine angustifolia* Jacq. and *R. rigidula* founded on Burchell No. 2587. The first of these is synonymous with, and must take precedence of, *R. uniflora*, Harvey, the second is evidently closely allied to *R. Totta*, D.C., but there is a mistake in the Burchell number, and I am unable to find it in the Kew Herbarium. Perhaps the other most important publications anterior to the "Flora" are E. Meyer's "Commentorium de Plantis Africae australioris" and Ecklon and Zeyher's "Enumeration." Subsequent to the "Flora" a good many species have been described, and a few preliminary notes on some of these novelties may not be out of place. In the "Journal of Botany," Vol. XVI, 131 (1878), Mr. Spencer Moore described *R. divorum* from near Pilgrims Rest. It has been subsequently found in Gazaland by C. F. M. Swynerton. It is a curious species with erect stem, showy flowers, and with the upper calycine lobes shorter than the lowest.

In 1888 Dr. Schinz described, in "Verhand. Bot. Brand.," Vol. XXX, two species, *R. hirsuta*, from Olukonda in Amboland, and *R. longiflora* from Great Namaqualand, the latter being allied to *R. Totta*, D.C., but with much finer flowers, and in 1894 the same author described in "Bull. Herb. Boiss.," Vol. II, *R. Woodii*, from Natal, an ally of *R. Orthodanum*, Benth. Dr. Harms in 1899, in Engler's "Bot. Jahrb.," Vol. XXVI, described *R. komatiensis* from between Spitzkop and the Komati River, and *R. longipes*, founded on Wilms No. 395, from Lydenberg. I have compared the latter with the type of *R. crassifolia*, Benth., and do not consider it specifically distinct. In 1897, in the "Journal of Botany," Dr. Schlechter described the very distinct *R. monophylla*, and in the same journal, in the same year, Messrs. Wood and Evans described the equally distinct *R. ovata*. In 1905 Dr. Schlechter, in "Ann. Nat. Hoffmus. Wien," described three species, *R. chrysantha*, *R. Harmsiana*, and *R. Penthieri*. In 1907, in the "Vierteljahrschrift Nat. Ges. Zurich," Dr. Schinz described *R. cinnamomea*, *R. congestiflora*, *R. elegantissima*, *R. Fleckii*, *R. namaensis*, and *R. Rehmanni*, and in 1910

the same author, in the same publication, described the very distinct *R. spectabilis*. I am doubtful whether *R. cinnamomea* is distinct from *Dolicholus venulosus*, Hiern. (*R. venulosus*, K. Schum.), and I think *R. Rehmanni* is synonymous with *Fagelia bituminosa*, D.C. In 1906 Dr. Bolus, in "Trans. S. Afr. Phil. Soc.," XVI, 383, described the curious and aberrant *R. pauciflora*. I hardly think *R. uncinata*, Schlechter, from Packhuis Berg, is separable from *R. bullata*, Benth. In 1913, in "Bull. Soc. Bot. Fr.," LX, 461, Gandoger quite correctly segregated the *R. memmonia* of Harvey, calling the silvery plant with a rigid stem *R. albissima*.

Several plants have been issued with ms. names to which I must allude. Wilms No. 375 I would place as synonymous with *R. monophylla*, Schlechter; Wilms No. 387 appears to me to be a form with more angled leaves than usual of *R. sigmodes*, Benth. I have retained *R. graciliflora*, Harms, founded on Wilms No. 367, as a variety of *R. Totta*, D.C.; it differs in the shape of the leaflets from the typical form. Quite recently in the "Kew Bulletin" Mr. Burt-Davy described *R. unifoliolata*. It is also a close ally of the true *R. Totta*, D.C., but is sometimes unifoliolate, and has shorter petioles. A little later in the same publication he also describes *R. confusa*, *R. Burkei*, *R. Harmsiana* Schl., var. *Burchellii*, and *R. Pentheri* Schl., var. *Hutchinsoniana*. With *R. Burkei* and *R. Pentheri*, Schl., var. *Hutchinsoniana*, he has associated my name.

NOTES ON TAXONOMY.

Harvey, in the "Flora Capensis," arranges the species of *Rhynchosia* then known to him in four sections: (i) **Chrysoscias**; (ii) **Polytropia**; (iii) **Copisma**; and (iv) **Orthodanum**. The first of these is well marked by the character of the inflorescence, which is generally umbellate, more rarely the flowers are solitary. The leaflets are narrow and either tomentose or villose below. This section I retain as originally delimited.

The section **Polytropia** is at once recognized by having pinnate or bipinnate leaves. But there is a difficulty with regard to the sections **Copisma** and **Orthodanum**. Certain plants in these sections are very distinct from one another; take, for instance, *R. adenodes*, Ecklon and Zeyher, and *R. minima*, D.C., with a distinctly voluble stem, which is the main character of Harvey's **Copisma**, and *R. Orthodanum*, Benth., which is erect, rigid, and suffruticose, and which is the type of Harvey's section **Orthodanum**. But there are certain intermediate plants which are difficult to place. Take, for instance, *R. viscidula*, Steud.: the plant from Little Namaqualand has a somewhat rigid stem and cannot be correctly placed in **Copisma**. I have, therefore, thought it advisable to unite the sections **Copisma** and **Orthodanum**, and am calling this section **Eurhynchosia**.

I have introduced the section **Cyanospermum**, Benth., for *R. cyanosperma*, Benth. It may be easily recognized by the robust voluble stem, the racemose inflorescence, the broad calycine segments, which are densely matted on the back with a grey tomentum, and the dark blue seeds. I am also introducing the section **Arcyphyllum** Torrey and Gray for two species which occur south of the Tropic. The stem in this section is voluble, the flowers are in dense, oblong, sessile, subsessile, or pedunculate clusters, and with the calycine teeth much narrower than in section **Cyanospermum**.

There are one or two anomalous species which require consideration. One of these is *R. pauciflora*, Bolus, with solitary pedicellate flowers and broad calycine lobes. It comes from the Transvaal and Swaziland. *R. ficifolia*, Benth., I am excluding from the genus. It is a *Pueraria*. Some species from Great Namaqualand such as *E. namaensis*, Schinz, I have not included, as it is recorded from Rehoboth, which is just north of the Tropic.

DESCRIPTION OF SPECIES.

SECT. I.—CYANOSPERMUM, Benth., in Benth. and Hook "Gen. Plant" (1865), 543. *Stem* voluble, robust. *Inflorescence* racemose. *Calyx* as long as or slightly longer than the corolla, segments rather broad, subequal in length, uppermost connate. *Seeds* dark blue.

1 **R. cyanospermum**, Benth., *ex* Baker in Oliver, "Fl. Trop. Afr.," II, 218.

Stems voluble, robust, densely grey or ferruginous-pubescent when young. *Stipules* ovate, pubescent. *Leaves* trifoliate; leaflets coriaceous; terminal leaflet ovate or sub-orbicular, below densely grey-tomentose; lateral leaflets unequalsided. *Flowers* rather large, in racemes. *Bracts* ovate, deciduous. *Calyx* 18–20 mm. long or rather more, densely matted on the back with grey tomentum; tube short; lobes uppermost, somewhat spatulate, the others blunt, 4–5 mm. broad. *Corolla* about as long as the calyx; standard obovate; carina boat-shaped, base unguiculate. *Pod* oblong, equalling the calyx; seeds dark blue.

NATAL.—Lower Illovo, Wood No. 6507, Natal Herb.

SECT. II.—ARCYPHYLLUM, Torr. and Gray, "Fl. North. America" (1838), 284.

Stem voluble. *Flowers* in dense oblong, sessile, or subsessile, or shortly-stalked clusters. *Calyx teeth* narrow, sometimes more or less connate, slightly shorter than, or same length as, the corolla. *Seeds* brown or black, not blue.

+ *Terminal leaflet ovate, subrotund, acuminate.*

2. **R. densiflora**, D.C., "Prod.," II, 386; Baker in Hooker, "Fl. Brit. India," II, 226; Baker *ex* Oliver, "Fl. Trop. Afr.," II, 222.

Glycine densiflora, Roth., nov. sp. 348; *Desmodium punctatum*, D.C., "Prod.," II, 338.

Stem slender, firm-herbaceous, voluble, grey pilose. *Stipules* lanceolate. *Leaves* trifoliate, papyraceous; terminal leaflets broadly ovate acuminate; petiolules of terminal leaflets 10–18 mm. long; lateral leaflets unequalsided. *Flowers* in dense, oblong, subsessile clusters. *Bracts* lanceolate. *Calyx* 12–15 mm. long, teeth narrow, subequal, two uppermost more or less connate; tube short. *Standard* glabrous, auriculate, about 15 mm. long; wings unguiculate, distinctly shorter than the carina; carina boat-shaped, obtuse. *Pod* 12–14 mm. long, rather narrowed at both ends, grey pubescent.

NATAL.—Gerrard No. 1737; Camperdown Div., Wood No. 11706.

TRANSVAAL.—Makapansberg, Rehmann No. 5548; Potgietersrust, Leendertz No. 146, 1216; Barberton, Thorncroft No. H. 11274; Pienaars River, Leendertz No. 783.

+ + *Terminal leaflet rhombeo-ovate, generally obtuse.*

3. **R. connata**, sp. nov.

Caulis volubilis, pubescens, longitudinaliter striatus. *Stipulae* ovatae, acutae; stipellis ad basin petiolorum minutis linearibus. *Folia* trifoliolata; foliolis terminalibus rhombeo-ovatis apice obtusis vel rarius subacutis, 3–4 cm. longis et latis; lateralibus inaequilateralibus, petiolo communi 4–5 cm. longo praedita; foliolis terminalibus petioulis 8–10 mm. longis suffultis. *Flores* flavi in capitula densa disposita; capitulis breviter pedunculatis. *Calyceis lobis* 2 superioribus plus minus connatis, reliquis lineari-lanceolatis. *Vexillum* apice emarginatum; alae oblongae basi unguiculatae; carina naviculariformis apice obtusa. *Ovarium* pilis vestitum; stylo gracili. *Legumen* ignotum.

NATAL.—Camperdown, Miss Franks No. 1250.

Ally of *R. Stuhlmanni*, Harms in Engler's "Bot. Jahrb.," XXVI, 308.

SECT. III.—CHRYSOSECIAS, Benth., *ex* Harvey and Sonder, "Fl. Cap.," II, 249.

Stem voluble. *Leaves* pinnately trifoliate. *Stipules* broad. *Flowers* in axillary umbels or solitary.

Series 1.—*Calyx* large, 15–18 mm. long. *Flowers* in umbels rarely solitary. *Underside of leaf fulvous-villose.*

4. **R. chrysoscias**, Benth., *ex* Harvey and Sonder, "Fl. Cap.," II, 249; Oliver in "Bot. Mag.," t. 5913.

Chrysoscias grandiflora, E. Mey., "Comm.," 139; *Glycine erecta*, Thunbg., "Fl. Cap.," 592; *Cylista lancifolia*, Ecklon and Zeyher, Herb. No. 1690.

A climbing suffrutex with voluble stem, and the younger parts clothed with tawny or yellowish hairs. *Stipules* broad. *Leaves* pinnately trifoliolate, thinly pubescent above, tomentose or villose below; terminal leaflets oblong-lanceolate with revolute margins 3.5–5 cm. long, 8–15 mm. broad; petioles 5–15 mm. long. *Peduncles* bearing 3–4 flowers at the summit. *Calyx* 15–18 mm. long; tube short; segments lanceolate; the two upper somewhat connate at the base. *Pod* not much longer than the calyx, pilose.

CAPE.—Districts George and Knysna, Plettenbergs Bay, Zeyher No. 1690.

Chrysoscias media, E. Mey., from Kaymans Gat, is a close ally.

Series 2.—*Calyx* silky-silvery, 15–20 mm. long, upper lobes connate to or beyond the middle. *Underside of leaf* white tomentose.

5. *R. leucoscias* Benth., ex Harvey, l.c. II, 249.

Cylista argentea, Ecklon and Zeyher, 1688; *Cylista angustifolia*, E. Mey., in "Linnaea," VII, 171; *Chrysoscias calycina*, E. Mey., "Comm.," 140.

Suffruticose voluble, albo-villose. *Stipules* broadly oblong, obtuse or subacute. *Leaves* trifoliolate, very shortly petiolate; terminal leaflets with revolute margins, oblong-lanceolate or linear, white tomentose below, 4–5.5 cm. long, 5–7 mm. broad in the middle; common petioles 2–4 mm. long. *Peduncles* umbelliferous. *Calyx* silky-silvery; tube very short; segments lanceolate, leafy, silky and silvery. *Corolla* two uppermost segments connate to or beyond the middle, glabrous; standard 14–16 mm. long. *Pod* not seen.

CAPE.—Van Stadensberg, Ecklon and Zeyher.

Var. *angustifolia*, Harvey, l.c. 249.

Leaflets narrower, narrow-linear. *Flowers* sub-solitary.

CAPE.—Caledon Div., River Zondereinde, Zeyher No. 2410.

Distinguished from *R. chrysoscias*, Benth., by its much whiter and woolly tomentum and the connate upper calyx lobes.

Series 3.—*Calyx* cano-pubescent, 6–9 mm. long; upper lobes connate to middle. *Umbels* 4–8-flowered.

6. *R. microscias*, Benth., ex Harvey, l.c. 249

Chrysoscias parviflora, E. Mey., "Comm.," 139.

Suffruticose voluble, silky-canescens. *Stipules* ovate-lanceolate, 4–6 mm. long. *Leaves* trifoliolate, on very short petioles; terminal leaflets linear-lanceolate or linear, 3–4 cm. long, 4–7 mm. broad, canescent and tomentulose beneath; common petioles 2–4 mm. long; lateral leaflets slightly unequaled at the base. *Peduncles* bearing umbels of 4–8 flowers. *Flowers* much smaller than in *R. leucoscias*, Benth. *Calyx* tube very short; segments lanceolate, cano-pubescent, shorter than the standard. *Standard* 10–12 mm. long, glabrous; carina glabrous, obtuse. *Pod* pilose, 15–20 mm. long, mucronate.

CAPE.—Mountain sides, near George, Drège; Knysna, Tyson, Bowie; Humansdorp Div., Zitzikama, E. Galpin No. 3989.

Known at once from the preceding by the smaller flowers and shorter indumentum.

Series 4.—*Flowers* solitary.

7. *R. angustifolia*, D.C., "Prod.," II, 388.

R. uniflora, Harvey, l.c. II, 249; *Glycine angustifolia*, Jacq., "Hort. Schoenb.," II, t. 231; *Cylista angustifolia*, Ecklon and Zeyher No. 1689.

Stem voluble, silky-canescens. *Stipules* ovate, subacute. *Leaves* trifoliolate, very shortly petiolate; leaflets lanceolate or linear, with revolute margins, whitish or tomentose below; terminal leaflets 15–25 mm. long and about 2–3 mm. broad; terminal petiolule silky canescent, 2–3 mm. long; lateral leaflets shorter than terminal. *Flowers* solitary,

axillary, on short pedicels. *Calyx* 13–15 mm. long, grey-villous; tube very short; segments lanceolate, not much shorter than the petals; two uppermost somewhat connate. *Standard* glabrous, 13–15 mm. long. *Pod* not seen.

CAPE.—Purpas Valley and Zwarteberg, near Caledon, *Ecklon* No. 1689.

This plant is easily recognized by the voluble stem, the very narrow trifoliate leaves, and the solitary axillary flowers with a silky villous calyx. It must take De Candolle's earlier name of *R. angustifolia*.

SECT. IV.—POLYTROPIA, Harvey and Sonder, "Fl. Cap.," II, 248.

Stem prostrate or trailing. Leaves pinnate or bipinnate, plurijugate. Flowers racemose.

Series 1.—Leaves pinnately decompose.

8. ***R. ferulaefolia***, Benth., ex Harvey, l.c.

Polytropia ferulaefolia, "Presl. Symb.," t. 13; *Psoralea prostrata*, "Linn.," sp. pl. 762; *Galega pinnata*, Thunberg, "Fl. Cap.," 602; *Polytropia umbellata*, Ecklon and Zeyher No. 1627.

Prostrate, nearly glabrous. *Stipules* rather short. *Leaves* pedately bipinnate, paucijugate or supra-decompound; leaflets sessile, narrow-lanceolate or linear, acute at each end, varying from about 5–20 mm. long; common petiole generally 2–4 cm. long, glabrous or nearly so. *Peduncles* longer than the leaves, bearing a cluster of 5–12 flowers towards the apex; pedicels 1–3 mm. long. *Flowers* viscoso-pubescent. *Calyx* with lowest segment conspicuously narrower than the rest. *Standard* 7–9 mm. long. *Pod* 12–16 mm. long, subfalcate or nearly straight, nearly glabrous.

CAPE.—Common on the Cape Flats and in several parts of the western districts. There is a specimen of this in Herb. Sloane, from the Cape, gathered by Oldenland. This will certainly be one of the earliest records for this species.

Series 2.—Leaves simply pinnate.

9. ***R. pinnata***, Harvey, "Thes.," t. 79; Harvey in Harvey and Sonder, l.c. 250.

Polytropia pinnata, Ecklon and Zeyher No. 1628.

Whole plant minutely downy, prostrate. *Stipules* ovate, striate, deflexed. *Leaves* simply pinnate, bi- to tri-jugate; terminal leaflet rhomboid-ovate or obovate, 6–12 mm. long; lateral leaflets generally elliptic-oblong, impresso-punctate. *Peduncles* elongate, 5–6-flowered near the summit. *Calyx* pubescent, 6–7 mm. long; lowest segment narrower than the others, shorter than the petals. *Standard* yellow, 6–8 mm. long, glabrous or nearly so; carina about same length as the standard. *Pod* not seen, stated by Harvey to be the same as in *R. ferulaefolia*, Benth.

CAPE.—Saldanha Bay, *Ecklon and Zeyher* No. 1628.

SECT. V.—EURHYNCHOSIA (incl. *Copisma* and *Orthodanum*).

Stem voluble or rigid. *Leaves* trifoliate or rarely unifoliate. *Inflorescence* generally racemose, but flowers sometimes solitary or subsolitary. *Calyx* generally shorter than the corolla; segments generally lanceolate. *Seeds* brown, or reddish-brown, or black, not blue (in those species in which the seeds are known).

CLAVIS OF SPECIES.

A.—Stems voluble or subvoluble.

(a) Bracts persistent.

—Leaves trifoliate.

Glabrous or nearly so.....

Hairy.....

— —Leaves unifoliate.....

R. rotundifolia, Walp.

R. grandifolia, Steud.

R. simplicifolia, E. Mey.

- Leaflets in typical form, ovate-oblong or elliptical, in forma *angustifolia*, lanceolate-oblong. Peduncles 1-3-flowered
 Leaflets narrow-lanceolate. Peduncles 2-flowered. Flowers larger than in *R. Totta*, D.C. *R. Totta*, D.C.
 Leaflets elliptical. Racemes very lax, few-flowered. Stem voluble. Indumentum cinnamon coloured. *R. longiflora*, Schinz.
 Leaflets lanceolate, more or less sericeous. Raceme very lax. Stem voluble. *R. venulosa*, K. Schum. (*R. cinnamomea*, Schinz.)
 Leaflets oblong-lanceolate or ovate. Peduncles 1-6-flowered near the summit. Stem voluble, glabrous, or minutely downy
 Leaflets linear or linear-lanceolate. Peduncles generally one-flowered. *R. elegantissima*, Schinz.
 *R. glandulosa*, D.C.
 *R. rigidula*, D.C.
- B.—Stem rigid or subrigid.
- ++ Peduncles generally shorter than the leaves, few-flowered.
 Silky villous or glabrous. Leaflets elliptic-oblong or lanceolate-oblong
 Canotomentose. Terminal leaflets broader than long.... *R. Orthodanum*, Benth.
 Cinnamon-coloured tomentum. Terminal leaflets oval or oval-suborbicular. *R. spectabilis*, Schinz.
 Pubescent. Terminal leaflets ovate or orbicular-ovate
 Velvety when young. Terminal leaflets ovate-lanceolate. Flowers large. *R. Rudolphi*, Harms.
 Silky silvery. Leaflets broadly ovate or cordate-ovate. Flowers 16-18 mm. long. *R. komatiensis*, Harms.
 Silky silvery. Leaflets larger, oblong-obovate. Flowers 20-23 mm. long. *R. clivorum*, S. Moore.
 *R. nitens*, Benth.
 *R. Galpini* Bak. fil.
- +++ Flower pedicellate.
 Terminal leaflets rhombic-ovate or suborbicular-ovate. Flowers 17-19 mm. long. *R. Bakeri*, Schinz.
- ++++ Peduncles scarcely longer than the leaves, 1-2-flowered.
 Shrubby, rigid, viscidly hairy and resin-dotted.
 Flowers 13-16 mm. long
 *R. bullata*, Benth.
 (*R. uncinata*, Schltr.)
- ++++ Peduncles about as long as the leaves, or slightly longer, several to many flowered. Flowers 8-16 mm. long.
 Hairy and somewhat silvery. Terminal leaflets elliptical. Flowers 14-16 mm. long
 Pubescent. Terminal leaflets oblong-obovate. Flowers 11-13 mm. long. *R. angulosa*, Schinz.
 Canotomentose. Terminal leaflets obovate. Flowers 8-12 mm. long. *R. Woodii*, Schinz.
 Pubescent. Terminal leaflets rhombic-ovate. Flowers 10-11 mm. long. *R. albissima*, Gandoger.
 *R. Pegleri*, Bak. fil.
- ++++ Peduncles longer than the leaves, always one-flowered.
 Leaflets rigid, narrow, margins distinctly revolute. *R. pauciflora*, Bolus.
- ++++ Peduncles longer than the leaves, few-flowered glandular plants.
 Lateral leaflets obtuse. *R. viscidula*, Steud.
 Lateral leaflets ovate-lanceolate, acute. *R. Schlechteri*, Bak. fil.

A.—STEM VOLUBLE. BRACTS PERSISTENT.

Series 1.—Leaves trifoliolate, leaflets small, orbicular or ovate.

10. **R. rotundifolia**, Walp. Rep. I, 787; Harvey, l.c. 250.

Copisma rotundifolium, E. Mey., "Comm.," 137.

Prostrate. Stem flexuous, angled. *Stipules* small, striate, about 4 mm. long. *Leaves* trifoliolate, on rather long filiform petioles; leaflets ovate or orbicular, netted-veined and resin-dotted, glabrous or nearly so; terminal leaflets 10-13 mm. long, 9-11 mm. broad; common petiole 3-4 cm. long. *Peduncles* elongate, densely racemose near the summit. *Flowers* rather large, 12-14 mm. long. *Bracts* broadly ovate, persistent. *Calyx* 5-6 mm. long; lowest segment longest, subglabrous. *Corolla* glabrous; standard broadly ovate or orbicular; keel broadly falcate. *Pod* not seen.

CAPE.—Pondoland : Omsamcaba, *Drège*.

Known by the broadly ovate persistent bracts, the trifoliolate leaves on rather long filiform petioles.

Series 2.—Leaves trifoliolate or unifoliolate ; leaflets rather large, pubescent.

+ *Leaves trifoliolate.*

11. **R. grandifolia**, Steud., "Nom., ed. 2, II, 454 ; Harvey, l.c. 250.

Copisma grandifolium, E. Mey., "Comm.," 138.

Procumbent robust hirsuto-pubescent. *Stipules* broadly cordate. *Leaves* trifoliolate, subsessile or petiole very short ; terminal leaflets ovato-subrotund, hispid-pubescent, 5–6 mm. long, 4–5.5 cm. broad ; lateral leaflets unequalsided. *Peduncles* elongate, longer than the leaves, closely spicate-racemose beyond the middle. *Bracts* ovate-oblong, acute, persistent, hairy. *Calyx* segments lanceolate ; uppermost semiconnate ; lowest subulate. *Corolla* glabrous ; standard striped on the back ; wings shorter than the standard ; keel boat-shaped. *Young pod* hirsute.

CAPE.—Near Kat River, *Drège* ; at foot of Winterberg, *Mrs. Barber* ; Kingwilliamstown, *Flanagan* No. 2137 ; Kreil's Country, *Bowker*.

Known by the elongate peduncle spicate-racemose beyond the middle, the rather showy flowers, and persistent ovate-oblong acute bracts.

+ + *Leaves unifoliolate.*

12. **R. simplicifolia**, E. Mey., "Comm.," 138 ; Harvey, l.c. 251.

Described as being very closely allied to *R. grandifolia*, Walp., but the leaves are simple. *Stipules* cordate-ovate, acuminate. *Bracts* ovate-lanceolate.

TRANSKEI.—Umtata, *Drège*.

Only known to me from E. Meyer's description. It is possibly a unifoliolate form of *R. grandifolia*, Walp.

B.—STEM VOLUBLE. BRACTS CADUCOUS. FLOWERS FASCICULATE.

Series 3.—Leaflets unifoliolate or trifoliolate, suborbicular, or ovate, or ovate-oblong, puberulous. *Flowers* 12–14 mm. long.

13. **R. monophylla**, Schlechter in "Journ. Bot." (1897) 279.

Stem voluble, branches decumbent, filiform, elongate. *Leaves* simple or trifoliolate, suborbicular, or ovate, or ovate-oblong, reticulate nerved, 2.5–4 cm. long, 2–3 cm. broad. *Flowers* in axils of leaves, fasciculate. *Bracts* deciduous. *Calyx* $\pm$ 6 mm. long ; segments linear, acute. *Standard* suborbicular-rhomboid, $\pm$ 11 mm. long, 9 mm. broad ; wings unguiculate ; keel obtuse, apex incurved, glabrous. *Ovary* villous. *Pod* obliquely ovate-elliptical, very compressed, base acute, apex acuminate, $\pm$ 11 mm. long, 5–6 mm. broad.

TRANSVAAL.—Near Barberton, *E. E. Galpin* No. 489 ; Pretoria, *Rehmann* No. 4582 ; *Rogers* No. 18942 ; *A. O. D. Mogg*, in Govt. Herb., No. 10556 ; near Witbank Station, *Gilfillan* No. 7184 ; *Lydenberg*, *Wilms* No. 375 ; near Barberton, *Bolus* Nos. 1142 and 7603.

Wilms No. 375 was issued with a ms. name, but it is not specifically distinct from the above.

Occurs also in Gazaland.

Var. nov. **Eylesii**, Bak. fil.

Caulis procumbens, pubescens. *Stipulae* latae, ovato-lanceolatae, 5–7 mm. longae. *Folia* saepissime trifoliolata, suprema unifoliolata, foliolis terminalibus late ovatis vel suborbiculari-ovatis, 2.5–4 cm. longis, 2.5–3 cm. latis ; petiolulis foliolorum terminalium 8–10 mm. longis ; foliolis lateralibus inaequilateralibus. *Flores* fasciculati. *Vexillum* flavum, $\pm$ 12 mm. longum. *Legumen* immaturum pilis vestitum.

RHODESIA.—Salisbury, *F. Eyles* No. 2094. Alt. 4900 ft.

This is a procumbent herb with rather broad stipules. It differs principally from the type in the leaves being mostly trifoliolate. The standard is yellow and the wings bright red. The calyx is ± 13 mm. long (total length), the lowest segment narrow and much longer than the others. Calyx tube, ± 3 mm.

C.—STEM VOLUBLE OR SUBVOLUBLE. BRACTS CADUCOUS, RARELY SUBPERSISTENT. LEAFLETS BROAD. FLOWERS RACEMOSE.

Series 4.—*Leaflets small. Peduncles nearly as long as or slightly longer than the leaves. Flowers small, in unilateral racemes.*

+ *Leaflets softly pubescent or velvety. Stipules broadly ovate.*

14. **R. Harveyi.** (*R. secunda* Harvey, l.c. ii. 251.)

Prostrate softly pubescent or velvety. Branches flexuous. Leaves trifoliolate, petiolate; terminal leaflets orbicular-rhomboid, obtuse, 10–15 mm. long, 12–17 mm. broad; lateral leaflets unequalsided. Peduncles shorter or scarcely longer than the leaves, laxly 2–4-flowered. Calyx 6–7 mm. long; segments longer than the tube, shorter than the glabrous standard. Standard ± 9 mm. long. Keel ± 9 mm. long. Pod 15–18 mm. long, narrowed towards the base, pubescent.

CAPE.—Smaldeel, *Zeyher* No. 514; near Grahamstown, *P. MacOwan* No. 160.

Flanagan No. 164 from Komgha is a form of this species.

++ *Leaflets fulvous, resin-dotted below.*

15. **R. secunda** (*Glycine secunda*, Thunberg, "Prod. Fl. Cap.," 131).

Stem herbaceous, filiform, pubescent. Stipules narrower than in the preceding species, lanceolate, longitudinally striate. Leaves trifoliolate, petiolate; terminal leaflets rhomboid-ovate, 10–15 mm. long, and nearly the same breadth, rounded at the apex; petiolules 2–3 mm. long; lateral leaflets unequalsided, sessile or subsessile, all resin-dotted below; petioles pubescent, 8–10 mm. long. Flowers small, in 5–10-flowered secund racemes, which are often 3–4 cm. long. Pedicels very short. Calyx 6–7 mm. long, hairy, the lowest segment longer than the rest. Standard 6–8 mm. long, glabrous; keel about the same length. Pod not seen.

CAPE.—In campis graminosis, *Thunberg*, Herb. Thunberg.

This is a rare plant, the only specimen I have seen being in Herb. Thunberg.

Series 5.—*Leaves trifoliolate, terminal leaflets broadly obovate. Peduncles short, 5–7-flowered.*

16. **R. Burkei**, Burt-Davy and Bak. fil. in "Kew Bull.," 1921, 193.

R. adenodes, Ecklon and Zeyher, var. ? *robusta*, Harvey, l.c.

Stems prostrate, robust, longitudinally striate, pubescent. Stipules lanceolate. Leaves trifoliolate, petiolate; terminal leaflets broadly obovate, 20–25 mm. long, 23–28 mm. broad; terminal petiolules 6–8 mm. long, on both sides except the nerves nearly glabrous; lateral leaflets unequalsided, sessile; common petiole 15–25 mm. long. Peduncles laxly 5–7-flowered, generally 25–30 mm. long. Flowers small, 7–8 mm. long. Calyx pubescent, + 4 mm. long. Standard 6–7 mm. long. Pod pubescent, about 12 mm. long.

ORANGE FREE STATE.—Thaba 'Nchu, *Zeyher* No. 280; *Burke* in Herb., Kew.

This is quite distinct from *R. adenodes*, Ecklon and Zeyher. It has broad, obovate terminal leaflets and a short inflorescence of quite small flowers. It does not seem to be common.

Series 6.—*Leaves trifoliolate, terminal leaflets ovate or obovate. Racemes short, few-flowered.*

+ *Terminal leaflets obovate, 20–25 mm. broad.*

17. **R. Fleckii**, Schinz in "Vierteljahrsschrift Nat. Ges. Zurich," LII, 428 (1907).

Stem slender, filiform, pubescent. *Stipules* small, linear-lanceolate, pubescent. *Leaves* trifoliolate, petiolate, grey pubescent; terminal leaflets obovate, about 25 mm. long and 20–25 mm. broad; terminal petiolules pubescent, 8–10 mm. long; lateral leaflets somewhat unequalsided, also about 25 mm. long, petiolules much shorter. *Racemes* short, few-flowered. *Calyx* grey pubescent, segments narrow-lanceolate, shorter than the corolla. *Standard* 7–9 mm. long; keel boat-shaped, apex blunt. *Pod* pubescent, mucronate, 23–25 mm. long, 8–9 mm. broad at the broadest point.

KALAHARI REGION.—Uschi, *Fleck* No. 693.

Noticeable on account of the slender stem, small stipules, trifoliolate leaves covered with grey pubescence, obovate terminal leaflets, and short few-flowered inflorescence. An ally of *R. confusa*, Burt-Davy.

++ *Terminal leaflets narrower, 12–16 mm. broad.*

18. **R. confusa**, Burt-Davy, in "Kew Bull.," 1921, 192.

R. nervosa, Benth., var. *pauciflora*, Harvey, l.c.

Perennial prostrate herb, with puberulent stems. *Stipules* subulate. *Leaves* trifoliolate, cano-pubescent; terminal leaflets ovate or obovate, longer than broad, 13–20 mm. long, 12–16 mm. broad; petiolules about 5 mm. long; lateral leaflets unequalsided, common petioles tomentose, 5–10 mm. long. *Peduncles* slender, 2–4-flowered, shorter than the leaves or only slightly longer. *Pedicels* slender, 1–2 mm. long. *Flowers* about 8 mm. long. *Calyx lobes* lanceolate, acuminate, pubescent. *Standard* glabrous. *Pod* puberulous, about 2 cm. long and 7 mm. broad.

TRANSVAAL.—Magalies River, *Zeyher* No. 510; Aapjes River and Vaal River, *Zeyher* Nos. 508, 511.

ORANGE FREE STATE.—Kroonstad Dist., Rhenoster Kop, *Zeyher*; Bloemfontein, *Burt-Davy* No. 11767.

BRITISH BECHUANALAND.—Kuruman Dist., *Burchell* Nos. 2298, 2457, 2548B; Vryburg Div., near Geluk, *Burt-Davy* No. 11121.

This species is easily recognized by its trifoliolate cano-pubescent leaves and short 2–4-flowered inflorescence generally shorter than the leaves.

It also occurs in Rhodesia, Victoria Div., *Munro* No. 549.

Series 7.—*Flowers 12–15 mm. long in lax racemes. Terminal leaflets ovate, triangular, acute, or broadly rhomboid. Stem tomentulose.*

19. **R. quadrata**, Harvey, l.c. ii, 254.

Stem herbaceous, flexuous, thinly tomentulose. *Stipules* small, lanceolate. *Leaves* thin, trifoliolate, petiolate; leaflets broadly rhomboid-acute or subacuminate; terminal leaflets 20–35 mm. long, 20–38 mm. broad; terminal petiolule 5–8 mm. long, lateral leaflets 20–35 mm. long, 20–38 mm. broad; terminal petiolule 5–8 mm. long. *Racemes* shorter or a little longer than the leaf, 5–8-flowered. *Bracts* narrow, caducous. *Calyx* glandular, 14–15 mm. long; lowest lobe longest and nearly same length as carina. *Standard* glabrous or nearly so, 13–15 mm. long. *Ovary* glandular, biovulate. *Pod* substipitate, falcate, twice as long as calyx.

NATAL.—Port Natal, *T. Williamson*, *Peddie*; Durban, *Wood* No. 1656.

Easily recognized by its thin, rhomboid, acute leaflets, and glandular calyx and ovary.

Series 8.—*Leaves trifoliolate, leaflets ovate or broadly ovate. Peduncles longer than the leaves, towards the apex, lastly 2–6-flowered.*

+ *Terminal leaflets 3–4.5 cm. long, 2.5–4.0 cm. broad.*

20. *R. reptabunda*, N. E. Br. in "Kew Bull." 1906, 103.

Stem herbaceous, creeping, pubescent or pilose. *Stipules* rather large, 7–11 mm. long. *Leaves* trifoliate, petiolate; leaflets ovate or broadly ovate, 3–4.5 cm. long, 2.5–4 cm. broad, on both sides pubescent; common petiole 3–6 cm. long, puberulous and laxly pilose. *Peduncle* longer than the leaves, 5–17.5 cm. long, towards the apex laxly 2–6-flowered. *Bracts* lanceolate, 4–5 mm. long. *Pedicels* $\pm$ 3 mm. long. *Calyx* pubescent; tube 3 mm. long, upper teeth 3–4 mm. long, lowest lanceolate-subulate, $\pm$ 7 mm. long. *Corolla* glabrous, yellow; standard 8.5 mm. long, broadly oblong; keel about the same length as the standard, obtuse. *Ovary* densely tomentose.

NATAL.—Near Charlestown, Wood No. 5734; Giant's Castle, Wood No. 10673.

Easily recognized by the trifoliate pubescent leaves, and the long peduncle laxly 2–6-flowered towards the apex, and large stipules.

+ + *Terminal leaflets smaller, 2.5–3.0 cm. long, 28–40 mm. broad.*

21. *R. Jacottetii*, Schinz. in Herb. Zurich.

Stem herbaceous, pubescent or pilose. *Stipules* conspicuous, oblong or ovate-oblong, hairy, 7–10 mm. long. *Leaves* trifoliate, petiolate; terminal leaflets broadly ovate or suborbicular-ovate, hairy, 2–3 cm. long, 2.5–4.0 cm. broad, generally broader than long; terminal petiolules 4–5 mm. long, lateral leaflets unequalsided, broad, common petioles pilose, 2–3 cm. long. *Peduncles* longer than the leaves, pilose, laxly 3–6-flowered towards the summit. *Bracts* lanceolate, pilose, subpersistent. *Pedicels* pilose. *Calyx* pilose, tube short, segments lanceolate, pilose. *Standard* glabrous, 8–10 mm. long; keel boat-shaped, obtuse, a little shorter than the standard. *Pod* not seen.

CAPE PROVINCE.—Komgha, *R. Schlechter*, Flanagan No. 854; Morugha, *R. Schlechter* No. 6170.

This plant is very closely allied to the preceding and perhaps not specifically distinct. The noticeable features are the trifoliate leaves, the terminal leaflets generally broader than long, the long peduncles much longer than the leaves, laxly 3–6-flowered near the summit.

Series 9.—*Flowers 10–12 mm. long in very lax raceme. Terminal leaflets suborbicular or broadly ovate, hirsute. Stem rufo-pilose.*

22. *R. ovata*, Wood and Evans in "Journ. Bot." (1897), 350.

Stem suffruticose, suberect, densely rufo-pilose. *Stipules* linear lanceolate, $\pm$ 6 mm. long. *Leaves* long petiolate, trifoliate; leaflets suborbicular or ovate, entire, base rounded, hispid, and with long hairs ciliate, about 3 cm. long, 2.5 cm. broad; lateral shortly petiolulate. *Racemes* axillary and terminal longer than the leaves, bearing a few distant flowers. *Calyx* hairy, about 7 mm. long. *Standard* obovate, 10 mm. long, base auriculate and clawed, glabrous; carina boat-shaped, apex obtuse. *Pod* oblong, acute, base narrowed, about 18 mm. long.

ZULULAND.—Near Entumeni, Wood No. 3987 and Wood No. 8534.

NATAL.—Lower Tugela, Wood No. 9275.

Easily recognized by the flexuose pilose stems, the suborbicular or broadly ovate leaflets, and the very lax inflorescence of yellow flowers.

Series 10.—*Flowers small, 9–10 mm. long, in var. grandiflora larger in a lax raceme. Leaflets ovate, 2–3 cm. long, 15–25 mm. broad, in typical form. Standard glabrous.*

23. *R. Harmsiana*, Schlechter, ex A. Zahlb. in "Ann. Mus. Nat. Hofmus Wien." XX (1905), 26.

Stem voluble, branches filiform, puberulous. *Stipules* lanceolate, subulate, acuminate. *Terminal leaflets* 2–3 cm. long, 15–25 cm. broad; common petiole pubescent. *Racemes* axillary, laxly several-flowered. *Calyx* campanulate, $\pm$ 8 mm. long; segments subulate

longer than the tube. *Standard* about 1 cm. long, lamina obovate, base auriculate; wings obliquely oblong; carina boat-shaped, obtuse. *Ovary* pilose, style filiform, glabrous.

EAST GRIQUALAND.—Between Mt. Insiswa and the River Umzinhlawa, *Krook* No. 2584.

NATAL.—*Rudatis* No. 1617; near Boston, *Wood* No. 11824.

Var. **Burchellii**, Burt-Davy in "Kew Bull.," 1921, 193.

Slender voluble herb. Leaves trifoliate, terminal somewhat rhomboid, 13–20 mm. long, 9–15 mm. broad, below sparsely glandular. *Peduncles* longer than the leaves, 5–7.5 cm. long, generally laxly 5–7-flowered. *Calyx* glandular, upper lobes short, lower longer, subulate, acuminate. *Standard* glabrous, with purple-brown lines. *Pod* strigose, reaching 2 cm. long.

CAPE PROVINCE.—Near Komgha, *Flanagan* No. 796; Van Stadensberg, *Burchell* No. 4746; Kentani, *Pegler* No. 2032; Humansdorp Div., Kronme River, *Burchell* No. 4860.

EAST GRIQUALAND.—Near Clydesdale, *Tyson* No. 2073.

NATAL.—Umzumbi, *Wood* No. 3143.

Var. nov. **grandiflora**.

Folia trifoliolata, foliolis terminalibus rhombico-triangularibus apice acutis 30–45 mm. longis, 25–55 mm. latis. *Racemi* multiflori cum pedunculo, 8–10 cm. longi. *Flores* lutei, 12–15 mm. longi, longiuscule pedicellati, pedicellis 3–4 mm. longis. *Calyx* in toto 9–10 mm. longus, hirtus et glandulosus, dentibus superioribus 3–5 mm. longis, inferioribus 6–7 mm. longis. *Vexillum* extus glabrum.

NATAL.—Lundies Hill, *Wood* No. 11401, Natal Herb. No. 12427.

The leaflets are thin and covered with scattered hairs. The two uppermost teeth of the calyx are more or less connate, the lowest segment very narrow. It is allied to *R. flavissima* Hochst., but the carina is shorter and the calyx is longer. The wing petals are oblong with a short spur on one side.

Series 11.—*Flowers*, 10–13 mm. *Leaves* trifoliate or unifoliate, leaflets rigid, elliptic-oblong or subrhomboid. *Peduncles* generally long, laxly 3–8-flowered. *Standard* glabrous.

24. **R. nervosa**, Benth., ex Harvey, l.c. 253.

Stem prostrate or twining hispid-pubescent. *Stipules* small, narrow lanceolate or subulate. *Leaves* in typical form trifoliate, shortly petiolate, terminal leaflets elliptic-oblong or subrhomboid, rigid, 3-nerved at base, hispid-pubescent 30–60 mm. long, 15–35 mm. broad, common petioles hispid-pubescent or villous, 3–15 mm. long or sometimes longer. *Peduncles* generally long, laxly 3–8-flowered. *Calyx* lobes lanceolate acuminate, longer than the tube, shorter than the standard. *Standard* glabrous, 8–12 mm. long. *Pod* sessile, villous.

CAPE.—Sand River and Langspruit, *Zeyher*; Winterberg, *Ecklon and Zeyher*.

NATAL.—Near Intshanga, *Wood* Nos. 6231 and 6545; near Gillits, *Wood* No. 7990; Claremont, *Wood* No. 7994.

EAST GRIQUALAND.—Clydesdale, *Tyson* No. 1056.

TRANSVAAL.—Lydenberg, *Wilms* No. 381; Hoogeveld, *Wilms* No. 385; Pretoria, *A. O. D. Mogy* No. 15636; Vereeniging, *Rogers* No. 11904; Kaalfontein, *Pole-Evans* No. H.16796; Jeppesdorp Ridge, Johannesburg, *Gilfillan* No. 54; Barberton, *E. Galpin*, No. 1003.

Forma **unifoliolata**, Bak. fil.

Caules teretes, flexuosi. *Folia* saepissime unifoliolata, late ovata, petiolo brevi suffulta. *Racemi* laxi, 2–5-flori, foliis longiores. *Flores* + 1 cm. longi. *Vexillum* glabrum.

NYLSTROOM.—Mrs. De Jongh No. 6521.

R. nervosa, Benth., differs from *R. hirsuta*, Ecklon and Zeyher, in its shorter petioles, its glabrous standard, and narrower stipules. The petals are yellow, with reddish striations. *R. nervosa*, Benth., var. *pauciflora*, Harvey, is *R. confusa*, Burt-Davy. Zeyher No. 515, referred here doubtfully by Harvey, is *R. venulosus*, K. Schum.

Series 12.—Flowers, 12–18 mm. Leaves trifoliolate, terminal leaflets ovate-rhomboid, acute. Racemes longer than the leaves, laxly several-flowered.

25. **R. Caribaea**, D.C., "Prod.," II, 384; Baker in Oliver "Fl. Trop. Afr.," II.

E. gibba, E. Mey., "Linnaea," VII, 170; *R. acuminata*, Ecklon and Zeyher, 1666; *Copisma gibbum* E. Mey., "Comm.," 137.

Stems elongate, voluble or prostrate, generally pubescent. Stipules minute. Leaves trifoliolate, petiolate; terminal leaflet ovate-rhomboid, acute or sometimes obtuse, membranous, generally 2–4 cm. long and 2–3 cm. broad; lateral leaflets unequally sided; common petiole 1–2 cm. long. Racemes longer than the leaves, laxly several-flowered, pedicels 2–4 mm. long. Calyx upper segments short and broad, lowest conspicuously longer, subulate. Standard glabrous, broadly ovate; wings oblong, base unguiculate; keel boat-shaped, base unguiculate. Pod falcate, much longer than the calyx, 24–30 mm. long.

CAPE.—Districts Uitenhage, Ecklon No. 1667, Alexander No. 117; Somerset Div., Burchell No. 3344, Bowker, Burchell Nos. 3096 and 2997; East London, E. Galpin No. 5684; Schoornstrom and Doornkop, Burke; Mossel Bay, Rogers No. 13348.

NATAL.—Gerrard No. 1902; Weenen, Wood No. 4470; Karkloof, Wood No. 7209; near Durban, Wood No. 7576.

BRITISH KAFFRARIA.—Mrs. Hutton.

DELAGOA BAY.—J. Monteiro No. 14.

TRANSVAAL.—Wakkerstroom, Beeton No. 12526.

Var. *pictam* (*E. gibba*, E. Mey., var. *pictum*, Benth.).

Copisma pictum, E. Mey. "Comm.," 135.

A more densely pubescent or tomentulose plant.

CAPE.—Somerset, Bowker; Howisons Poort, Mrs. Hutton; East London, E. Galpin No. 5685; Port Alfred, Burchell No. 4094; Komgha, Flanagan No. 719.

NATAL.—Inanda, Wood No. 692; Dist. Alexandra, Rudatis No. 896.

SWAZILAND.—Miss Stewart No. 4. ZULULAND.—Ngoye, Wood No. 10333.

Var. nov. **Thorncroftii**.

Leaves trifoliolate, terminal leaflets rhomboid-ovate, subacuminate. Flowers 15–18 mm. long, larger than type.

TRANSVAAL.—Saddleback Mts., Barberton, alt. 5000 ft., G. Thorncroft No. 849; Barberton, E. Galpin No. 1308, is allied to var. *picta* (Benth.).

Series 13.—Flowers small, 9–11 mm. long, in 3–8-flowered racemes. Terminal leaflets triangular-ovate, apex acute.

26. **R. Dieterlenae**, Bak. fil., sp. nov.

Caulis volubilis, gracilis, pubescens. Stipulae angustae, 4–6 mm. longae. Folia trifoliolata, foliolis terminalibus triangulari-ovatis, apice acutis, 3–4 longis, 2·5–3·0 cm. latis. lateralibus inaequilateralibus, petiolo 20–25 longo suffulta. Flores parvisculi in racemos pauci vel plurifloros dispositi, 9–11 mm. longi. Racemi 3–7 cm. longi, saepius 3–8-flori. Calyx in toto 5 mm. longus, dentibus superioribus breviusculis. Vexillum 9–10 mm. longum, extus glabrum. Carina 9–10 longa. Ovarium pilis vestitum. Legumen ignotum.

BASUTOLAND.—Leribe, Mrs. Dieterlen No. 840, Natal Herb. No. 15418.

An ally of *R. minima*, D.C. The racemes are 3-8-flowered, the stipules narrow, and the flowers 9-11 mm. long. The terminal leaflets are longer and larger than in *R. minima*, D.C., and the carina is tipped with blue. The standard is glabrous externally. The young inflorescence is very hirsute.

Series 14.—Flowers, 10-12 mm. Leaflets broadly ovate-orbicular, rigid, hispid-pubescent. Peduncles laxly 3-12-flowered. Standard pubescent.

27. *R. hirsuta*, Ecklon and Zeyher, "Enum.," 256; Harvey, l.c. 252.

Copisma diversifolium, E. Mey., "Comm.," 135.

Stem widely spreading, flexuous, hirsute. *Stipules* ovate, acute. *Leaves* trifoliolate, petiolate; terminal leaflets broadly ovate-orbicular or rhombico-ovate, rigid hispid-pubescent, generally 2-5 cm. long, 2.5-4 cm. broad; petioles 1-2 cm. long, hirsute. *Peduncles* elongate, laxly and distantly 3-8 or sometimes more flowered. *Flowers* 10-12 mm. long. *Standard* densely pubescent, about 10 mm. long; carina boat-shaped, 8-10 mm. long. *Pod* sessile, villous, 18-24 mm. long.

CAPE.—Philipstown, Ecklon and Zeyher No. 1676; Queenstown, E. Galpin No. 1587.

EAST GRIQUALAND.—Near Clydesdale, Tyson Nos. 1448 and 2193.

BASUTOLAND.—Leribe, Mrs. Dieterlen No. 6841.

TRANSVAAL.—Magaliesberg, Burke and Zeyher.

Var. *angustifolia*, Harvey, l.c. 252.

Upper leaves oblong, lower ovate.

NATAL.—Port Natal, Gueinzius.

Series 15.—Flowers, 13-15 mm. Leaves trifoliolate, terminal leaflets ovate, thickish, canotomentose below. Peduncles generally longer than the leaves, laxly few-flowered.

28. *R. argentea*, Harvey, l.c. ii, 253.

Glycine argentea, Thunb., "Fl. Cap.," 292.

Stem voluble, tomentose. *Stipules* ovate, acute. *Leaves* trifoliolate, petiolate; terminal leaflets ovate, thickish, canotomentose below, 15-33 mm. long, 13-22 mm. broad; terminal petiolules 3-6 mm. long; common petioles 10-16 mm. long. *Peduncles* generally longer than the leaves, occasionally shorter, laxly few-flowered. *Pedicels* rather long, 4-6 mm. *Calyx* pubescent, its upper segments broadly lanceolate, lowest rather longer subulate. *Standard* pubescent. *Pod* not seen.

CAPE.—Thunberg, Burchell No. 5122; Knysna, Newdigate No. 207; Stockenström, W. C. Scully No. 4886.

There is a specimen of *Glycine argentea* in Thunberg's Herbarium which agrees with that of Burchell.

This species is easily recognized by the broadly ovate leaflets canotomentose below. The flowers are rather large, 13-15 mm. long.

Series 16.—Flowers small, 6-8 or 9 mm. long. Terminal leaflets rhombico-ovate or rhombico-ovate.

29. *R. minima*, D.C., "Prod.," II, 385; Harvey, l.c. ii, 254.

Dolichos minimus, "Linn.," sp. pl. 726; *Copisma tenue* and *Copisma falcatum*, E. Mey., "Comm.," 136.

Stem slender, voluble, thinly downy or glabrescent. *Stipules* small, narrow. *Leaves* trifoliolate, petiolate; terminal leaflets rhombico-ovate or rhombico-ovate, varying much in size and shape; lateral leaflets unequalsided; common petiole 5-25 mm. long. *Racemes* laxly several or many flowered, longer than the leaves. *Flowers* small, 6-9 mm. long. *Calyx* 3-4 mm. long, lowest lobe longer than the others, much shorter than the petals. *Standard* orbicular-ovate, 5-7 mm. long. *Carina* boat-shaped. *Pods* falcate, 10-17 mm. long, about 4 mm. wide, puberulent, seeds brown.

NATAL.—Durban, *Wood* No. 7556; Port Natal, *Dr. Grant*; near Durban, *Wood* No. 6383.

TRANSVAAL.—Barberton, *E. Galpin* No. 1196; Shilouvane, *Junod* No. 1114; Potgietersrust, *Leendertz* No. 2328, *H. Bolus* No. 11120; Komatipoort, *R. Schlechter* No. 11773.

LOURENCO MARQUES.—*R. Schlechter* No. 11566.

A widely spread little plant. May be recognized by the lax raceme of small flowers, 6–8 or sometimes 9 mm. long.

Series 17.—*Flowers* 10–12 mm. long in lax 10–24-flowered racemes. *Terminal leaflets* ovate-triangular, acute. *Stem* rufous hirsute.

30. *R. stenodon*, sp. nov.

Caulis volubilis rufo-hirsutus. *Stipulae* lanceolatae, acuminatae, rufo-hirsutae, 5–6 mm. longae. *Folia* trifoliolatae, foliolis terminalibus ovate-triangularibus $\pm$ 5 cm. longis, et 4–4.5 cm. latis, lateralibus inaequilateralibus, omnibus utrinque pubescentibus, subtus nerviis rufo-hirtis, petiolo communi rufo-hirsuto 25–30 mm. longo praedita. *Racemi* saepius 10–24-flori, laxi. *Flores* flavi 10–12 mm. longi. *Calyx* in toto, 8–9 mm. longus, glandulosus et hirsutus, dentibus infimis angustissimis et longioribus, dentibus supremis connatis. *Vexillum* obovatum, unguiculatum. *Alae* oblongae uno latere calcaratae. *Carina* naviculariformis. *Ovarium* pilis vestitans.

NATAL.—Oliver's Hoek Pass, *Wood* No. 3344. Alt. 4800 ft. In flower Jan., 1886.

An ally of *R. Pentheri*, Schlechter. The flowers are yellow, the calyx is hirsute and glandular, the lowest segment conspicuously longer than the rest and very narrow. The wing petals have a prominent tooth on one side.

Series 18.—*Flowers*, 10–12 mm. *Leaves* trifoliolate, terminal leaflets rhomboid-ovate. *Racemes* lax, many flowered.

31. *R. memnonia*, D.C., var. *prostrata*, Harvey, l.c. 253.

Stem flexuous, longitudinally striate, pubescent. *Stipules* lanceolate deciduous. *Leaves* trifoliolate, petiolate, terminal leaflets rhomboid-ovate generally, 10–12 mm. long and about the same breadth, terminal petiolules pubescent, 3–4 mm. long; lateral leaflets unequaled; common petiole 5–10 mm. long. *Racemes* longer than the leaves, lax, many-flowered. *Flowers* pendent. *Calyx* glandular, about 10 mm. long; segments longer than the tube, lanceolate. *Standard* pubescent on back, 10–11 mm. long; carina about 10 mm. long, boat-shaped. *Pod* 15–20 mm. long, pubescent.

TRANSVAAL.—Blomberg, *R. Schlechter* No. 4647; Magaliesberg, *Burke* No. 101; Shilouvane, *Junod* No. 768; Komatipoort, *R. Schlechter* No. 11750.

DELAGOA BAY.—*Bolus* No. 1143.

NATAL.—*Gerrard* No. 1949.

The plant described by Harvey as typical *R. memnonia* I have placed under *R. albissima*, Gandoger. It is in all parts tomentose-canescens or silky.

Series 19.—*Flowers*, 10–12 mm. *Leaflets* densely velvety, thick. *Peduncles* elongate, laxly several-flowered. *Standard* pubescent.

32. *R. crassifolia*, Benth. in Harvey, l.c.

R. longipes, Harms in Engler's "Jahrb.," XXVI, 306.

Stem procumbent. *Branches, petioles*, etc., densely velvety. *Stipules* oblong obovate. *Leaves* trifoliolate, petiolate; terminal leaflets elliptic-ovate, subobtusely, softly velvety, 50–55 mm. long, 30–42 mm. broad; petioles 10–25 mm. long, hairy. *Peduncles* elongate, laxly several-flowered or sometimes many-flowered. *Calyx* silky, villous, 6–7 mm. long; segments longer than the tube, shorter than the standard. *Standard* pubescent, 8–10 mm. long; carina boat-shaped. *Pod* sessile, villous, 20–25 mm. long, 7–8 mm. broad.

TRANSVAAL.—Magabiesberg, *Burke and Zeyher*; between Kaap and Crocodile Rivers, *E. Galpin* No. 1067; Lydenberg, *Wilms* No. 395; near Pretoria, *McLea* No. 3153; Shanskop Fort, *Pole-Evans* No. 186; Cerro de Pasco, near Barberton, *E. Galpin* No. 1067.

Easily recognized by the thick velvety leaves, the elongate lax-flowered peduncle, the hairy standard, and the villous pod.

Series 20.—*Flowers 10–12 mm. long in lax racemes much longer than the leaves. Terminal leaflets suborbicular, obtuse. Standard glabrous.*

33. **R. Pantheri**, Schlechter, *ex* A. Zahlb. in "Ann. Nat. Hofmus. Wien.," XX (1905), 26.

Decumbent branches filiform, elongate, voluble, subvillose, puberulous. *Stipules* minute, obliquely lanceolate, acuminate. *Leaves* trifoliolate; terminal leaflets rhomboid-orbicular or orbicular, obtuse, 15–25 mm. long, 12–25 mm. broad; common petiole about 5 mm. long in type, villose-puberulous. *Racemes* elongate, much exceeding the leaves, laxly several or many flowered, longly peduncled. *Calyx* campanulate, about 7 mm. long, somewhat glandular; segments subulate. *Standard* obovoid, base auriculate, about 11 mm. long; wings oblong; carina boat-shaped, obtuse. *Ovary* pilose; style filiform, glabrous. *Pod* not seen.

CAPE.—Albany Division, *Dr. Atherstone*.

EAST GRIQUALAND.—Near Newmarket, *Krook* No. 2631; near Currie's Post, *Schlechter* No. 6087.

NATAL.—Van Reenens Pass, *Krook* No. 2523; near Charlestown, *Wood* No. 5707, *Gerrard* No. 1736.

BASUTOLAND.—Leribe, *Dieterlen* No. 112. KAFFRARIA, *Baur*.

Var. **Hutchinsoniana**, Burt-Davy and Bak. fil. in "Kew Bull.," 1921, 193.

Prostrate herb. Stems hirsute. *Leaves* sessile or subsessile; terminal leaflets suborbicular, obtuse, 30–34 mm. long, 30–33 mm. broad. *Racemes* reaching 15–21 cm. long, laxly-flowered. *Flowers* about 1 cm. long. *Calyx* hirsute, more or less glandular; lobes acuminate. *Corolla* apex purplish; standard yellow, with purple lines. *Pod* not seen.

NATAL.—Karkloof, *J. Wylie* in Herb. *Wood* No. 7212; Edendale, *Wood* No. 4601; Van Reenen, *Wood* No. 6674; near Northdene, *Wood* No. 3180.

EAST GRIQUALAND.—Near Clydesdale, *Tyson* No. 2068.

TRANSVAAL.—Johannesburg, *Gilfillan* No. 211.

Series 21.—*Leaflets ovate-orbicular or subrhomboid, below densely gland-dotted. Flowers 10–12 mm. long, in racemes longer than the leaves. In var. Cooperi, Harvey, flowers larger.*

34. **R. adenodes**, Ecklon and Zeyher, "Enum.," 254; Harvey, l.c. ii, 254.

R. amatymbica, Ecklon and Zeyher No. 1671; *Copisma effusum*, E. Mey., "Comm.," 135.

Prostrate, subglabrous or minutely puberulous. *Stipules* lanceolate. *Leaves* trifoliolate, petiolate; terminal leaflets ovate-orbicular or subrhomboid, acute or obtuse, resinous-dotted, generally 7–25 mm. long, 10–25 mm. broad; terminal petiolule 2–3 mm. long; lateral leaflets unequalsided; common petiole 10–24 mm. long. *Peduncles* longer than the leaves, shortly and closely racemose near the summit. *Calyx* pubescent, its segments lanceolate, longer than the tube. *Standard* glabrous or puberulent, glandular, longer than the calyx. *Pod* hairy, 15–18 mm. long.

CAPE.—Somerset East Division, Boschberg, *Bolus* No. 322; East London, *E. Galpin* No. 7338; Aliwal North, *Bolus* No. 10472; Queenstown Dist., *Mrs. Barber*, *E. Galpin* No. 1595; Komgha, *Flanagan* No. 764; near Philipstown, *Ecklon* No. 1670.

ZULULAND.—Mckenzie. TEMBUKILAND.—*Ecklon* No. 1671.

NATAL.—Krauss No. 301; Drakensberg, *Rehmann* No. 7067; Inanda, *Wood* Nos. 244, 419, 297; between Greytown and Newcastle, *Wilms* No. 1950.

EAST GRIQUALAND.—Near Clydesdale, *Tyson* No. 1447-1055; Umzumwubu Flats, *Tyson* No. 1132.

KAFFRARIA.—*T. Cooper* No. 155; *Baur* Nos. 24, 125, 761.

TRANSVAAL.—Pretoria, *Rehmann* No. 4648; Lydenberg, *Wilms* Nos. 281, 282; Fairy Glen, *Leendert* No. 318; Wakkerstroom, *Beeton* No. 12393; Heidelberg, *Gilfillan* No. 76; Vereeniging, *Gilfillan* No. 142; Standerton, *Burt-Davy* No. 2323.

Var. **laxiflora**, *Burt-Davy* in *Herb.*, Kew.

Pedunculi elongati, laxiflori. *Flores* parviusculi, 7-8 mm. longi.

NATAL.—Pietermaritzburg, *Dr. Sutherland*, *Herb.*, Kew.

The leaflets are rhombic-ovate pointed, 10-17 mm. long, 9-13 mm. broad.

Var. **Cooperi** (*R. Cooperi*, Harvey, ms.)

Folioli majora late ovata vel suborbicularia. *Pedunculi* longi. *Flores* majores, 14-18 mm. longi.

EAST GRIQUALAND.—Near Fort Donald, *Tyson* No. 1637; Clydesdale, *Tyson* No. 2065.

ORANGE FREE STATE.—*T. Cooper* No. 1012.

NATAL.—Gillit's, *Wood* No. 11503; Lynedoch, *Wood* No. 4538; Nottingham Road, *Wood* No. 6042. KREILS COUNTRY.—*Bowker*.

This is a climbing plant with orange-coloured flowers.

Series 22.—*Flowers*, 12-15 mm. *Leaves* trifoliolate, leaflets large, softly villous. *Peduncles* elongate, densely racemose beyond the middle.

35. **R. sigmodes**, Benth., ex Harvey, l.c. ii. 251.

Sigmodostyles villosa, Benth. in Hook, "Lond. Journ.," II, 93.

Stem robust, prostrate, softly villous. *Stipules* broadly triangular. *Leaves* trifoliolate, subsessile or shortly petiolate; terminal leaflets cordate-ovate or suborbicular, rugose-thick, velvety, 7-11 cm. long, 6-12 cm. broad. *Peduncles* elongate, densely racemose beyond the middle. *Calyx* 7-9 mm. long, villous, its segments longer than the tube. *Standard* glabrous, 12-13 mm. long; keel 13-14 mm. long. *Pod* falcate, hairy, 20-28 mm. long, sessile, obtuse.

NATAL.—Near Pietermaritzburg, *Krauss*; near Howick, *Wood* Nos. 8384 and 8665; Inanda, *Wood* No. 1059; near Pinetown, *Wood* No. 3406.

EAST GRIQUALAND.—Zuurberg, *Tyson* No. 1581.

TRANSVAAL.—*Wilms* No. 387 (forma).

Wilms No. 387, issued with a ms. name, seems a form of this species, with more angled leaves than usual.

Series 23.—*Leaflets* glandular. *Racemes* dense, many-flowered. *Flowers* about 15 mm. long. *Standard* glandular.

26. **R. chrysanthia**, Schlechter, ex A. Zahlb. in "Ann. Nat. Hofmus. Wien.," XX (1905), 27.

Decumbent at length, vobule; branches filiform, more or less compressed, angular, puberulous. *Stipules* lanceolate, acuminate. *Leaves* trifoliolate, petiolate; terminal leaflets ovate-suborbicular, 18-30 mm. long, 15-30 mm. broad, below glandular; common petiole 20-35 mm. long. *Racemes* longly peduncled, densely many-flowered (15-25), much exceeding the leaves. *Calyx* campanulate, about 10 mm. long, glandular; segments linear-lanceolate. *Standard* obovoid, about 13 mm. long; wings obliquely subfalcate-spathulate; carina boat-shaped. *Ovary* pilose; style filiform, glabrous. *Pod* not seen.

NATAL.—Ixopo, *Krook* No. 2547, *Schlechter* No. 6641; District Alexandria, *Wilms* No. 1305.

Easily recognized by its dense raceme, glandular leaves, and glandular standard.

D.—STEM VOLUBLE. LEAVES NARROWER.

Series 24.—*Leaves trifoliolate, leaflets elliptic-oblong or oblong. Peduncles 1-2-flowered, shorter than leaves.*

37. *R. puberula*, Harvey, l.c. 255.

Copisma tottum, E. Mey., "Comm.," 133; *Eriosema puberulum*, Ecklon and Zeyher, 1677; *Hedysarum ciliatum*, Thunbg. "Fl. Cap.," 594; *Desmodium? ciliatum*, D.C., "Prod.," II, 333.

Root woody. Stems several, spreading, prostrate or subvoluble. *Stipules* lanceolate. *Leaves* trifoliolate, very shortly petiolate; leaflets elliptic-oblong or oblong-rigid; terminal leaflets 24-30 mm. long, 8-16 mm. broad. *Peduncles* 1-2-flowered, shorter than the leaves. *Flowers* 10-13 mm. long. *Calyx* villous, 6-7 mm. long, its segments longer than the tube, shorter than the petals. *Standard* glabrous, longer than the calyx, 10-12 mm. long; carina shorter than the standard. *Pod* sessile, very villous, 12-16 mm. long seeds.

CAPE.—Galgebosch, *Thunberg*; Albany Division, *Bowie*, *Burchell* Nos. 3498 and 4189; Visch River, *Hutton*; near Grahamstown, *P. MacOwan* No. 473; Mossel Bay Division, *Burchell* No. 6210; Uitenhage, *Ecklon* No. 1577; Riversdale Division, near Spiegel River, *Burchell* No. 7206; Kingwilliamstown Division, *Tyson* No. 853. NATAL.—Camperdown, *Miss Franks* No. 12528. BRITISH KAFFRARIA. —*T. Cooper* No. 539. TRANSVAAL.—Magaliesberg, *Burke*.

Easily recognized by the several spreading subvoluble stems, the trifoliolate leaves with rigid elliptic-oblong or oblong leaflets, and the few-flowered inflorescence shorter than the leaves.

Series 25.—*Leaves usually trifoliolate, rarely unifoliolate; leaflets ovate, or oblong or lanceolate, petiolate. Peduncles 1-3-flowered.*

38. *R. Totta*, D.C., "Prod.," II, 388; Harvey, l.c. ii, 255.

Glycine Totta, Thunberg, "Flora Cap.," 591.

Procumbent or subvoluble. *Stem* filiform, hispid-pubescent or glabrate. *Stipules* small, ovate-lanceolate, rigid. *Leaves* in typical form, trifoliolate, petiolate; leaflets in typical form, ovate or ovate-oblong, reticulate, terminal leaflets 10-25 mm. long, 5-12 mm. broad; common petioles 5-15 mm. long; lateral leaflets somewhat unequelsided. *Peduncles* filiform, equalling the petioles or longer, 1-3-flowered. *Calyx segments* longer than the tube, but shorter than the standard. *Standard* glabrous, 6-9 mm. long. *Pod* oblong, sessile, pilose, pubescent.

CAPE.—Galgebosch, *Thunberg*, type; Grahamstown, *MacOwan* No. 462; Humansdorp Division, *Burchell* No. 4824, *Burt-Davy* No. 12088.

BECHUANALAND.—*Burchell* Nos. 2433, 2487, 2569.

NATAL.—Ladysmith, *Wood* No. 8254; Phoenix, *Schlechter* No. 3027.

The type of *R. Totta* in Thunberg's Herbarium has ovate or ovate-oblong leaves. *Copisma paniculatum*, E. Mey., from Magaliesberg in the Transvaal, and Katrivierspoort, has longer and narrower leaflets, but is not specifically distinct.

Var. *Fenchelii*, Schinz in "Vierteljahrsschrift Nat. Ges. Zurich," LII, 426 (1907).

Stem grey puberulous. *Leaves* trifoliolate.

GREAT NAMAQUALAND.—Keetmanshoop, *Fenchel* No. 195.

Var. *namaensis*, Schinz. in Herb.

Leaves trifoliolate, leaflets lanceolate or ovate-lanceolate, grey puberulous.

GREAT NAMAQUALAND. Seheini, alt. 4000 ft., *Pearson* No. 3741; Keetmanshoop, *Fenchel*. Intermediate between *R. Totta*, D.C., and *R. longiflora*, Schinz.

Var. **pilosa**, Harvey, l.c. ii, 256, pro species.

Copisma pilosum, E. Mey., "Comm.," 133.

Stem patently pilose. *Leaves* trifoliate, leaflets of lower leaves ovate-oblong of the upper linear.

PORT NATAL—*Gueinzus*; Zureberg and between the Omtendo and Omsamculo, Drège.

This is hardly specifically distinct from *R. Totta*, D.C.

Var. **unifoliolata**, Burt-Davy, in "Kew Bull.," 1921, 49, pro species.

Leaves unifoliate or trifoliate, leaflets ovate, common petiole short, 1–2 mm. *Flowers* 7–8 mm. long.

TRANSVAAL.—Ermelo, *Burt-Davy* No. 17403. HOOGEVELD.—*Rehmann* No. 6836.

Var. **brevipetiolata**, Bak. fil.

Caules flexuosi, pilosi vel tomentosi. *Stipulae* lanceolatae, striatae. *Folia* trifoliolata, anguste oblonga vel elliptica, foliolis terminalibus 15–23 mm. longis, 5–12 mm. latis, apice acutis, petiolo communi brevi, 2–4 mm. longo suffulta. *Flores* 8 mm. longi, pedunculati, saepissime solitarii rarius bini. *Calyx* 6–7 mm. longus, pilosus. *Legumen* 10–15 mm. longum, pilosum.

TRANSVAAL.—Johannesburg, Jeppestown Ridge. *Gilfillan* Nos. 22 and 65; Nat. Herb. Pretoria. Nos. 6103 and 6104.

Ally of *R. Totta*, D.C., var. *pilosa*, Harvey. The noticeable features are the pilose or tomentose stems, the short petioles. It was distributed as *R. puberula*, Harvey.

Var. **graciliflora** (*R. graciliflora* Harms) in Herb. Mus. Brit.

Stems flexuous, filiform, pubescent or nearly glabrous. *Stipules* lanceolate, longitudinally striate, 4–5 mm. long. *Leaves* trifoliate, petiolate; terminal leaflets linear-lanceolate, glabrous above, with a few scattered hairs below, 3–4.5 cm. long, 3–5 mm. broad; terminal petiolules 4 mm. long; lateral leaflets slightly unequelsided at base; common petioles 15–25 mm. long. *Peduncles* slender, generally one-flowered. *Calyx* hairy; segments lanceolate, longer than the tube, but shorter than the corolla. *Standard* glabrous, 7–9 mm. long. *Pod* about 16 mm. long, hairy outside, seeds reddish brown.

CAPE.—Queenstown, *E. Galpin* No. 1547. NATAL.—Mooi River, *J. Webb* No. 7011. EAST GRIQUALAND.—Kokstad, *Tyson* No. 1336. BASUTOLAND.—Leribe, *Mrs. Dieterlen* No. 6839. ORANGE FREE STATE.—Bethlehem, *T. Potgieter* No. 59.

TRANSVAAL.—Lydenberg, *Wilms* No. 367; Modderfontein, *Conrath* No. 259; near Pretoria, *Bolus* No. 7750; Ermelo, *G. McCall* No. 7592; Davel, *W. J. Wayland*, sine no.; near Pretoria, *Pole-Evans* No. 51.

A plant from Camperdown, *Franks* (Com. *Medley Wood* No. 11826), is a form of this species.

Differs from the type of *R. Totta* in Thunberg's Herbarium by the linear lanceolate leaves.

39. **R. longiflora**, Schinz in "Bot. Verh. Brand.," XXX, 168 (1889).

Stems slender, voluble, pilose. *Leaves* trifoliate, petiolate; terminal leaflets narrow, lanceolate, acute, mucronate, 13–30 mm. long, 2.5–5 mm. broad; terminal petiolules 2 mm.; lateral leaflets unequelsided; common petiole reaching 8 mm. long. *Peduncles* axillary, two-flowered. *Calyx* bilabiate; segments as long as or a little longer than the tube. *Standard* glabrous, 22 mm. long; keel about the same length; wings shorter, about 13 mm. long. *Pod* pilose, sessile, about 23 mm. long and 7 mm. broad.

GREAT NAMAQUALAND.—Osis.

Noticeable on account of the narrow lanceolate terminal leaflets and two-flowered racemes.

R. amboense, Schinz., is allied, but the flowers are smaller.

R. Totta, D.C., var. *namaensis*, Schinz., seems intermediate between *R. Totta* and the present species. Flowers longer than in *R. Totta*, D.C.

Series 26.—Leaflets lanceolate or oblong-lanceolate, when young sericeous. Flowers 8–10 mm. long, in a very lax raceme.

40. *R. elegantissima*, Schinz. in "Vierteljahrsschrift Nat. Ges. Zurich." LII, 428 (1907).

R. sericea, Conrath in Herb., Kew.

Suffrutex with flexuous, filiform branches. *Stipules* longitudinally striate. *Leaves* trifoliolate, petiolate; terminal leaflets lanceolate, 2–4 cm. long and 8–12 mm. broad; terminal petiolules 2–3 mm. long; lateral leaflets slightly unequilateral; common petiole 8–15 mm. long. *Racemes* few-flowered, very lax. *Flowers* 8–10 mm. long. *Calyx* segments acuminate, shorter than the standard. *Standard* 7–9 mm. long. *Pod* 15–17 mm. long, pubescent, seeds dark coloured.

TRANSVAAL.—*Rehmann* No. 5545; near Pretoria, *Conrath* No. 258; Onderstepoort, *A. O. D. Mogg* No. 15674.

A suffrutex with flexuous branches. The young foliage is silvery, the leaflets are lanceolate.

Series 27.—Leaflets oblong-ovate or ovate-lanceolate, tomentose. Flowers 10–12 mm. long, in very lax few-flowered racemes. Stems rufo-tomentose.

41. *R. venulosus*, K. Schum. in "Just Jahrb.," XXVII, 496.

Dolicholus venulosus, Hiern. in "Cat. Welw. Afr. Pl.," I, 271; *R. cinnamomea*, Schinz. in "Vierteljahrsschrift Nat. Ges. Zurich." LII, 429 (1907); *R. remota*, Conrath in "Kew Bull." (1908), 224.

Stem terete, densely rufo-tomentose, scandent. *Stipules* small. *Leaves* trifoliolate, oblong, ovate or ovate-lanceolate tomentose; terminal leaflets 20–35 mm. long, 10–14 mm. broad; lateral leaflets base more or less oblique; common petioles 10–15 mm. long. *Racemes* axillary, very lax, 2–5-flowered; pedicels densely pubescent, 2–3 mm. long. *Calyx* 7 mm. long; segments acuminate, longer than the tube. *Standard* 8–11 mm. long. *Pod* oblong, pilose, about 2 cm. long.

TRANSVAAL.—Magaliesberg, *R. Schlechter* No. 4162; Witpoortje, near Johannesburg, *Conrath* No. 262; Mooi River, *Burke*; Rustenburg, *Olive Nation* No. 157.

BECHUANALAND.—Kuruman Hills, *Burchell* No. 2172.

Easily recognized by the terete, scandent, cinnamon-coloured, hirsute or tomentose stem, and very lax 2–5-flowered racemes.

Zeyher No. 515 is this species.

Series 28.—Leaves trifoliolate, leaflets oblong or oblong-lanceolate, or ovate. Peduncles slender, longer than the leaves, 1–6-flowered at the summit. Pod stipitate.

42. *R. glandulosa*, D.C., "Prod.," II, 388; Harvey, l.c. 256.

Glycine glandulosa, Thunberg, "Cap.," 591; *Glycine heterophylla*, Thunberg, "Cap.," 592; *Copisma glandulosum*, E. Mey., "Comm.," 133; *Rhynchosia glandulosa*, riparia, *glabra*, Ecklon and Zeyher; *Glycine glabra*, Spreng; New Emdack, iii, 54. *Rhynchosia glabra*, D.C., "Prod.," II, 387; *Copisma glabrum*, E. Mey., "Comm.," 134.

Prostrate or *volute*, glabrous or downy. *Stipules* ovate or sub lanceolate. *Leaves* trifoliolate, petiolate; leaflets oblong or oblong-lanceolate, or ovate; terminal leaflets 10–30 mm. long, 3–10 mm. broad, resinous-dotted and paler beneath. *Peduncles* slender,

longer than the leaves, 1-6-flowered at the summit. *Calyx* tube shorter than the lanceolate segments. *Standard* glabrous. *Pod* stipulate, glabrous or downy, 16-18 mm. long, tapering at base into a stipe 2-4 mm. long.

CAPE.—Common on stony hills from Capetown to Uitenhage and Albany.

Series 29.—*Leaflets* linear, below resinous-punctate. *Pedicels* axillary one-flowered.

43. **R. rigidula**, D.C., "Prod.," II, 388.

Stem erect, retrorsely cano-pubescent. *Leaflets* linear, mucronate, pubescent, below resinous-punctate. *Pedicels* axillary, one-flowered, at the middle inflexed, then deflexed. the length of the petiole. *Pod* subvillose, one-seeded, oblong.

CAPE.—*Burchell* No. 2587.

This is only known to me from De Candolle's description, of which the above is a translation. I think there is some error in the number, as this number belongs to a plant of another family.

E.—STEMS RIGID OR SUBRIGID. USUALLY ERECT.

Series 30.—*Leaves* trifoliolate, subsessile or shortly petiolate, terminal leaflets elliptic-oblong or lanceolate-oblong. *Peduncles* shorter than the leaves, closely few-flowered in var. *densiflora*, mostly terminal and more densely flowered.

44. **R. Orthodanum**, Benth., ex Harvey, l.c. 257; Wood, "Natal Plants," t. 220.

Orthodanum latifolium, *sordidum*, and *argenteum*, E. Mey., "Comm.," 131-2; *Eriosema sericeum*, Ecklon and Zeyher, "Enum.," 256.

Erect suffruticose, silky-villous or glabrescent. *Stipules* lanceolate. *Leaves* trifoliolate, subsessile, or shortly petiolate; leaflets elliptic-oblong or lanceolate-oblong, or oblanceolate, 20-60 mm. long, 5-20 mm. broad. *Peduncles* shorter than the leaves, closely few-flowered. *Calyx* 8-10 mm. long; lobes lanceolate-acuminate, much longer than the tube, either shorter or longer than the standard. *Standard* glabrous, suborbicular-obovate; keel boat-shaped. *Pod* 12-16 mm. long, silky, sometimes glabrescent.

CAPE.—Winterberg, above Philipstown, Kat River, Ecklon and Zeyher; Kentani, Miss Pegler.

NATAL.—Port Natal, Drège, Krauss No. 374; in open ground all over the Colony, fide Wood.

BRITISH KAFFRARIA.—*T. Cooper* No. 294.

TRANSVAAL.—Near Crocodile River, R. Schlechter No. 3921; Johannesburg, Dr. Rand No. 951.

Var. **Muhlenbeckii**, Harvey, l.c. 257.

Orthodanum Muhlenbeckii and *O. glabratum*, Meisner in Hook, "Lond. Journ.," II, 91-2.

Stems dwarf. *Leaves* and *flowers* smaller, terminal; leaflets generally 10-15 mm. long, 3-7 mm. broad.

CAPE.—Near Kromme River, Uitenhage; Albany Division; Uitenhage, Bowie; Queenstown. E. Galpin No. 1958.

Var. nov. **densiflora**.

Pedunculi saspissine terminale copiose floriferi.

NATAL.—District Alexandra, H. Ridatis No. 1349.

This species is variable in size and shape of leaflets. It occurs as far north as Gazaland, where it was collected near Chipete by C. F. M. Swynnerton.

Series 31.—*Terminal leaflets* broad, cinereous-tomentose, *Racemes* few-flowered. *Flowers* 16-17 mm. long.

45. **R. spectabilis**, Schinz in "Vierteljahrsschrift Nat. Ges. Zurich." LV, 236 (1910).

Fruticose, stems canotomentose. *Stipules* small, hairy. *Leaves* trifoliolate, petiolate; terminal leaflets broadly ovate or suborbicular ovate, base cordate or subcordate, 3-5 cm. long, 3-5.5 cm. broad; terminal petiolules 8-10 mm. long; lateral leaflets unequalsided, all markedly reticulate and canotomentose below. *Racemes* few-flowered, shorter than the leaves. *Calyx* canotomentose externally, about 11 mm. long; lobes lanceolate. *Standard* covered with white hairs externally; keel about as long as the standard. *Pod* 20-25 mm. long, apiculate, hairy.

TRANSVAAL.—Moord Drift, *R. Schlechter* No. 4310, *Leendertz* No. 2228.

A canotomentose shrub with few-flowered racemes, and flowers 16-17 mm. long.

Series 32.—*Leaves* shortly petiolate, terminal leaflet oval or suborbicular. *Raceme* shorter than the leaves, few-flowered. *Stem* terete.

46. **R. Rudolphi**, Harms, in "Notizblatt. Bot. Gard.," Berlin, B.Ld. VIII, 156 (1922).

Species ad *R. ambacensem*, K. Schum (*Dolicholum ambacensem*, Hiern) affinis. *Caulis* rigidus, erectus, ferrugineo-tomentosus. *Stipulae* ferrugineo-tomentosae mox delapsae. *Folia* trifoliolata, foliis terminalibus ovalibus vel ovali-suborbicularibus utrinque tomentosis, 4-4.5 cm. longis, 25-33 mm. latis, foliolis lateralibus saepius minoribus, 3-3.5 cm. longis, 15-30 mm. latis, petiolo communi ferrugineo-tomentoso 5-10 mm. longo suffulta petiolulis foliolorum terminalium 6-8 mm. longis etiam ferrugineo-tomentosis. *Racemi* foliis breviores saepius 2-3-flori, rhachi ferrugineo-tomentosa. *Calycis* tubus brevis extus ferrugineo-tomentosus, lobis angustis lineari-lanceolatis tubo multoties longioribus. *Petala* delapsa. *Ovarium* pilis albidis dense villosum.

TRANSVAAL.—Near Crocodile River, alt. 1600 m., *R. Schlechter* No. 3910, Herb. Zurich

This is an erect plant, the stem being ferrugineo-tomentose. The leaves are trifoliolate, the terminal leaflets are longer than they are broad. The racemes are 2-3-flowered. *Calyx* 13-14 mm. long, of which the tube is only 4 mm. It differs from *R. ambacensis*, K. Schum. by the shape of the terminal leaflets and the tomentum. It is also allied to *R. spectabilis*, Schinz.

Series 33.—*Leaflets* ovate or orbicular-ovate. *Peduncles* axillary, two-flowered. *Flowers* 18-20 mm. long.

47. **R. komatiensis**, Harms in Engler's "Jahrb.," XXVI, 306.

Fruticose erect, branches pubescent. *Leaves* trifoliolate, shortly petiolate; terminal leaflets ovate or orbicular-ovate, 15-27 mm. long, 12-20 mm. broad; lateral leaflets unequalsided; common petiole 5-10 mm. long, tomentose. *Peduncles* axillary, short, about two-flowered; pedicels equalling calyx or shorter. *Calyx* obliquely campanulate; teeth ovate; lowest longest. *Corolla* about 18 mm. long, about three times the length of the calyx; standard pilose externally. *Ovary* and young pod densely hirsute.

TRANSVAAL.—Between Spitzkop and the Komati River, *Wilms* No. 374; *Shilovane*, *Junod* No. 541.

A shrub with ovate or ovate-orbicular leaflets and two-flowered peduncles. The corolla is about three times the length of the calyx, which is an interesting feature of this species.

Section 34.—*Leaves* petiolate, terminal leaflet ovate-lanceolate. *Racemes* shorter than the leaves. *Flowers* large. *Standard* 15-18 mm. long in type. *Upper calycine lobes* shorter than lowest lobe.

48. **R. clivorum**, S. Moore in "Journ. Bot.," XVI, 131 (1878).

Suffruticose erect. *Stipules* rather large, oblong-acuminate. *Leaves* trifoliolate, petiolate; terminal leaflet ovate-lanceolate, below conspicuously nerved, 5-6 cm. long, 18-25 mm. broad; lateral leaflets unequalsided. *Racemes* shorter than the leaves, few-flowered,

Flowers yellow, perfumed. *Calyx lobes* oblong, acute, unequal; lower longer, slightly longer, than the keel; upper somewhat broader; tube about 5 mm. long; lowest lobe 11–12 mm. long. *Standard* 15–18 mm. long, ovate, about as long as the carina; carina boat-shaped, obtuse, base longly unguiculate. *Pod* about 15 mm. long, seeds subreniform.

TRANSVAAL.—Near Pilgrims Rest, *J. McLea* No. 3033; Ivy Range, alt. 5000 ft., *Thorncroft* No. 342.

Var. nov. **longicalycina**.

Folia trifoliolata, foliolis ovato-lanceolatis. *Calycis lobis* inferioribus, 20–22 mm. longisquam reliquis et carinam longioribus, calyx in toto 26–28 mm. longus.

TRANSVAAL.—Pietersburg, *Isobel MacCallum* No. 1415, Herb. Kew.

This variety differs in having a longer calyx and rather longer leaflets.

This species is a small shrub and is easily recognized by the erect habit, large flowers, and ovate-lanceolate leaflets.

R. pycnantha, Harms, is doubtfully distinct.

Series 35.—*Silky-silvery plants. Leaves trifoliolate. Raceme few-flowered, generally shorter than leaves, rarely longer. Flowers 16–23 mm. long. Standard pubescent.*

(a) *Leaflets broadly ovate or cordate-ovate.*

49. **R. nitens**, Benth., in Harvey, l.c. 257.

Erect suffruticose, everywhere silky-silvery. *Stem* terete or subterete. *Stipules* obsolete. *Leaves* trifoliolate, petiolate; terminal leaflets broadly ovate or cordate-ovate; lateral leaflets unequalsided; common petiole, 6–12 mm. *Racemes* 2–3-flowered, shorter than the leaf. *Flowers* 16–18 mm. long, yellow. *Calyx* much shorter than the petals, 7–8 mm. long, lobes shorter than the tube. *Standard* pubescent, 16–18 mm. long, about twice as long as the calyx; carina boat-shaped, a little shorter than the standard. *Pod* densely pilose, 3 cm. long.

TRANSVAAL.—Magaliesberg, *Burke and Zeyher*; Waterkloof, Pretoria, *I. C. Verdoorn* No. 40; *Mentjes Kop, Pole-Evans* No. 356.

This is an erect suffrutex with silky-silvery foliage and yellow flowers. The standard is pubescent and about twice as long as the calyx.

(b) *Leaflets longer, oblong-obovate. Flowers larger.*

50. **R. Galpini**, sp. nov.

Suffrutex erectus, caulibus juvenilibus tomentosis. *Folia* trifoliolata, praecique subtus argenteo-sericea, foliolis terminalibus oblongo-obovatis, 4–5 cm. longis, 2–2.5 cm. latis, foliolis lateralibus inaequalateralibus, angustioribus 3.5–4 cm. longis, 12–15 mm. latis, petiole communi 3–6 mm. longo praedita. *Racemi* pluriflori. *Flores* majusculi, 20–23 mm. longi. *Vexillum* extus pilis vestitum calyce distincte longius. *Calyx* extus pubescens, in toto 10 mm. longus, dentibus lanceolatis 5 mm. longis tubo subaequilongis. *Ovarium* pilis flavidis vestitum. *Legumen* ignotum.

TRANSVAAL.—Barberton, mountain sides, *Musidora, E. E. Galpin* No. 442. Differs from *R. nitens* in the much longer and differently shaped leaflets.

Series 36.—*Leaves petiolate, terminal leaflets rhomboid-ovate, softly pubescent below. Flowers pedicellate. Standard 18 mm. long.*

51. **R. Bakeri**, Schinz in "Bull. Herb. Boiss," Ser. II (1908). 628.

Fruticose probably erect, branches pilose. *Stipules* not seen. *Leaves* petiolate trifoliolate, terminal leaflets rhomboid-ovate or suborbicular-ovate, coriaceous apex shortly acuminate, about 6 cm. long and 5 cm. broad; lateral leaflets unequalsided; common petiole 25 mm. long. *Inflorescence* racemose. *Calyx* hirsute, the lowest segment being longer than the rest. *Standard* glandular externally, about 18 mm. long and 12 mm. broad; wings about 3 mm. broad; carina 15 mm. long, blunt and somewhat longer than the wings. *Ovary* and young pods densely hirsute,

TRANSVAAL.—Terminus of Selati Railway, *Junod* No. 1585.

A shrub with trifoliate leaves and rhomboid-ovate leaflets. The calyx is glandular, also the standard.

Series 37.—*Leaves viscidly hairy and resin-dotted, small, oblong, bullate. Peduncles short, 1-2-flowered.*

52. **R. bullata**. Benth., *ex* Harvey and Sonder, l.c. ii, 257.

Hydrosia bullata, E. Mey., "Comm.," 89; *R. uncinata*, Schlechter in Herb. Mus. Brit.

Shrubby erect, branched, viscidly hairy, and resin-dotted. *Stipules* ovate-acuminate, longer than the petiole. *Leaves* trifoliate, on very short petioles; leaflets oblong, small, bullate, with recurved margins; terminal leaflets 5-8 mm. long, 2-3.5 mm. broad. *Peduncles* scarcely longer than the leaves, 1-2-flowered. *Calyx* glandular; lobes scarcely longer than the tube. *Standard* glabrous, 12-15 mm. long, with a longish claw; carina equalling the vexillum. *Pod* 10-12 mm. long, oblong, compressed, sparsely pilose.

CAPE.—Clanwillian Div., Wupperthal, *Drège*, *Bolus* No. 8990; Packhuis Berg, *R. Schlechter* No. 8661.

A very distinct species, being an erect, viscidly hairy little shrub. The peduncles are short and 1-2-flowered.

Series 38.—*Terminal leaflets elliptical. Racemes rather longly peduncled. Stems distinctly angled.*

53. **R. angulosa**, Schinz in "Bull. Herb. Boiss.," Ser. II (1908), 628.

Rootstock woody, branches somewhat flexuous, angled, canopilose. *Stipules* lanceolate. *Leaves* trifoliate, very shortly petiolate; terminal leaflets elliptical hairy on the back, 3-4.5 cm. long, 10-20 mm. broad; lateral leaflets somewhat smaller; common petiole canopilose 2-4 mm. long. *Racemes* axillary, longly peduncled, about as long as leaves or slightly shorter. *Calyx* pilose with long hairs; tube 3 mm. long; the lowest segment is about 9 mm. long, and is longer than the upper segments. *Standard* glabrous, 13-15 mm. long, 10 mm. broad; carina blunt; wings narrow. *Pod* 20-23 mm. long and about 7 mm. broad.

TRANSVAAL.—Mountain sides, Barberton, *E. Galpin* No. 1095, alt. 4500-5000 ft.; Messina, near Barberton, *Galpin* No. 507.

This is a dwarf shrub with angular stems, trifoliate very shortly petiolate leaves, and a raceme of 3-8 orange-coloured flowers, the standard being streaked and reddish.

Series 39. *Terminal leaflet obovate, base cuneate. Racemes longer than the leaves, many-flowered.*

54. **R. Woodii**, Schinz in "Bull. Herb. Boiss." (1894), 197, tab. iii, figs. A-F.

Stem erect, cano-pubescent, longitudinally striate. *Stipules* lanceolate, subacuminate. *Leaves* trifoliate, subsessile or petiole very short, terminal leaflets obovate or oblong-obovate; lateral leaflets unequalsided. *Racemes* longer than the leaves, many-flowered. *Peduncles* canescent-tomentose. *Flowers* yellow, the standard being striate with brown lines. *Calyx* hairy, segments lanceolate, two uppermost connate, 6-7 mm. long, distinctly shorter than the petals. *Standard* obovate glabrous, 10-12 mm. long; wings spurred unguiculate; carina boat-shaped, apex obtuse, base unguiculate. *Pod* 18-20 mm. long, glabrous, pointed, seeds black, reniform, 6 mm. long.

NATAL.—Inanda, *Wood* No. 1884, *Rehmann* No. 8411, *Wood* No. 6627.

An easily recognized shrub with obovate terminal leaflets, racemes longer than the leaves, and yellow flowers, the standard being striate with brown lines.

Series 40.—*Leaves silky-silvery or cinereous pubescent. Flowers 8-12 mm. long, in long racemes longer than the leaves.*

55. *R. albissima*, Gandoger in "Bull. Soc. Bot., France," LX, 461, 1913.

R. memnonia, Harvey pro-parte; Wood, "Natal Plants," t. 349.

Caulis rigidus, cano-tomentosus, longitudinaliter striatus. *Stipulae* lanceolatae, caducae. *Folia* trifoliolata petiolata, foliolis terminalibus rhombeo-obovatis, 2-3 cm. longis, 2-2.5 cm. latis, petiolulis terminalibus 6-8 mm. longis, foliolis lateralibus inaequilateralibus, petiolo communi saepissime, 15-20 mm. longis. *Racemi* axillares, folio longiores, multiflori. *Flores* penduli, 8-12 mm. longi. *Calyx* canescenti-tomentosus, 6-7 mm. longus, segmentis acuminatis. *Vexillum* extus pilis vestitum. *Carina* vexillo subaequilonga. *Legumen* canescenti-tomentosum, 12-15 mm. longum.

TRANSVAAL.—Magaliesberg, *Burke*; near Aapies River, *Schlechter* No. 3619; Gezina, near Pretoria, *A. O. D. Mogg* No. 10082; above Aapies River, *Rehmann* No. 4374; Hartbeestpoort, *Miss Findlay* No. 21.

ZULULAND.—Ntondweni, *J. Wylie* (Wood No. 9197); Umsundusi, *Wood* No. 3954.

Easily recognized by the whole plant being tomentose-canescens; the stems are rigid, the terminal leaflets rhombeo-obovate, the racemes longer than the leaves, many-flowered; the flowers are yellow.

This is not *R. memnonia*, D.C.

Between Lourenco Marques and Komatipoort, *Bolus* No. 7751.

56. *R. Pegleri*, sp. nov.

Caulis strictus, virgatus, pubescens, longitudinaliter striatus. *Stipulae* angustae lineari-lanceolatae. *Folia* trifoliolata, foliolis terminalibus rhombeo-ovatis, 18-22 mm. longis et latis, punctatis foliolis lateralibus inaequilateralibus, petiolo pubescente, 14-25 mm. longo suffulta. *Racemi* densi multiflori, pedunculi folio longiores. *Flores* 10-11 mm. longi. *Calyx* in toto 9-10 mm. longus, pubescens, dentibus infimis, 7 mm. longis. *Vexillum* extus pubescens, in toto 10 mm. longum. *Carina* apice obtusa. *Ovarium* pilis vestitum. *Legumen* ignotum.

KENTANI.—Valleys, *Miss Alice Pegler* No. 156.

A plant allied in some respects to *R. adenodes*, from which it differs in its much thicker virgate stem. The standard is hairy externally. The calyx tube is 3 mm. long, the upper teeth 6 mm. long, the lower 7 mm.

Series 41.—Leaves trifoliolate, leaflets narrow. Flowers solitary on peduncles longer than the leaves.

57. *R. pauciflora*, Bolus in "Trans. S. Afr. Phil. Soc.," XVI, 383 (1906).

Suffrutes with virgate stems. *Stipules* small. *Leaves* trifoliolate, subsessile or shortly petiolate; leaflets rigid, narrow, either linear-oblong or linear, margins distinctly revolute; terminal leaflets 4-6 cm. long, 3-5 mm. broad, midrib below conspicuous. *Peduncles* longer than the leaves, bent just below the flower. *Calyx* lobes lanceolate, distinctly longer than the corolla, total length of calyx 15-16 mm. *Petals* 10-12 mm. long; keel boat-shaped; standard glabrous. *Ovary* hairy. *Pod* 3.5-4 cm. long, apiculate.

SWAZILAND.—*Burtt-Davy* No. 2825. TRANSVAAL.—Spitskop, *R. Pott* No. 5062.

A virgate shrub with trifoliolate leaves and narrow rigid leaflets; the peduncles are one-flowered and longer than the leaves, and the calyx is longer than the corolla.

Series 42.—Leaves trifoliolate. Peduncles longer than leaves, few-flowered. Stem rigid or flexuous.

58. *R. viscidula*, Steud., "Nom.," ed. 2, II, 254; Harvey, l.c. 252.

Copisma viscidulum, E. Mey., "Comm.," 134.

Stem fruticulose, rigid or subvoluble. *Stipules* lanceolate or ovate or subrotund, below glandulose-punctate, 8-11 mm. long, 10-18 mm. broad; common petioles 10-22 mm. long;

lateral leaflets unequalsided. *Racemes* longly peduncled, few-flowered. *Flowers* 10–12 mm. long; pedicels 3 mm. long. *Calyx* lobes longer than tubes. *Carina* boat-shaped, blunt, about 11 mm. long. *Pods* sessile, oblong, viscoso-pubescent.

NAMAQUALAND.—Khamiesberg, W. Pearson; Ceres Division, Koude Bockeveld, R. Schlechter No. 8875.

E. MEYER divides this species into :—

(a) *nemorale* magis volubile, foliolis majoribus tenuioribus subcordatis piloso-pubescentibus.

LITTLE NAMAQUALAND.—Kasparskloof, Drège.

(b) *campestrevix* voluble, foliolis minutis firmioribus subortundis sutomentose-pubescentibus.

LITTLE NAMAQUALAND.—Between Koussie and Garip, Drège.

59. **R. Schlechteri**, sp. nov.

Species ad *R. viscidulam*, Steud., valde accedens. *Caulis* flexuosus. *Stipulae* lanceolatae acuminatae. *Folia* trifoliolata, petiolata, foliola terminalia ovata acuta, 12–15 mm. longa, 10–12 mm. lata, longiora quam lata, foliolis lateralibus ovato-lanceolatis, 10–16 mm. longis, 9–12 mm. latis, petiolulia terminalibus 5 mm. longis, petiolo communi 8–20 mm. longo suffulta. *Pedunculi* 2.6–6.5 cm. longi, pauciflori. *Flores* lutei, 10–12 mm. longi. *Calycis* tubus campanulatus, segmentis linearilanceolatis quam petala brevioribus. *Vexillum* glabrum. *Carina* obtusa, 11 mm. longa.

NAMAQUALAND.—Between Spektakel and Komaggas, Bolus No. 6559; Brachdamm, Schlechter No. 1115. A glandular plant.

Differs entirely from the preceding in the shape of the leaflets; the lateral leaflets are unequalsided, ovate, acute, 10–16 mm. long, 9–12 mm. broad.

IMPERFECTLY KNOWN SPECIES.

R. trichodes, Harvey, l.c. 258 (*Copisma trichodes*, E. Mey., "Comm.," 134).

Stem procumbent, flexuous, very hairy, with patent yellow hairs. *Leaflets* elliptic-oblong, subacute, netted veined, yellow hairy on both sides, gland-dotted beneath, equalling the petiole. *Peduncles* one-flowered, equalling the leaf. *Legumes* unripe, straightish, villous.

CAPE.—Drège, no locality assigned.

The "Index Kewensis" refers this species to *R. Totta*, D.C.

R. nitida, Harvey, l.c. (*Copisma nitidum*, E. Mey., "Comm.," 136).

Stem terete, filiform, voluble, glabrescent. *Leaves* broadly triangular, lateral scarcely unequalsided, glabrous, and shining on both sides, ciliate. *Racemes* equalling the leaves, few-flowered. *Flowers* suberect. pedicels equalling the pubescent, nearly glandless calyx. *Pod* falcate, glabrescent, about three-seeded.

CAPE.—Grassy hills near Omsamculo, Drège.

If the description is correct, this cannot be a species of *Rhynchosia*, as the ovary in this genus only has two ovules.

EXCLUDED SPECIES.

R. Rehmanni, Schinz = *Fageha bituminosa*, D.C.

R. ficifolia, Benth. = *Pueraria ficifolia*, C.B., Clarke,

THE GENERA ALOE AND MESEMBRYANTHEMUM

AS REPRESENTED IN THUNBERG'S HERBARIUM.

By N. E. BROWN.

ALOE.

THE Kew Herbarium having acquired, through the courtesy of Professor H. O. Juel, Director of Upsala Botanic Garden, the loan of the type specimens of the genera *Aloe* and *Mesembryanthemum* collected in South Africa by Thunberg, I have had the opportunity to carefully examine and compare them with the material at Kew, and have thought it would be useful to future workers upon these genera to give an account of the specimens and my identifications of them.

During my long career at Kew I have on several previous occasions had the privilege of being able to examine Thunberg's types of various orders and genera, and in all cases have made notes of my comparisons upon the sheets in the Kew Herbarium. In all the genera that I have previously examined Thunberg's material has usually been quite satisfactory and often excellent, but in the genus *Aloe* this is unfortunately not the case, for the specimens of most of the species consist of detached leaves and flower-stems, and in some instances the leaves and flowers of what are supposed to represent one species belong to two different species or even different genera. This is so unlike the usual excellence of Thunberg's specimens that it must be attributed to the succulent nature and often large size of the plants and the difficulty Thunberg must have had in properly making good, dried specimens of them while travelling as he did at that date. It seems evident that his specimens of detached leaves and flowers must have become mixed in the press, and that he sometimes forgot which leaves belonged to the flowers; hence, the mixture.

As I have very rarely found a locality written upon the sheet containing the specimen, Thunberg must have had some separate notes connecting the localities, as given in his "*Flora Capensis*," ed. Schultes, with the specimens.

From Professor Juel's account of Thunberg's Herbarium, it would appear that Thunberg brought his collection, or some part of it, to Europe mounted upon small sheets of thin paper, and remounted them on larger sheets of thicker paper after his return to Sweden. In doing this it is quite possible that some of the mixtures of species may have been made.

In the account of them which follows, the species have been arranged alphabetically under the names that Thunberg used in his "*Flora Capensis*," ed. Schultes, for convenience of reference: and for the sake of brevity I have quoted this work throughout as Thunb., "*Fl. Cap.*"

ALOE, LINN.

A. arachnoides, Thunb., "*Diss. Aloe*," p. 7 (1785), and "*Fl. Cap.*," p. 311.

The specimen consists of three tufts of leaves and three racemes of flowers; all of the latter, I believe, are detached, but have been inserted among the leaves, and only that of the middle specimen can belong, as the other two flower-stems certainly belong, to a

different plant. The leaves belong to *Haworthia arachnoidea*, Duval, "Pl. Succ. Hort. Alenconia," p. 7 (1809); *H. arachnoides*, Haw., "Synop.," p. 96 (1812); *Aloe arachnoidea*, Miller, "Gard. Dict.," ed. 8, No. 17 (1768), as generally understood.

The locality given by Thunberg is Karroo near Zwartkops Zoutpan, in Uitenhage Division.

A. dichotoma, Thunb., "Fl. Cap.," p. 309.

The type specimen of this species consists of two short, longitudinal sections of the stem, one showing the very smooth bark and the other the interior fibre, and one leaf. It is the plant understood by this name, but the earliest publication of it is *Aloe dichotoma*, Masson in "Phil. Trans. Roy. Soc.," Lond. (1776), Vol. LXVI, Part I, p. 310, where the following description is given: "We found a new species of Aloe here (Masson and Thunberg were then, on 2nd November, 1774, ascending the Bokkeveld Mountains), called by the Dutch *Koker Boom*, of which the Hottentots make quivers to hold their arrows; it being of a soft, fibrous consistence, which they can easily cut out, leaving only the bark, which is hard and durable. These trees were about 12 feet high, with a straight, smooth trunk, about 10 inches or a foot (in) diameter and 5 or 6 feet in length, which divided into two branches, and those were again sub-divided into two more branches, which terminated in a bunch of thick, succulent leaves surrounding the stem, spear-shaped, entire, without spines, and hanging down like the leaves of *Dracaena draco*. We did not see it in flower, but by the above characters took it for a new species, and called it *Aloe dichotoma*." This name is usually quoted as having been given by Linnaeus fil., "Suppl.," p. 206, but that work was not published until 1781, five years later than the publication of Masson's description, which is also a far better one than that of the younger Linné.

A. disticha, Thunb., "Diss. Aloe," p. 7 (1785), and "Fl. Cap.," p. 311, not of Linnaeus.

There are two sheets bearing this name, numbered 1 and 2. Both contain very good specimens of a new species of *Gasteria*, nearly allied to *G. verrucosa*, Haw., but differing from that species by the leaves being little more than half as broad at the base, with more parallel sides and more acute and somewhat spine-pointed at the apex, and with smaller and more prominent white tubercles than those of *G. verrucosa*. The following is a description of it:—

Gasteria Thunbergii, N.E.Br.

Leaves strictly two-ranked, 3-7 inches long, 6-8 lines broad just above the sheath, and of nearly equal width to about 6-9 lines below the tip, where they taper sharply into a fine spine-like point. The surface on both sides and the margins are very rough from being thickly covered with small, white, very prominent tubercles. The *flower-stems* are detached, but probably belong to the leaves; they are about 18 inches long, but were evidently longer, and are unbranched, $2\frac{1}{2}$ lines thick. *Flowers* 7-9 lines apart. *Bracts* 3-4 lines long, reflexed, deltoid, acuminate, membranous. *Pedicels* 6-8 lines long, reflexed. *Perianth* (above its articulation with the pedicel) 11 lines long, curved, inflated at the lower part.

Thunberg does not give the locality of this species.

A. humilis, Thunb., "Diss. Aloe," p. 6 (1785), and "Fl. Cap.," p. 311.

This is *A. humilis*, Miller, "Abr. Gard. Dict.," ed. 6, No. 10 (1771).

The specimen consists of a small tuft of leaves and two good flowering stems. The sheet is labelled on the back as being from a plant cultivated in the Botanic Garden at Upsala.

A. lingua, Thunb., "Diss. Aloe," p. 8 (1785), and "Fl. Cap.," p. 312.

This is *A. plicatilis*, Miller, "Gard. Dict.," ed. 8, No. 7 (1768); *A. linguaeformis*, Linn. f., "Suppl.," p. 206 (1781), not of Miller.

The specimen consists of two leaves and two flowering stems. It was collected on mountains at Roo de Zand, near Tulbagh Waterfall.

A. lingua, Thunb., is wrongly quoted by Baker in the "Fl. Cap.," Vol. VI, p. 291 (although omitted from the index), as being partly *Gasteria disticha*, whilst *A. linguaeformis*, Linn. f., is not quoted at all, no attention having been paid to Haworth's correct statement in "Phil. Mag.," Nov. (1827), Vol. II, p. 353, that Thunberg's plant was *Rhipidodendrum distichum*, Willd. in "Ges. Naturf. Fr. Berl. Mag.," Vol. V, p. 165 (1811), which is a synonym of *Aloe plicatilis*.

A. maculata, Thunb., "Diss. Aloe," p. 8 (1785), and "Fl. Cap.," p. 311.

There are two sheets of this marked *a* and *β*.

Sheet *a* contains portions of two leaves and a raceme of what I believe to be **Gasteria pulchra**, Haw., "Synop.," p. 86 (1812).

Sheet *β* (which also has the name "*obliqua*, Haw.," written upon it in pencil) contains the apical half of some large and broad-leaved species of *Gasteria*, totally different from *G. pulchra*, which I cannot identify, and two inflorescences of some species of *Aloe* that are indeterminable.

Thunberg states that *A. maculata* grows on the Outeniqua Mountains, and often cultivated at Capetown, so that his specimens may have been partly obtained from both places.

A. perfoliata, Thunb., "Diss. Aloe," p. 5 (1785), and "Fl. Cap.," p. 310.

There are three sheets bearing this name, marked *a*, *β*, and *γ*.

Sheet *a* contains one leaf about 7 inches long and $2\frac{3}{4}$ inches broad, a tuft of four small leaves about $2\frac{1}{2}$ inches long and 1 inch broad, and two single flowers. These fragments may really belong to three different species: it is not possible to name them with certainty.

Sheet *β* contains part of a leaf and two inflorescences, which I believe to belong to *A. latifolia*, Haw., "Synop.," p. 82 (1812).

Sheet *γ* contains part of a leaf and a flower-spike of *A. vera*, Linn., "Sp. Pl.," p. 320.

The locality given by Thunberg for *A. perfoliata* is the Karroo beyond Swellendam, on the sides of mountains. Sheet *γ*, however, is labelled on the back as having been sent from the West Indies by Forsström, where *A. vera* has been introduced from the Canaries.

A. picta, Thunb., "Diss. Aloe," p. 6 (1785), and "Fl. Cap.," p. 310.

This species is not represented by any specimen so named in Thunberg's Herbarium. He founded it upon *A. perfoliata*, vars. *θ*, *λ*, *μ*, *ν*, Linn., "Sp. Pl.," ed. 1, p. 320.

A. pumala, Thunb., "Diss. Aloe," p. 7 (1785), and "Fl. Cap.," p. 311.

There are two sheets so named, marked *a* and *β*.

Sheet *a* contains a tuft of leaves of *Haworthia fasciata*, Haw., "Suppl.," p. 57 (1819), on the left-hand side, and on the right-hand side a tuft of leaves of *Haworthia granata*, Haw., "Suppl.," p. 57 (1819). Each of these tufts has a flower-stem placed among them, but detached, which may or may not belong to the leaves: I am rather doubtful if they are properly placed.

Sheet *β* contains two leaves and a paniculately branched flowering-stem of what I believe to be *Haworthia margaritifera*, Haw., "Suppl.," p. 55 (1819), but the material is too poor to make the identification certain.

No locality is given for this species, but sheet *a* is labelled on the back as being from plants cultivated in Upsala Botanic Garden.

N.B.—*Aloe pumala*, Linn., "Sp. Pl.," ed. 1, p. 322, is likewise a mixture of two or three species of *Haworthia* with white tubercles, so that it is probable that at that date several species of this type were considered to be forms of one species.

A. retusa, Thunb., "Diss. Aloe," p. 10 (1785), and "Fl. Cap.," p. 311.

The specimen of this consists of part of a plant with two leaves and a flower-stem attached, five detached leaves, and a detached inflorescence, which all belong to *Haworthia retusa*, Duval. "Pl. Succ. Hort. Alenconio," p. 7 (1809), and Haw., "Synop.," p. 95 (1812); *Aloe retusa*, Linn., "Sp. Pl.," ed. 1, p. 322. No locality is mentioned by Thunberg for this species, but the specimen is labelled on the back of the sheet as being from a plant cultivated in Upsala Botanic Garden.

A. sinuata, Thunb., "Diss. Aloe," p. 6 (1785), and "Fl. Cap.," p. 311.

This name was founded by Thunberg upon the plant enumerated by Linnaeus in his "Species Plantarum," ed. 1, p. 320, and ed. 2, p. 458, as *A. perfoliata*, var. ξ , which is based upon "*Aloe succotrina angustifolia spinosa. flore purpurea*," of Commelin, "Horti Medici Rariorum Plantarum," Vol. I, p. 94, t. 48 (1697). This plant of Commelin's is *Aloe succotrina*, Weston. "Universal Botanist and Nurseryman," Vol. I, p. 5 (1770), a book that seems to have been overlooked by all modern authors, yet it contains names that antedate some that have been credited to Lamark, whose first volume of his "Encyclopédie" was not published until 1783, or thirteen years later than Weston's book.

No specimen exists in Thunberg's Herbarium bearing the name *A. sinuata*, but there is a sheet bearing the name *A. succotrina* in Thunberg's handwriting, which Professor Juel, in his "Plantae Thunbergianae," p. 119, seems to have considered to represent *A. sinuata*, Thunb. This sheet, however, contains (a) part of a leaf of *Aloe fruticosa*, Lam., (b) part of a leaf of *Aloe spicata*, Thunb., and (c) part of a raceme of flowers of some species of *Aloe* that is quite indeterminable, but most certainly does not belong to either species represented by leaves upon that sheet.

So that if *A. sinuata*, Thunb., is considered to be founded upon *A. perfoliata*, var. ξ of Linnaeus, it must be placed as a synonym of *A. succotrina*, Weston. But if it is held to be founded upon the sheet bearing the name *A. succotrina* in Thunberg's Herbarium, then the name *A. sinuata* must disappear altogether.

In the synonymy that Thunberg gives under *A. sinuata*, he quotes "*Aloe barbadensis mitior, laete virens et splendens*," Dill., "Hort. Elth.," p. 23, t. 19, f. 24." The figure should be 21, not 24. This figure represents one of the American species of *Agave*.

Although *Aloe succotrina* is not mentioned by Thunberg in his "Prodromus" or "Floras," yet in his "Travels," English translation, ed. 3, Vol. I, p. 213, he mentions, under the date of 14th December, 1772, when at a farm near Slange River, in Oudtshoorn Division, that "Here we saw quickset hedges of *Aloe succotrina*." What is this plant? Can it be a species distinct from *Aloe fruticosa*?

A. spicata, Linn. f., "Suppl.," p. 205 (1781); Thunb., "Fl. Cap.," p. 309.

Thunberg's specimen of this is the type of this species, and consists of portions of two leaves and six detached flowers. The leaves are respectively $10\frac{3}{4}$ and $13\frac{3}{4}$ inches long, and 14 and 16 lines wide at their basal ends, gradually tapering thence into a long and slender subulate point, which at 3 inches below the acute or bluntish tip is only 2 lines broad. The part of the leaf on the sheet named *A. succotrina* above-mentioned (under *A. sinuata*) is just 1 foot long and 14 lines broad at its basal end, and has just the same long, subulate apex. The teeth on the margins are 6-8 lines apart and very small, being not more than half a line long; the apical spine is also small. The margin between the spines is straight or very faintly concave. The flower-stem and pedicels are absent, but the flowers may have been sessile. One flower has what appears to be a bract attached to it, which is 6 lines long and 5 lines broad, and is broadly elliptic, obtuse, 3-nerved. The perianth is campulate in shape, 7 lines long and 5 lines in diameter as pressed, but is probably of nearly the same dimensions when alive. The segments are nearly 3 lines broad, ovate, oblong, obtuse, 3-nerved; they all appear to be free to the base, and are not recurved at the apex. According to Thunberg they are white, with three green veins.

The stamens are exerted 3-4 lines beyond the tips of the perianth-segments, with stout, purple filaments half a line broad, but, according to Thunberg, the filaments are yellowish at the upper part and white at the base. He states that the flower is filled with a purple fluid, which, in the process of drying the specimen, may have stained the filaments, especially as I note that some of the perianth-segments are also tinted with purple. The style protrudes 3-4 lines beyond the stamens.

Thunberg states that this very distinct and hitherto unknown plant grows "in the interior regions, flowering in August": and that it has a thick stem 3-4 feet high, bearing a crown of spreading leaves 2 feet long, and a very dense spike, a foot long, of crowded horizontally spreading flowers.

In the "Gardeners' Chronicle" (1921), Vol. LXX, p. 6, I gave an account of the history of *Aloe spicata* so far as known to me at the time, in which I detailed how I had vainly tried to get a clue from his travels to the locality where Thunberg found this plant, which still awaits rediscovery. There is nothing at all like it in the Kew Herbarium. I am now inclined to think that Thunberg may have seen the plant in cultivation in some garden at or near Capetown.

A. spiralis, Thunb., "Diss. Aloe," p. 9 (1785).

The sheet of this species in Thunberg's Herbarium contains a tuft of leaves of what I believe to be *Apicra spiralis*, Baker in "Jour. Linn. Soc.," Vol. XVIII, p. 217 (1880), and a detached inflorescence of some species of *Gasteria*, which is quite indeterminable.

No locality is mentioned by Thunberg for this species, and the specimen is from a plant cultivated in the Botanic Garden at Upsala.

I do not know if *Apicra spiralis*, Baker, is identical with *Apicra spiralis*, Willd., in "Ges. Naturf. Fr. Berl. Mag.," Vol. V, p. 273 (1811); it is a point that requires investigating.

A. variegata, Thunb., "Diss. Aloe," p. 9 (1785), and "Fl. Cap.," p. 312.

This is *A. variegata*, Linn., "Sp. Pl.," ed. 1, p. 327.

Thunberg's specimen consists of two leaves, with the variegation upon them well preserved, and a single flower. Thunberg does not mention a locality for it, and no information is given on the back of the sheet.

A. viscosa, Thunb., "Diss. Aloe," p. 9 (1785), and "Fl. Cap.," p. 312.

Thunberg's sheet of this species contains three flowering specimens. The central specimen, which I have marked (A) on the sheet, is larger than the others, and is *Haworthia viscosa*, Haw., "Synop.," p. 90 (1812). The lateral specimens, marked (B), may belong to a variety of *Haworthia viscosa*, Haw., or are possibly a distinct species allied to it. I have not seen any plant alive that is quite like them, so that living plants are required for their proper identification.

From the above account it will be seen that Thunberg's collection of the genus *Aloe* is rather a poor one in comparison with other genera in his Herbarium, as it contains little of interest except the type specimens of *Aloe spicata*, *A. dichotoma*, and the new species of *Gasteria* described above.

MESEMBRYANTHEMUM.

Thunberg's collection of this genus is a very interesting one and consists of 109 sheets of specimens, many of which are excellent, although often small, some are scrappy, and, as in the genus *Aloe*, in some instances there is a mixture of species either upon the same sheet or upon different sheets bearing the same name.

When Sonder was preparing his monograph of this genus for the "Flora Capensis," he had Thunberg's specimens to work with, but, upon examining them, I find that he has dealt with them in a most imperfect and unsatisfactory manner. In some cases he has wrongly identified Thunberg's plant with specimens collected by Zeyher in a totally

different region, and described from Zeyher's plant, so that occasionally Thunberg's species is not described in the "Flora Capensis" at all. Added to this, Sonder has accepted the nomenclature of Salm Dyck's fine work on this genus as unimpeachable, whereas it is often entirely wrong, and therefore as the original descriptions of Thunberg and of Haworth are often entirely ignored and those of Salm Dyck, belonging to quite a different plant, substituted, much confusion of nomenclature has been made and the descriptions in the "Flora Capensis" are often quite untrustworthy. And as these errors have not been corrected by Berger in his monograph of the genus, since that work was chiefly compiled from the works of Salm Dyck and Sonder, there is really needed at the present time a carefully worked-out monograph of this very remarkable and highly interesting group of plants. This, as I have already stated in the "Journal of the Linnean Society of Botany," Vol. XLV, pp. 54-55, can only be properly and fully accomplished by a comparison with and study of the unique collection of original drawings of a large number of Haworth's types preserved at Kew.

In the process of comparison and identification of Thunberg's types with specimens in the Kew Herbarium, where Thunberg has mixed two or more species under one name, it has become necessary to take as being the plant he intended to bear the name that specimen which most accurately agrees with his description. As an example of this kind, *M. articulatum*, Thunb., may be cited. This name, as represented by his specimens, includes three distinct species, all very similar in general appearance, and upon a superficial examination they might easily be supposed to be one species, so that his description will apply fairly well to all of them, with the exception of the words "punctate-scarid" as applied to the stem and calyx. As these words only apply to one of the specimens, neither of the others having this character, that particular specimen must be accepted as the type of *M. articulatum*, Thunb.

In the account of Thunberg's specimens which follows, the names are arranged in alphabetical order as they appear upon his sheets; and to the headline I have only given the reference to the well-known edition by Schultes of Thunberg's "Flora Capensis," published in 1823, other references being given under the determination of the specimens, except in those cases where Linnaeus or Linnaeus fil. is the author, as these, of course, take precedence over the names of Thunberg, Aiton, or Haworth. Thunberg gives no authority for the name on the sheets.

For the localities mentioned by Thunberg, I have used the modern form of spelling. Many of his more interesting species were collected in the Van Rhynsdorp and Calvinia Divisions, in the Karroo between the Olifants River and the Bokkeveld Mountains, and on or near Hantam Mountains.

The following is an explanatory list of the abbreviations used for the books quoted:—

Ait., "Hort. Kew."	W. Aiton, "Hortus Kewensis," ed. 1, Vol. II (1789)
Berger, "Mesemb."	A. Berger, "Mesembryanthemum and Portulacaceen" (1908).
"Bot. Mag."	"Botanical Magazine."
D. C., "Plant Grass"	A. P. de Candolle, "Histoire des Plantes Grasses" (1799-1829).
Eckl. & Zeyh., "Enum."	C. F. Ecklon & K. Zeyher, "Enumeratio Plantarum Africae Australis Extratropicæ" (1834-37).
Haw., "Misc."	A. H. Haworth, "Miscellanea Naturalia" (1803).
Haw., "Obs."	A. H. Haworth, "Observations on the genus Mesembryanthemum," Part II (1795).
	N.B.—Part I was published in 1794, but does not contain any descriptions of species, only introductory matter.
Haw., "Rev."	A. H. Haworth, "Revisiones Plantarum Succulentarum" (1821).
Haw., "Suppl."	A. H. Haworth, "Supplementarum Plantarum Succulentarum" (1819).
Haw., "Synop."	A. H. Haworth, "Synopsis Plantarum Succulentarum" (1812).

- Houtt., "Handl."..... M. Houttuyn, "Handleiding tot de Plant- en Kruidkunde," Vol. IX (1778).
- Houtt., "Nat. Hist."..... M. Houttuyn, "Natuurlyke Historie of uitvoerige Beschryving der Dieren, Planten en Mineraalen," Deel II (1773-83).
- Jacq., "Hort. Schoenbr."..... N. J. Jacquin, "Plantarum rariorum Horti Caesarei Schoenbrunnensis" (1797-1804).
- Jacq., "Hort. Vind."..... N. J. Jacquin, "Hortus Botanicus Vindobonensis" (1770-76).
- Lam., "Encyc."..... Chevalier Lamarck, "Encyclopédie Methodique," Vol. I-IV (1783-97).
- Linn., "Sp. Pl."..... C. Linnaeus, "Species Plantarum," ed. 1 (1753).
- Linn. f., "Suppl."..... C. von Linne, "Supplementarum Plantarum Systematis Vegetabilium (editionis decimae tertiae), Generum Plantarum (editionis sextae), et Specierum Plantarum (editionis secundae)" (1781).
- "Nov. Act. Acad. Leop.-Car. Ephem."..... Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum exhibentia Ephemerides," Vol. VIII. Appendix (1791).
- Salm Dyck, "Mesemb."..... N.B.—Twenty-one species of *Mesembryanthemum* are described in this rare book.
J. Salm-Reifferscheid-Dyck, "Monographia generum Aloes et Mesembryanthemi" (1836-63).
- Sond., "Fl. Cap."..... C. W. Sonder in Harvey & Sonder, "Flora Capensis," Vol. II (1861-62).
- Thunb., "Fl. Cap."..... C. P. Thunberg, "Flora Capensis," ed. Schultes (1823).
- Thunb., "Mus. Nat. Acad. Upsal."..... C. P. Thunberg, "Museum Naturalium Academiae Upsaliensis," Auctum, Part II (1827).
- Thunb., "Prodr."..... C. P. Thunberg, "Prodromus Plantarum Capensium," Part II (1800).
- Willd., "Enum. Hort. Berol."..... C. L. Willdenow, "Enumeratio Plantarum Horti Regii Botanici Berolinensis" (1809); Supplementum (1813).
- Willd., "Hort. Berol."..... C. L. Willdenow, "Hortus Berolinensis" (1803-16).
- Willd., "Sp. Pl."..... C. L. Willdenow, "Species Plantarum," Vol. II (1799).

MESEMBRYANTHEMUM, L.

M. acinaciforme, Thunb., "Fl. Cap.," p. 422.

This name appears in both Thunberg's "Prodromus" and his "Flora," but no specimen exists in his Herbarium representing that name as used by him. There is, however, a specimen named "*M. forficatum*" by Thunberg, which is the true *M. acinaciforme*, Linn., and that name is also written upon the sheet as "*acinaciforme*, Mus."

M. albidum, Thunb., "Fl. Cap.," p. 423.

The specimen consists of two well-preserved flowering branches of *M. albidum*, L. The locality of this species is not mentioned.

M. angulatum, Thunb., "Fl. Cap.," p. 426.

There are three sheets of this, marked α , β , and γ . They all belong to the same species, viz. :—

M. Aitonis, Jacq., "Hort. Vind.," Vol. III, p. 8, t. 7 (1776).

M. angulatum, Thunb., "Prodr.," p. 91 (1800).

M. crystallophanes, Eckl. & Zeyher, "Enum. Pl. Afr. Austr.," p. 322 (1836), and Salm Dyck, "Mesemb.," § 60, fig. 2.

Thunberg collected this near the Sundays River in Uitenhage Division, and his specimens are identical with those of MacOwan, "Herb. Austr. Afr." (1873) from near Grahamstown. It is somewhat remarkable that this very distinct plant has not previously been identified with Jacquin's excellent figure.

Thunberg describes the branches as opposite, and they are so on the lower part of the specimen on sheet *a*, but on the upper part of that specimen, and on that on sheet *β*, they are alternate, just as they are on MacOwan (1873).

Sonder in "Fl. Cap.," Vol. II, p. 454, quotes the specimen on sheet *γ* as being "*M. angulatum*, var. *gracile*," but it is merely a small lateral branch 2 inches long, broken from a larger specimen, and very possibly from one of those on sheet *a* or *β*. The Bethelsdorp specimen which Sonder also quotes for his variety *gracile*, and from which he doubtless described, is probably a different plant. Jacquin raised *M. Aitonis* from seeds sent to him by Aiton, so that in all probability they were collected and sent to Aiton by Masson, who accompanied Thunberg on his journeys.

M. apetalum, Thunb., "Fl. Cap.," p. 417.

This is represented in Thunberg's Herbarium by a scrap about 3 inches long, being evidently a short branch from an annual, but it is undoubtedly:—

M. apetalum, Linn. f., "Suppl.," p. 258 (1781): *M. copticum*, Jacq., "Hort. Vind.," Vol. III, p. 7, t. 6 (1776), not of Linnaeus.

M. apetalum was described from a plant cultivated in Upsala Botanic Garden, doubtless raised from seeds supplied by Thunberg. As no specimen exists in the Herbarium of the younger Linné, that in Thunberg's Herbarium must be accepted as the type of the species. It quite agrees with the description of Linnaeus fil., but I cannot match it with any specimen at Kew. It and *M. papulosum*, Linn. fil., are evidently allied species, and require to be searched for and made better known. To this end I here give a translation of the description of *M. apetalum*, Linn. fil., with details concerning Thunberg's specimen included.

The plant is an annual. Stems or branches 1–2 inches long, prostrate, terete, reddish, covered with glittering papillae, as is the whole plant. The branch of Thunberg's specimen including leaves, is 3 inches long and $\frac{3}{4}$ line thick, and glabrous; the papillae have all shrunk and disappeared from the whole of the specimen except on one pedicel, where they are very evident and rather large, but, under a strong lens, in one or two places evidence that the leaves were also papillate can just be traced. Leaves opposite, about $1\frac{1}{2}$ –2 inches long and 1–2 lines broad, linear, or, according to Linnaeus fil., slightly lanceolate, obtuse, succulent, channelled above, convex and somewhat roughly papillate beneath, glabrous. Pedicels $\frac{1}{2}$ –1 inch long, 1-flowered. As pressed, the flower has the calyx-lobes and petals infolded upon the top of the ovary, and is about $3\frac{1}{2}$ –4 lines in diameter. Calyx-lobes unequal, with subulate-cylindric tips. Petals 24, distant, very short, setaceous, white, resembling filaments without anthers, incumbent upon the top of the ovary. Stamens with filaments like the petals, but shorter; anthers fuscous. Ovary large, convex and smooth above. Stigmas 5, about $1\frac{1}{2}$ line long, erect in a column, with recurved tips.

This is evidently a remarkable and very interesting species, although with insignificant flowers. Any reader who may discover it would confer a favour if he or she would kindly send seeds of the plant to me, addressed to the Royal Botanic Gardens, Kew, or through Dr. Pole Evans or Mrs. Bolus.

Jacquin's figure of *M. copticum* seems so exactly to represent the plant of which Thunberg's specimen is only a single branch that I think there can be no doubt of their identity. Linnaeus fil. seems to have described the petals from flowers that had passed into the young fruiting stage, and become rolled up and "setaceous" and bent down upon the top of the ovary.

M. copticum of Linnaeus is a totally different plant and identical with *Aizoon hispanicum*, Linn.

M. articulatum, Thunb., "Fl. Cap.," p. 415.

There are two sheets bearing this name, marked 1 and 2, which contain three distinct species, all very similar in general appearance, and Thunberg's description will apply to all of them, with the exception of the words "punctato-scabri" and "punctato-scabridum"

as applied to the stem and calyx. These words I interpret to mean, rough from being covered with raised points. The only specimen having this character is on the left-hand side of sheet 2; it seems to have been turned over when remounted, therefore the surface-structure is obscured by the adhesive matter upon it, so that the minute acute points or minute conical hairs can only be clearly seen on a few parts of the specimen, which has alternate branches, all directed to one side of the stem. Each branch ends in an oblique compound cyme composed of 3-6 branchlets bearing 1-3 flowers, and these branchlets are also directed to one side. The specimen seems to be from a growing plant just coming into flower; with apparently linear, half-terete, or perhaps channelled, acute leaves $2\frac{1}{2}$ -5 lines long.

The right-hand specimen on this same sheet has its branches placed at right-angles to each other, and they and the calyx were smooth when alive. But this smoothness is obscured on the dried specimen by minute, whitish blisters in places, caused by some secretion of lime or salt. The calyx seems also to be different in form from that of the right-hand scabrid specimen.

Sheet 1 contains two specimens, both alike, which differ from the specimens on sheet 2, by having the stems and calyx covered with blunt papillae (not points), which Thunberg in other cases of the same structure describes as "papilosis." The branches also differ from those of the scabrid plant in being placed at right-angles to each other. Therefore, as Thunberg's description of *M. articulatum* does not fully apply to any other than the left-hand specimen on sheet 2 bearing that name in his Herbarium, and especially as it well agrees with that specimen in the most important character he mentions, that specimen must be taken as being the type of that species, and its synonymy will be:

M. articulatum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, append., p. 10 (1791), and "Fl. Cap.," p. 415.

M. secundum, Thunb., "Mus. Nat. Acad. Upsal.," Part II, p. 12, name only (1827).

This specimen of *M. articulatum* seems to me to be the same plant as *M. secundum*, Thunb., which shows the scabrid character much more clearly. And I consider both to be the same species as Burke No. 468, from Dikkop Flats near the Fish River, in Albany Division; and Burke and Zeyher No. 705, which is probably from the same region. As Thunberg collected near the Fish River on the Uitenhage side of it, he may have obtained his specimen from that region. But the localities he gives for *M. articulatum* are "the Karroo near the Olifants River, towards the Bokkeveld Mountains, in Zwartland, and elsewhere." As none of his specimens have the locality attached to them, it is impossible to assign the particular locality to any of them.

The other two species under "*M. articulatum*" I am unable to identify, and consider it inadvisable to describe them from those specimens.

At the original place of publication four varieties are enumerated, and var. δ is described as "*major procumbens*" instead of "*minor procumbens*" as given in Thunberg's "Fl. Cap.," ed. Schultes.

Sonder in "Fl. Cap.," Vol. II, pp. 434 and 435, quotes *M. articulatum*, Thunb., as a synonym partly of *M. junceum*, Haw., and partly of *M. granulicaule*, Haw. These determinations are utterly wrong, for neither of Thunberg's specimens is like either *M. junceum* or *granulicaule* of Haworth, both of which have been completely misunderstood and wrongly described by Sonder, both being much more slender species than the plants mistaken for them by Sonder, and by Berger, who has mainly copied from Sonder and other works.

It may not be out of place to call attention to the fact that in consequence of Haworth's *M. junceum* being completely misunderstood, and because many of the species belonging to this group are very similar in appearance, at least half a dozen different species are to be found in Herbaria named *M. junceum*, not one of them, so far as I have seen; being correct.

M. aureum, Thunb., "Fl. Cap.," p. 425.

The type consists of one small specimen, collected on the Karroo, between the Olifants River and the Bokkeveld Mountains. As it is not the *M. aureum* of Linnaeus, Sonder renamed it:—

M. auratum, Sonder in "Fl. Cap.," Vol. II, p. 449.

M. aureum, Thunb. in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 16 (1791), not of Linnaeus.

It is evidently an erect shrub, with straight stems or main branches $1-1\frac{1}{2}$ line thick, swollen at the nodes, with short, opposite, nearly erect or very ascending branchlets, that are densely papillate and of a pale biscuit colour as described by Sonder, but the main stem has a smooth, greyish-brown bark. Leaves opposite, 3–5 lines long and 1 line thick, obtuse or subacute, apparently subterete, channelled down the face and seemingly papillate, glabrous. Flowers solitary or 2–3 in a cyme at the ends of the branchlets. Pedicels $1-1\frac{1}{2}$ line long, papillate. Calyx papillate; tube (ovary) obconic, about 5 lines in diameter in young fruit; lobes about 3 lines long, broadly ovate, with a blunt dorsal point below the apex of the membranous margin. Petals wanting on the specimen, but described by Thunberg as linear and yellow. He also states that the stigmas are 5, subulate, acute, and erect.

I have given the above description of Thunberg's type specimen because Sonder has evidently confused it with another species collected by Zeyher, from which he has partly described, since he gives as a character, "rudiments of old remaining leaves very spinous," which is not at all the case on Thunberg's specimen.

M. barbatum, Thunb., "Prodr.," p. 89, and "Fl. Cap.," p. 418.

One sheet containing three branches of—

M. barbatum, Linn., and of "Bot. Mag.," t. 70.

M. stelligerum, Haw., "Synop.," p. 278 (1812), and "Rev.," p. 190; Sond. in "Fl. Cap.," Vol. II, p. 447; excluding synonyms, not *M. stelligerum*, Haw. in "Phil. Mag.," (1824), Vol. LXIV, p. 61.

The modern monographers of this genus, Sonder and Berger, do not seem to have noted that Haworth, finding he had misunderstood the plant to which Linnaeus gave the name *M. barbatum* and had originally described that plant under the name of *M. stelligerum*, and another species under the name of *M. barbatum*, has corrected that error in the "Philosophical Magazine" (July, 1824), p. 61, where he transposes the two names, so that the correct synonymy for *M. barbatum*, Linn. is as given above, and for *M. stelligerum* as follows:—

M. stelligerum, Haw., in "Phil. Mag." (1824), Vol. LXIV, p. 61, not of Haw., "Synop.," p. 278.

M. barbatum, Haw., "Synop.," p. 277; Sonder in "Fl. Cap.," Vol. II, p. 446, excluding synonyms; not of Linnaeus nor of Berger.

The *M. barbatum*, Berger, "Mesemb.," p. 80, on account of its having acute papillae is certainly neither *M. barbatum*, Linn., nor *M. stelligerum*, Haw., both of which have blunt papillae. It is possibly *M. intonsum*, Haw. I do not know what *M. stelligerum*, Berger, "Mesemb.," p. 82, may be, but from the more numerous bristles described as present at the apex of the leaf, it is probably distinct from both *M. barbatum* and *M. stelligerum*, and is possibly a garden hybrid.

Thunberg states that *M. barbatum* grows upon Paardeberg, at Plantam, and elsewhere. But doubtless he confused two or more species in his mind, from seeing similar species in different localities.

M. bellidiflorum, Thunb., "Fl. Cap.," p. 418.

This seems to be represented by a sheet bearing the name "*M. bellidifolium* (?)," containing a single flower with two bracts at the base of its pedicel. It is quite indeterminable. No locality is mentioned for it.

M. bicolorum, Thunb., "Fl. Cap.," p. 427.

This is represented by a sheet labelled "*M. bicolor*," containing three short flowering branchlets from a plant cultivated in Upsala Botanic Garden, which seem to belong rather to *M. coccineum*, Haw., than to *M. bicolorum*, Linn.

M. bracteatum.

There are three sheets bearing this name, one of which has been named by Sonder, the others are respectively marked *a* and *β*. Sheet *a* contains four pieces of a plant cultivated in Upsala Botanic Garden, the others were collected in South Africa. This species is not enumerated in Thunberg's "Flora Capensis."

Sheets *a* and *β* contain specimens of—

M. mutabile, Haw., "Obs.," p. 377 (1795).

The other sheet is correctly named by Sonder—

M. bracteatum, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 185.

M. calamiforme, Thunb., "Fl. Cap.," p. 417.

One small specimen, which is *M. calamiforme*, Linn. Thunberg does not mention a locality for it.

M. capillare, Thunb., "Fl. Cap.," p. 419.

There are two sheets bearing this name, marked *a* and *β*, and the specimens on each sheet are separately described by Thunberg as forms *a* and *β* of *M. capillare*.

Sheet *a* contains two specimens, which have been marked by Sonder "*a*" and "*β*." The right-hand specimen, marked "*β*" quite agrees with Thunberg's diagnosis of *M. capillare* and his description *a*. This plant is not *M. capillare*, Linn. f., but is—

M. brevisfolium, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 188 (1789), and of Haw., "Obs.," p. 274 (1795), but not of Haworth's later works nor of other authors.*

M. capillare, Thunb. in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 13 (1791), and "Fl. Cap.," p. 419, as to description *a*, which is the only description given at the original place of publication, description *β* being absent from that work; not of Linnaeus fil.

M. subglobosum, Haw., "Misc.," p. 62 (1803).

The specimen is a branch or portion of an erect glabrous shrublet, with the main stem about 1 line thick, and the branchlets about $\frac{1}{2}$ line thick, distinctly papillate. Leaves opposite, about $1\frac{1}{4}$ – $1\frac{1}{2}$ line long and 1 line thick, somewhat subglobose or very shortly

* I here wish to point out that *M. brevisfolium* of Salm Dyck, of Sonder, and of Berger is a totally different plant from the *M. brevisfolium*, Ait., differing from it by having longer leaves, which are not terete or subglobose, but flat above and keeled beneath, and has very different flowers. Its correct synonymy is—

M. erigeriflorum, Jacq., "Hort. Schoenbr.," Vol. IV, p. 39, t. 477 (1804).

M. lateriflorum, D. C., "Plant. Grass.," t. 164 (1828).

M. brevisfolium, Salm Dyck, "Mesemb.," § 50, f. 4 (after 1836); Sonder, "Fl. Cap.," Vol. II, p. 342; and Berger, "Mesemb.," p. 95, not of Aiton, nor of Haworth.

This is only one example out of many of the confusion that has been made by later authors accepting as correct the names given by Salm Dyck, apparently without the slightest investigation.

In Haworth's time there were evidently two plants in cultivation under the name of *M. brevisfolium*, neither of them being the same as *M. erigeriflorum*. In 1795, Haworth considered them to be forms of one species, but afterwards separated them.

One of them, as described by Haworth ("Obs.," p. 276), had leaves which "rarely measure a quarter of an inch," and were covered with rounded (not acutely pointed) papillae, or, as Haworth states, the papillae "were by no means pubescent." This plant was the true *M. brevisfolium*, Ait. The other Haworth describes at first as *M. brevisfolium* var. *longum*, and states that it has leaves up to $1\frac{1}{4}$ inch long, with "pubescent papulae which point downwards and looked in a microscope like minute hooks of glass or ice." This plant is quite unknown to me. In his "Synopsis" and later works Haworth retains this plant alone under the name *M. brevisfolium*, and places the true *M. brevisfolium*, Ait., as a synonym of *M. subglobosum*, Haw. But Aiton's name, being the older, must be retained.

ellipsoid, papillate, but apparently with smaller papillae than those on the stem. Flowers solitary at the ends of the short branchlets. Pedicel $2\frac{1}{2}$ -3 lines long, slender. Calyx 5-lobed, papillate; tube (ovary) with a very distinct constriction under the base of the lobes and thereabout 1 line in diameter, and there is also a slight constriction at its base, where it joins the pedicel; lobes unequal, $\frac{1}{2}$ -1 line long, broadly ovate, obtuse. The other parts of the flower do not admit of examination, and the petals are destroyed. There are some better specimens of this plant at the British Museum, collected by Masson, who travelled with Thunberg. These have leaves 1-2 lines long, varying from subglobose to cylindric. The pedicels vary from 1-6 lines long, and the corolla appears to be about 9-10 lines in diameter, with petals about 4 lines long. The constrictions on the calyx-tube are not so distinct as on Thunberg's specimen.

Thunberg describes the flowers as minute and red. His specimen quite agrees with an original drawing of the typical *M. brevifolium*, Ait., at Kew. The peculiar constriction of the calyx-tube is a very marked feature of both Thunberg's specimen and the drawing of the type plant, in which the corolla is represented as being 8 lines in diameter, with apparently 25-30 acute purple petals.

M. brevifolium, Ait., and *M. sessile*, Thunb., are evidently allied species, although one is papillate and the other smooth.

The left-hand specimen, marked "a," resembles *M. brevifolium*, but the branches spread more widely, the leaves are up to 4 lines long, and cylindric and covered with rounded papillae, and the pedicels 8 or more lines long. The calyx-tube below the lobes is also shorter, and the petals are only 2 lines long. It is doubtless the plant alluded to by Thunberg at the end of his description "a," by the words "*Alia est hujus varietas: foliis magis teretibus, subpapulosis*" ("another, with more terete subpapulose leaves, is a variety of this"). I am not able to identify this specimen with any at Kew, nor is there any like it among those collected by Masson at the British Museum. It is quite distinct from *M. erigeriflorum*, having terete (not trigonous) leaves. More ample material is needed to determine whether it is a variety of *M. brevifolium* or a distinct species.

Thunberg collected these two specimens on hills near the Olifants River, towards the north, in Van Rhynsdorp Division.

Sheet β contains two specimens of a plant stated by Thunberg, under description β , to grow near the Gamtoos River, in the Humansdorp Division. It is totally different from the specimens on sheet α , and is an exceedingly distinct species, differing from every other in the genus by its very slender filiform branchlets.

This plant is—

M. capillare, Linn. f., "Suppl.," p. 260 (1781); Thunb., "Fl. Cap.," p. 419, as to description β only. (This description, nor any corresponding to it, does not occur under Thunberg's original description cited above under *M. brevifolium*.)

An irregularly much-branched shrublet, with very slender, woody branches and branchlets, the ultimate of which are filiform and scarcely a quarter of a line thick, thickened at the nodes in a somewhat bead-like manner, of a dark reddish-brown or chocolate colour (possibly purple when alive), smooth and shining on the older parts, but marked with minute white dots on the younger parts, the dots being minute dried-up papillae. The leaves have mostly fallen; those that remain are 1-2 lines long and less than half a line thick, and, according to Thunberg, slightly flattened above, obtuse, papillate. Pedicels solitary, 3-6 lines long. The specimen is not in flower, and the few capsules upon it are damaged, but appear to be very shortly and broadly obconic, about 2 lines in diameter. Thunberg describes the flowers as being "minute, reddish, with reflexed sepals."

I do not match this plant with any specimen in the Kew Herbarium. The swollen nodes (resembling tiny oval beads) of its remarkably slender, brown branchlets distinguish it at once from all other species.

I find that there is no specimen of this plant in the Herbarium of Linnaeus fil. at the Linnean Society, so that, as in so many other cases, he undoubtedly described from Thunberg's specimens, which must be regarded as the type.

M. ciliatum, Thunb., "Fl. Cap.," p. 416.

One sheet of good specimens.

This plant has hitherto been included in the genus *Mesembryanthemum*, but its appearance is so distinctive that I expected it would differ in floral structure from that genus, and upon dissection I found this to be the case; for besides the very remarkable character of the ring of deflexed hairs at the base of the leaf-sheaths, I find the petals are united into a short tube at their base, the ovary is nearly superior, and the capsule of quite a different type from that of *Mesembryanthemum*. I therefore propose to found the genus *TRICHOCYCLUS* for its reception and that of two allied species, which will be described at the end of this account of Thunberg's species.

TRICHOCYCLUS, N.E.Br.

Dwarf bushy plants with erect succulent branches. Leaves opposite, united below into a short sheath, which is ciliate all around its very base with deflexed hairs, persistent. Calyx 5-lobed. Petals numerous, united into a short tube at the base. Stamens numerous, perigynous. Ovary nearly superior, with only a short basal portion immersed in the calyx-tube, 5 (-6?) -celled; stigmas 5 (-6?), subulate; ovules, few in each cell. Capsule 5 (-6?) -valved; valves sub-erect when open, wingless, and without expanding keels, but with a pair of erect, free, ovate, obtuse processes arising from their base within that close over the seeds and somewhat resemble valves of an inner capsule: they are probably the homologues of the cell-wings of the capsule of *Mesembryanthemum*. Seeds few in each cell, compressed. The name is derived from the Greek *Thrix*, a hair, and *Kyklos*, a circle, in allusion to the circle of deflexed hairs at the base of the leaf-sheaths.

T. CILIATUS, N.E.Br.—*Mesembryanthemum ciliatum*, Ait. "Hort. Kew," ed. 1, Vol. II, p. 179 (1789); Thunb. in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 11 (1791).

Collected between the Olifants River and Bokkeveld Mountains, in the Van Rhynsdorp Division.

This plant is identical with Pearson No. 3941, from south-west of Bitterfontein, in the same region.

M. corallinum, Thunb., "Fl. Cap.," p. 416.

One specimen in a flowerless condition. The original reference is

M. corallinum, Thunb. in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 12 (1791).

A shrublet, hard and woody except at the young parts, much-branched, and apparently stout at the basal part, the lower part of Thunberg's type being 4 lines thick, the branches much less stout, and the branchlets rather less than 1 line thick, constricted at the nodes into short joints 2-3 lines long, glabrous, evidently smooth when alive. Leaves 2-3 lines long and about $\frac{3}{4}$ line broad and as much in thickness, subterete, slightly flattened above, obtuse or subacute, incurved-spreading, glabrous. According to Thunberg, the flowers are solitary and terminal at the apex of the branchlets, sessile. Calyx 5-lobed, green; lobes terete erect. Petals many, linear, spreading, white.

Thunberg states that this grows on dry hills near the Olifants River, towards the north, and in the Karroo below the Bokkeveld, and elsewhere.

I have not been able to match this with any specimen in the Kew Herbarium, but there is a larger specimen of it in the British Museum, collected by Masson when travelling with Thunberg.

M. cordifolium, Thunb., "Fl. Cap.," p. 417.

One sheet of this plant, collected in a wood near Zeekoe River, in the Humansdorp Division. It is—

M. cordifolium, Linn. f., "Suppl.," p. 260 (1781), and is identical with Zeyher No. 2621.

M. crassifolium, Thunb., "Fl. Cap.," p. 421.

There is one sheet of this containing two specimens, for which Thunberg gives the locality: hills in Zwartland, Malmesbury Division. These specimens belong to—

M. filicale, Haw., "Misc.," p. 87 (1803).

M. crassifolium, Thunb., "Prodr.," p. 90 (1800), not of Linnaeus.

Thunberg's specimens are identical with MacOwan, "Herb. Austr.-Afr." (1731), from the foot of Devil's Peak.

Sonder in "Fl. Cap.," Vol. II, p. 407, refers *M. crassifolium*, Thunb., to *M. reptans*, Ait., but that is a very different species, with larger, prominently dotted (not smooth) leaves and much larger yellow or white (not rosy) flowers.

M. debile, Haw., also placed by Sonder as a synonym of *M. reptans*, Ait., is likewise totally distinct from the latter, and I believe it to be the plant distributed by Schlechter under No. 9241 as *M. filicale* (which it certainly is not) from French Hoek.

M. criniflorum, Thunb., "Fl. Cap.," p. 411.

One sheet containing four good specimens, collected on sandhills near Capetown. These specimens agree with those of MacOwan, "Herb. Austr.-Afr." (1745), from near Sea Point. Their correct synonymy is—

M. criniflorum, Linn. f., "Suppl.," p. 259 (1781); Thunb., "Prodr.," p. 88 (1800).

M. limpidum, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 182 (1789).

M. spatulatum, Willd., "Sp. Pl.," Vol. II, p. 1025 (1799), not of Thunberg.

M. tricolor, Willd., "Hort. Berol.," Vol. I, p. 22, t. 22 (1803), not of other authors.

(*N.B.*—Willdenow's work was issued in parts, so that the date 1816 on the title-page is very misleading, that being the date when Volume II was issued as a whole, but it was issued in parts previously.)

The colour of the flower varies.

Sonder and Berger have erroneously quoted *M. tricolor*, Willd., as belonging to *M. pyropaeum*, Haw. They also erroneously quote Houttuyn as being the author of *M. criniflorum*, whilst Thunberg ("Fl. Cap.," p. 411) attributes the name "*M. capense*" to Houttuyn, but neither of these names are to be found at the place quoted in Houttuyn's work; the plant referred to being *M. pugioniforme*, Houtt., "Nat. Hist.," Deel II, t. 53, separately issued as Handleid., "Planten tot de Kraidkunde," Vol. IX, p. 32, t. 53, which represents a form of *M. pomeridianum*, Linn., in fruit, with capsules having 16–20 valves and cells, whilst *M. criniflorum* has only 5 valves and cells to its capsules. But authors have paid little heed to fruit-characters in this genus.

M. crystallinum, Thunb., "Fl. Cap.," p. 413.

There are two sheets of this, containing specimens that appear to be *M. crystallinum*, Linn. The locality given for it by Thunberg is near the Sundays River, but sheet 1 is labelled as being from a plant cultivated in Upsala Botanic Garden.

M. decussatum, Thunb., "Fl. Cap.," p. 414.

One sheet with two specimens upon it, collected on dry hills near the Olifants River, towards the north, in the Karroo below the Bokkeveld. Its synonymy is—

M. brachiatum, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 191 (1789), and Haw., "Misc.," p. 58.

M. decussatum, Thunb., "Prodr.," p. 88 (1800).

This is the same plant as Pearson No. 5544, which has been erroneously distributed as *M. geniculiformum*, Linn., from north of Nieuwerust, in the Van Rhynsdorp Division, which at Kew is represented by a small and very poor specimen that has been grazed by sheep or cattle, but is undoubtedly the same plant as *M. decussatum*, Thunb.

M. deltoides, Thunb., "Fl. Cap.," p. 418.

One sheet with two good branches upon it of *M. deltoides*, Linn., from the mountains of Roodezand Kloof, now New Kloof, near Tulbagh.

M. difforme, Thunb., "Fl. Cap.," p. 423.

One specimen, in unripe fruit. This is a small plant scarcely an inch high, and quite different from *M. difforme*, Linn., as I have elsewhere pointed out, its synonymy being—

M. exiguum, N.E.Br., in "Journ. Linn. Soc. Bot.," Vol. XLV, p. 102 (1920).

M. difforme, Thunb., "Prodr.," p. 90 (1800), not of Linnaeus.

It seems to belong to the same group as *M. rostratum*, Linn. The specimen has a woody root stock bearing a tuft of four growths. Each growth (including flower) is 8-9 lines long and about 4 lines in diameter, with three pairs of leaves. The basal pair are 6 lines long and 1-1½ line broad (the dimensions here given are doubtless much less than those of the living plant), widely spreading, united at the base into a sheath only 1½ line long: they appear to be concave above, very rounded on the back, with a keel at the apex, which is truncate in side view, and has a minute point, glabrous, but covered with minute hard papillae or granules. The second pair are erect and united at the base into a cylindric sheath 4½-5 lines long, with two compressed, acute leaf tips 2 lines long, papillate like the lower leaves and keeled down the back; the sheath in the dried specimen is thin, somewhat membranous, and whitish. Within this sheath the third pair of leaves is enclosed, which appear to be free nearly to the base and just like the basal pair, but they are erect, with their upper surfaces pressed together. They probably ultimately spread widely and become in turn a basal pair. Pedicel 4-5 lines long, with a pair of small leaf-like bracts at its base. This bears a nearly ripe capsule, no flower being present on the specimen. Calyx apparently 5-lobed to the top of the ovary; lobes 2-2½ lines long, some leaf-like, keeled and minutely tuberculate or papillate, the others broadly ovate, with membranous margins. Petals numerous, their shrivelled remains about 5 lines long when wetted. Stigmas 5, about 1½ line long, subulate, recurved-spreading.

Sonder, in the "Flora Capensis," quotes this very distinct species on p. 395 as being a synonym of *M. denticulatum*, Haw., and on p. 399 as a synonym of *M. namaquense*, Sond. The former is a totally different plant, and the latter I have not seen, but it is described as having leaves 3 lines broad and a "peduncle three to four times longer than the leaves." So that it is evident that Thunberg's plant is different from both.

M. digitiforme, Thunb., "Fl. Cap.," p. 412.

One sheet, with two perfect growths and two flowers upon it. Collected in the Karroo between Olifants River and the Bokkeveld Mountains, in the Van Rhynsdorp Division. It is—

M. digitatum, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 181 (1789).

M. digitiforme, Thunb., in "Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 6 (1791)

M. dolabriforme, Thunb., "Fl. Cap.," p. 424.

Two sheets marked 1 and 2.

Sheet 1 contains a specimen of some stemless species in fruit, with two leaves. It is unknown to me, and probably new, but quite indeterminable.

Sheet 2 contains a flowering branch and a separate flower of *M. dolabriforme*, Linn.

The locality given by Thunberg is "on very dry hills at Hantam, near the Roggeveld Mountains."

M. edule, Thunb., "Fl. Cap.," p. 426.

One sheet, containing two flowering branches. Thunberg states that this plant, which is *M. edule*, Linn., grows near Capetown, Zwartland, and elsewhere.

M. emarcidum, Thunb., "Fl. Cap.," p. 415.

Two sheets, marked 1 and 2, stated to have been collected on the Bokkeveld and elsewhere in the Karroo.

Sheet 1 contains two species, a branch (which I have marked "A" on the sheet) and four pieces (which I have marked "B").

"A" is the plant described by Thunberg as *M. emarcidum*, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 9 (1791); *M. anatomicum*, Haw., "Misc.," p. 50 (1803).

This specimen appears to me to be the same as Pearson No. 6434, from Khoms Ravine, in the Khamiesberg. At the original place of publication, Thunberg states that this species has 4 erect stigmas, and grows in the Bokkeveld mountain region, without mention of any other locality. There is a good specimen of it at the British Museum, collected by Masson when travelling with Thunberg.

"B" is *M. expansum*, Linn.

Sheet 2 contains one good specimen of *M. expansum*, Linn.

M. emarginatum, Thunb., "Fl. Cap.," p. 421.

One specimen from Upsala Botanic Garden. The specimen is a poor one, but probably is *M. emarginatum*, Linn.

M. expansum, Thunb., "Fl. Cap.," p. 415.

Two sheets marked 1 and 2. Sheet 1 contains a very fine flowering specimen from Upsala Botanic Garden and sheet 2 a good specimen in fruit. The locality given for this species is near Dutoits Kloof. Both specimens are—

M. tripolium, Linn.

M. expansum, Thunb., "Prodr.," p. 88 (1800), not of Linnaeus.

Thunberg's specimens agree with those of Bolus No. 7930, from Orange Kloof on Table Mountain.

M. falcatum, Thunb., "Fl. Cap.," p. 422.

One specimen. No locality is quoted for it. The plant is *M. lacrum*, Haw., "Obs.," p. 383 (1795); *M. falcatum*, Thunb., "Prodr.," p. 90 (1800), not of Linnaeus.

M. fasciculatum, Thunb., "Fl. Cap.," p. 417.

One specimen, collected near Sundays River, in the Uitenhage Division.

This species has been wrongly quoted by Sonder in "Fl. Cap.," Vol. II, p. 432, as a synonym of *M. tetragonum*, Thunb., which is an utterly different plant of erect bushy habit, whilst *M. fasciculatum* is prostrate.

I have been unable to match *M. fasciculatum*, Thunb., with any specimen in the Kew Herbarium, but it is evidently allied to *M. crassifolium*, Linn., which grows in the region of Capetown (Wolley Dod No. 1625), and is probably the eastern representative of that species, but seems to root much more freely at the nodes than *M. crassifolium* does. The following is a description of Thunberg's plant:—

M. fasciculatum. Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 11 (1791).

Stem prostrate, rooting at the nodes, about $1\frac{1}{2}$ line thick, with arching internodes $1\frac{1}{2}$ –2 inches long, glabrous, smooth, of a pale biscuit colour, bearing at the nodes short, erect, leafy-flowering shoots 1–2 inches high, including the flower, each with 4–5 pairs of

crowded ascending-spreading leaves 6-7 lines long and 1 line thick, which appear to have been nearly terete and very obtuse at the apex, and are described by Thunberg as "scarcely trigonous." Like some other species from the same region, the dried leaves have numerous dot-like blisters formed by crystalline particles of some salt or lime. Pedicel terminal, solitary, bractless, about 5 lines long. Calyx 4-lobed, according to Thunberg, but the lobes cannot be properly seen on the specimen. The flower seems to be in a closed condition, and measures about 5 lines from the base of the calyx to the tips of the apparently lax petals, which are white, according to Thunberg.

M. fastigiatum, Thunb., "Fl. Cap.," p. 413.

One sheet containing two good specimens, collected on very dry hills near the Olifants River, towards the north, in the Van Rhynsdorp Division.

De Candolle, and following him Sonder, and Berger have maintained the name *M. fastigiatum*, Haw., which was published in 1795, and given to the four-years older *M. fastigiatum*, Thunb. the new name of *M. papuliferum*, even overlooking the fact that Haworth had also long before bestowed a new name upon it. Thunberg's name, however, must be maintained, and as no good description of the plant exists, the following is an account of Thunberg's type specimens:—

M. fastigiatum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 7 (1791).

M. erectum, Haw., "Misc. Nat.," p. 53 (1803).

M. papuliferum, D. C., "Prodr.," Vol. III, p. 448 (1828).

An annual $1\frac{1}{2}$ –2 $\frac{1}{2}$ inches high, and as dried, broadly obconic in outline and 2–3 inches in diameter across the flat top, densely covered with watery papillae on stems, leaves, and calyx. One specimen is divided at the base into two, and the other into three main branches, each of which at about half an inch above the base is divided into two or three branches $\frac{1}{2}$ – $\frac{3}{4}$ inch long: each of these either bear one to three flowers at the apex or divide once more before doing so, the whole forming a flat-topped cyme. All the branches thicken upwards in an obconic manner. The leaves have disappeared from the lower nodes, and the few that remain are much shrivelled, 5–7 lines long and 1 line thick; they seem to have been nearly terete, or perhaps more or less flattened above, obtusely rounded at the apex. The flowers seem to be nearly sessile or but shortly pedicellate. Their structure I have not examined, but Thunberg describes the calyx as 5-lobed; the lobes as being ovate, obtuse, purple, and the corolla is snow-white.

I find that there are much larger specimens of this plant at the British Museum, collected by Masson.

The plant later described as *M. fastigiatum*, by Haworth, requires to have a new name; therefore, I propose for it that of—

M. plenifolium, N.E.Br.

M. fastigiatum, Haw., "Obs.," p. 210 (1795), not of Thunberg.

M. filiforme, Thunb., "Fl. Cap.," p. 417.

One specimen, collected at Hantam.

This plant is not a *Mesembryanthemum*, but is a species of *Galenia*, and may therefore bear the following name:

Galenia filiformis, N.E.Br.

Mesembryanthemum filiforme, Thunb., "Prodr.," p. 89 (1800), and "Fl. Cap.," p. 417. *Aizoon microphyllum*, Bartl., in "Linnaea," Vol. VII, p. 541 (1832).

This plant was left undetermined by Sonder, and it seems to be distinct from *Galenia herniariaefolia*, Fenzl., under which Sonder places *Aizoon microphyllum* as a synonym. It was evidently covered with minute watery papillae when alive, and the shrunk cells in some places have somewhat the appearance of pubescence, but I do not perceive any real hairs upon this or on *G. herniariaefolia*, such as Sonder mentions in his Key to the species of *Galenia*.

M. forficatum, Thunb., "Fl. Cap.," p. 422.

The specimen representing this is—

M. acinaciforme, Linn.

M. forficatum, Thunb., "Prodr.," p. 90 (1800), not of Linnaeus.

It was collected on the Karroo.

M. geniculiflorum, Thunb., "Fl. Cap.," p. 414.

The specimen of this is the true *M. geniculiflorum*, Linn., and is labelled "*culta in horto gryphico*." It is identical with Bolus No. 9236, from near the Montagu Baths.

M. glaucum, Thunb., "Fl. Cap.," p. 427.

One sheet containing two specimens. The locality is not mentioned by Thunberg.

The right-hand specimen may possibly be *M. glaucum*, Linn., but the flower is detached.

The left-hand specimen is *M. roseum*, Willd., "Enum. Plant. Hort. Berol.," p. 535 (1809).

M. hispidum, Thunb., "Fl. Cap.," p. 418.

There are five sheets so named, marked α , β , γ , δ , and ϵ , of which Thunberg gives separate descriptions under letters α , β , and γ , which presumably correspond with his sheets so lettered. He evidently regarded them as forms of one species.

Sheet α contains a good specimen of *M. candens*, Haw., "Rev.," p. 186 (1821). It is stated to grow on sandy plains, and is identical with Zeyher No. 2592, from near Port Elizabeth.

Sheet β contains two branches of *M. calycinum*, Haw., "Rev.," p. 187 (1821). It is the plant described by Thunberg under β , for which he gives no locality, but it is the same plant as Bolus No. 5050, from near Mitchell's Pass.

Sheet γ contains one poor specimen of *M. micans*, Linn. It may be the plant described by Thunberg under γ , but scarcely agrees with the description. The locality given for γ is Karroo between Olifants River and the Bokkeveld Mountains, in the Van Rhynsdorp Division.

Sheet δ is named "*M. hispidum* (?)." It is an undescribed species, unlocalized by Thunberg, but identical with a branch mixed with another species collected by Bolus near the Montagu Baths, and distributed under No. 6715, which I describe under the following name :—

M. praecultum, N.E.Br.

An erect bushily-branched shrub, probably less than a foot high, the specimen seen being about 6 inches high, with the stoutest part of the stem less than 1 line thick. The young parts, including leaves and calyx, densely papillate, the old parts with a somewhat smooth greyish or brown bark. Internodes 3–6 lines long. Leaves opposite, 1–2 lines long, subglobose or nearly as thick as long, or shortly cylindric, very obtuse; most of them have fallen from the specimens. Flowers very numerous, produced all along the main and lateral branches at nearly every node, solitary in the axil of each fallen leaf. Pedicel ascending, 3–5 lines long. Calyx 5-lobed to the top of the ovary; lobes subequal, nearly 2 lines long, oblong, obtuse. Petals numerous, about 3 lines long and $\frac{3}{4}$ –1 line broad, obtuse, tapering downwards. Stamens numerous. Stigmas 5, spreading, $1\frac{1}{2}$ line long, slender, subulate, arising from a 5-lobed crown-like structure on the top of the ovary. Capsule $1\frac{1}{2}$ line long and 2 lines broad, shortly and broadly obconic, convex at the top, 5-valved.

Sonder in "Fl. Cap.," p. 442, has referred *M. hispidum* (δ) of Thunberg's Herbarium to *M. brevifolium*, Ait., but that is a very different plant, with totally different and smaller flowers (see under *M. capillare*). The *M. brevifolium*, Sonder, is a mixture of three or more species.

Sheet ϵ contains two branches of what I believe to be *M. floribundum*, Haw., "Misc.," p. 100 (1803). It is not localized.

From the above it will be noted that neither of Thunberg's specimens belong to *M. hispidum*, Linn.

M. laeve, Thunb., "Fl. Cap.," p. 425.

One specimen, collected near Sundays River, in the Uitenhage Division. As the name *M. laeve* had previously been used by Aiton, Haworth changed the name to—

M. Thunbergii, Haw., "Misc.," p. 86 (1803).

M. laeve, Thunb., in "Acad. Leop.-Car. Ephem.," Vol. VIII. Append., p. 16 (1791), not of Aiton.

Thunberg's specimen agrees with those of Rogers No. 4656, collected near Zwartkops, in the Port Elizabeth Division.

This eastern plant is similar to and might be mistaken (in the dried state) for *M. dissimile*, N.E.Br., which grows near Capetown, but differs from that species by having more obtuse leaves, no bracts at the base of the pedicel, and apparently smaller flowers. But the two require to be compared when alive, so that their distinctive characters can be properly contrasted. The prostrate stems of *M. Thunbergii* are 2 lines thick, very smooth, and pale brownish.

M. lanceum, Thunb., "Fl. Cap.," p. 417.

One sheet, containing two good specimens. The locality is not stated. There is another specimen of it from Upsala Botanic Garden, named "*M. tripolium*," and also another mounted on the sheet of *M. oratum* (which see). The original reference for it is *M. lanceum*, Thunb., "Prodr.," p. 89 (1800).

Sonder's description of this, in "Fl. Cap.," Vol. II, p. 455, is fairly correct, except that I believe the stems to be procumbent or straggling. Possibly the plant figured by De Candolle, "Plant. Grasses," t. 47, as *M. expansum*, may be *M. lanceum*, Thunb.; it does not seem to be *M. expansum*, Linn. It also seems closely allied to *M. varians*, Haw., and may prove to be that species when better known.

M. lineare, Thunb., "Fl. Cap.," p. 411.

One sheet of specimens, for which the localities Zwartland and Groenekloof, in the Malmesbury Division, are given. This plant is—

M. gramineum, Haw., "Obs.," p. 470 (1795).

M. lineare, Thunb., "Prodr.," p. 88 (1800).

M. pyropaeum, Haw., "Suppl.," p. 99 (1819).

M. tricolor, Sims, "Bot. Mag.," t. 2144 (1820), and Haw. "Rev.," p. 163 (1821), not of Willdenow.

Thunberg's specimens agree with those of Wolley Dod No. 1796, from sandhills at Duinefontein.

M. gramineum was founded upon the figure and description in Petiver, "Gazophylacium," Vol. I, p. 10, t. 88, f. 6, and "Cat.," Vol. II, p. 4, No. 488, which clearly represents this plant, although Petiver described the petals as yellow; but, as he seems to have had only dried specimens, he was perhaps misled by the yellowish colour they sometimes assume when dried.

Berger, "Mesemb.," p. 38, also quotes *M. clavatum*, Haw., as a synonym of this species, but that name was likewise founded upon a plant figured by Petiver on the same plate (t. 88, f. 7), which represents quite a different plant with creeping and probably perennial stems. The *M. tricolor*, Willd., quoted by the above authors for this species, is *M. crinitiflorum*, Linn. f. (which see).

M. linguaeforme, Thunb., "Fl. Cap.," p. 424.

This is represented by four detached leaves and three flowers, labelled as from a plant cultivated in Upsala Botanic Garden, which certainly do not belong to *M. linguaeforme*, Linn., but probably to—

Glottiphyllum longum, N.E.Br., in Gardener's "Chronicle" (1922), Vol. LXXI., p. 9

Mesembryanthemum longum, Haw., "Obs.," p. 177 (1795).

M. loreum, Thunb., "Fl. Cap.," p. 421.

One flowerless specimen from Upsala Botanic Garden. Thunberg states that it grows on the Karroo between Olifants River and the Bokkeveld. The specimen, however, does not belong to *M. loreum*, Linn., but is probably *M. crassifolium*, Linn., and is identical with Schlechter No. 8316 (from Zout River) and Wolley Dod No. 1625 (from Governor's Cottage).

M. micans, Thunb., "Fl. Cap.," p. 426.

One sheet containing three specimens, collected below the Bokkeveld Mountains.

This plant is not *M. micans*, Linn., and it is erroneously quoted as a synonym of *M. collinum*, Sond., by Sonder in "Fl. Cap.," Vol. II, p. 443. That species, however, was founded upon specimens collected by Zeyher near the Gauritz River, in the Mossel Bay Division, a widely different locality from that of Thunberg's plant, which is a different species and identical with Pearson No. 4903, from Hottentots Kloof, in the Ceres Division, which is stated to be a "weak herb, $\frac{1}{2}$ -1 foot; among bushes." But it is evidently weak and lanky and only partly woody from growing among or in the shade of bushes; otherwise, it is quite identical with Thunberg's shrubby specimens. I propose the following name for it:—

M. exspersum, N.E.Br.

M. micans, Thunb., "Prodr.," p. 91, not of Linnaeus.

A dwarf shrublet, 6-12 inches high, with woody branches about $\frac{3}{4}$ line thick and more slender branchlets. Branches opposite, rather widely diverging at their origin, with internodes 3-15 lines long, the young parts ascending, curved or flexuose, with a smooth brown bark on the old parts, and the young leafy branchlets purplish or brownish, covered with scattered longitudinally compressed papillae. Leaves opposite, 2-4 lines (5-6 lines on Pearson's specimen) long, $\frac{1}{2}$ line thick on the dried specimens, erect or more or less spreading, not united at the base, distinctly concave-channelled down the face, rounded on the back, obtuse, of equal thickness throughout, not thickened at the apex, mostly straight, very densely covered with smaller papillae than those on the stem, whitish. Flowers solitary, terminal; pedicels $1\frac{1}{2}$ -2 $\frac{3}{4}$ inches long, slender, brownish or purplish, with scattered papillae like those on the stem, which become crowded at the apex. Calyx 5-lobed down to the top of the short and very broadly obconic ovary, densely papillate; lobes subequal, $1\frac{1}{2}$ -2 lines long, deltoid or deltoid-ovate, acute, two (or three ?) of them with broad membranous margins, not appendaged. Petals numerous, 4-4 $\frac{1}{2}$ lines long, linear, obtuse or notched at the apex, apparently of some shade of red.

M. moniliforme, Thunb., "Fl. Cap.," p. 415.

One sheet, containing two good specimens of the plant in a resting condition, that is, without either leaves or flowers. The original reference is—

M. moniliforme, Thunb., in "Nov. Act. Acad. Lcop.-Car. Ephem.," Vol. VIII, Append., p. 7 (1791).

Thunberg's specimens represent the plant as about 3 inches high, with stout, crowded branches 4-6 lines thick, with numerous annular constrictions $1\frac{1}{2}$ line apart, giving them the appearance of being formed of a number of flat, button-like beads, dark brown, glabrous.

This is not quite like the plant I described and figured, partly from memory, in the "Journal of the Linnean Society," Vol. XLV, p. 116, t. 5, f. 10; as the joints of that plant, as I remember them, were certainly not so flattened as in Thunberg's specimens, yet for all that, it may be specifically the same.

Thunberg collected it on hills near the Olifants River, towards the north, in the Van Rhynsdorp Division.

M. noctiflorum, Thunb., "Fl. Cap.," p. 414.

One specimen from a plant cultivated in Upsala Botanic Garden. It is *M. noctiflorum*, Linn.

M. nodiflorum, Thunb., "Fl. Cap.," p. 413.

Three sheets of good specimens, all of which belong to *M. nodiflorum*, Linn. It is stated to grow near Olifants River, in the Karroo below the Bokkeveld, in Zwartland, and elsewhere. The specimens quite agree with Schlechter No. 11068, from Eenkokerboom.

Sonder in "Fl. Cap.," Vol. II, p. 452, places *M. apetalum*, Linn. f. and *M. copticum*, Linn. as synonyms of *M. nodiflorum* (in which Berger follows Linn.), but they are both perfectly distinct from it and from each other, *M. copticum*, Linn. being the same as *Aizoon hispanicum*, Linn., and *M. apetalum*, Linn. f. is described above.

M. ovatum, Thunb., "Fl. Cap.," p. 417.

This name is represented in Thunberg's Herbarium by one sheet containing two very distinct species.

The right-hand specimen is identical with Zeyher No. 2624, from the valley and hills of the Zwartkops River, in the Uitenhage Division, and is undoubtedly the plant intended by the name *M. ovatum*, in Thunb., "Fl. Cap.," p. 417, for which Thunberg gives no locality.

The left-hand specimen is a branch of *M. lanceum*, Thunb.

But upon noticing the more erect slope of the letters and the different ink with which they appear to have been written, that the specific name "*ovatum*" was written at a different time from the name "*Mesembryanthemum*," I consulted the original description of *M. ovatum*, Thunb., and found that he there describes a totally different plant, which is not represented by any specimen in his Herbarium, nor can I find any in the Kew Herbarium that corresponds to his description. From this I think it probable that his original specimen must have got lost or destroyed, and that at a later date, when compiling his "Flora Capensis," he mistook the ovate-leaved plant (like Zeyher No. 2624) for his *M. ovatum*, and because he found it did not correspond with his original description, wrote a fresh one, made from the specimen he had before him, which is flowerless, and added such details of the flower as are given, and the statement that it is papulose, from his original description. Thus we have two very distinct species described by Thunberg under the name of *M. ovatum* at different dates. The plant originally described appears to be a very small papillate annual, whilst that later described has long (prostrate ?) stems, and is glabrous and smooth or not conspicuously papillate. The following is a translation of the original description :-

M. ovatum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII. Append., p. 8 (1791), not of any other book or author.

Plant dwarf, diffuse. Root annual, fibrous. Stems several, radical, as long as a finger-nail or an inch long, somewhat erect, simple, terete, leafless or slightly leafy (probably with or without bracts) at the middle, papulose, covered with crystalline scales (can elongated dried watery papillae be intended ?). Radical leaves several (or many), larger, stem-leaves fewer, fleshy, opposite, connate, the length of a finger-nail or an inch long, flat, obovate, tapering below, and again widened at the base, obtuse, entire, papulose above, papulose and covered with hyaline scales beneath, viscid, green or purple. Flowers terminal, solitary, white. Calyx fleshy, 5-partite, papulose, crystalline and scaly: lobes 1 line long, with three a little longer, ovate, obtuse, concave, with a thin membranous margin. Petals numerous, inserted on the calyx and longer than it, linear, entire, white. Stamens very many, inserted at the base of the calyx and shorter than it; filaments distinct, filiform, glabrous, white or purple; anthers ovate, didymous, yellow. Nectary crowning the ovary, affixed in a wreath to the margin of the calyx, crenulate, green. Ovary depressed, 5-grooved, glabrous. Styles 5, erect, subulate, acute. Capsule fleshy, depressed-conical, 5-valved, 5-celled. Seeds numerous.

Thunberg states that this plant grows near the Cape, on sandhills, and that it is common. I cannot, however, as above stated, find any plant in the Kew Herbarium that agrees with the above description. Can this plant be a form of *M. criniflorum*, Linn. f. ?

As the above described plant must take precedence for the name *M. ovatum*, that in Thunberg's Herbarium now bearing that name requires to be renamed, and as I find it to be identical with a plant described by Berger, the following will be its synonymy:—

M. Haeckelianum, Berger in "Engler Bot. Jahrb.," Vol. XLV, p. 224 (1910).

M. ovatum, Thunb., "Fl. Cap.," p. 417 (1823), not of "Nov. Act. Acad. Leop.-Car.-Ephem.," Vol. VII, Append., p. 8 (1791).

M. elongatum, Eckl. & Zey., "Enum. Plant. Afr. Austr.," p. 321 (1836), and in "Linnaea," Vol. XIX, p. 658 (1847).

M. angulatum var. *ovatum*, Sond. in "Fl. Cap., Vol. II, p. 454, partly (1862).

Root-stock perennial. Stems probably annual ascending or perhaps prostrate with ascending branches, up to 18 inches long and $1\frac{1}{2}$ line thick, angular, glabrous, and apparently smooth, with internodes up to 2 inches long. Leaves opposite, $\frac{3}{4}$ – $1\frac{1}{2}$ inch long, 3–9 lines broad, lanceolate or ovate, acute, tapering into a petiole at the lower third, entire, glabrous, apparently smooth, fleshy. Flowers solitary in the forks of the stems or axillary. Pedicels $\frac{1}{2}$ –1 inch long, moderately stout, glabrous, bractless. Calyx very unequally 4-lobed, the two larger lobes 6–10 lines long and 3–4 lines broad, lanceolate or ovate-lanceolate, acute, leaf-like, the two smaller 4–5 lines long, with a membranous ovate basal part and a subulate point. Petals numerous, about 6 lines long and $\frac{1}{2}$ – $\frac{3}{4}$ line broad. Stamens numerous. Style $\frac{1}{2}$ – $\frac{3}{4}$ line long, stout: stigmas 4, short, apparently flat, acute.

Valley and hills of the Zwartkops River, in the Uitenhage Division, Zeyher No. 2624, and without locality, but probably from the same region, a flowerless specimen in Herb. (Thunberg).

Sonder in "Fl. Cap.," p. 454, quotes the plant named *M. ovatum* in Thunberg's Herbarium and Zeyher No. 2623 under *M. angulatum* var. *ovatum* as both being the same plant, whereas they are entirely different, the *M. ovatum*, Herb. (Thunberg), being *M. Haeckelianum*, just described, and Zeyher No. 2623 being *M. angulatum*, Thunb.

M. papulosum, Linn. f., "Suppl.," p. 259 (1781).

One sheet containing three specimens in fruit. This species is not mentioned in Thunberg's "Flora Capensis," nor is there any indication of locality on the back of the sheet. But as it was described by Linnaeus fil. from a plant cultivated in Upsala Botanic Garden, it may have been raised from seed obtained from these fruiting specimens, or these specimens may have come from Upsala Garden; and as they accurately agree with the description of Linnaeus fil., they must be accepted as the type of the species, for there is no specimen of the plant among those from the Herbarium of the younger Linné, and as he described many species from Thunberg's specimens, these may even possibly be those from which he described. He quotes, however, *M. Aitonis*, Jacq. as a synonym of *M. papulosum*. This is an error, for *M. Aitonis* has angular (not terete) stems, quite different leaves, straight (not deflexed) pedicels, and broad obovate (not subulate) tips to the calyx-lobes. *M. Aitonis* is also an eastern species from the Uitenhage Division, whilst *M. papulosum* grows near Capetown, for it is identical with Wolley Dod No. 2864, from the shore between Sea Point and Camps Bay. The plant is an annual, covered with large papillae, and is well marked by the peculiar curvature just below the calyx of the 6–8 lines long pedicels, causing the flowers to nod. The petals are small and yellow, "half as long as the shorter calyx-lobes."

M. pinnatifidum, Thunb., "Fl. Cap.," p. 427.

Two sheets; one contains a specimen from Upsala Botanic Garden. The localities given by Thunberg are Mountains of Roode Zand, near Tulbagh Waterfall, and on the Paardeberg. All belong to

M. pinnatifidum, Linn. f., "Suppl.," p. 260 (1781).

M. pinnatum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 15 (1791).

The specimens agree with those collected by Wolley Dod at Smitwinkel Vley.

M. pomeridianum, Thunb., "Fl. Cap.," p. 427.

One specimen from a plant cultivated in Upsala Botanic Garden. It is *M. pomeridianum*, Linn. To the synonymy of this plant must be added *M. pugioniforme*, Houtt., "Nat. Hist." Deel II, t. 53, separately issued as "Handleid. Plant. Kruidkunde," Vol. IX, p. 32, t. 53; *M. capense*, Houtt., ex Thunb. "Fl. Cap.," ed. Schultes, p. 411. (See note under *M. criniflorum*.)

M. pruinatum, Thunb., "Fl. Cap.," p. 425.

This is represented by one very poor specimen, which is *M. echinatum*, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 194 (1789); *M. pruinatum*, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 17 (1791).

Stated to grow in the Karroo near Loeri River, in the Uitenhage Division, and in Cannaland. Sonder has retained *M. pruinatum* as a distinct species, but it is unquestionably identical with *M. echinatum*.

M. pugioniforme, Thunb., "Fl. Cap.," p. 424.

Two sheets containing scrappy specimens, which belong to *M. pugioniforme*, Linn. The locality given for it by Thunberg is near Verloren Valley and elsewhere. Sheet 2 contains a specimen from Upsala Botanic Garden.

M. ringens, Thunb., "Fl. Cap.," p. 423.

Three sheets marked α , β , and γ . Under this name Thunberg describes four varieties.

Sheet α contains two species, the two upper specimens, which I have marked "1" on the sheet, evidently represent the plant described by Thunberg under var. α as growing below the Roggeveld. They have leaves about 9 lines long, with 4-5 acute teeth (not bristles or cilia) on the margins and 2-3 on the keel. It appears to be *M. murinum*, Haw., "Obs.," p. 165 (1795).

The lower specimen and two detached leaves I think belong to *M. felinum*, Hill.

Sheet β contains two species, the upper left-hand specimen, marked "1," appears to be *M. murinum*, Haw., like the specimens on sheet α . The other two specimens, marked "2," seem to belong to the plant from Hantam, described by Thunberg under var. γ , which is possibly *M. mustelinum*, Haw., "Suppl.," p. 87 (1819).

Sheet γ contains two specimens of a species that I cannot identify: it is probably new, but the material is inadequate for description.

M. rostratum, Linn., Thunb. "Fl. Cap.," p. 424, states that this grows in the Karroo between the Olifants River and the Bokkeveld. But the species is not represented in Thunberg's Herbarium.

M. sabulosum, Thunb., "Fl. Cap.," p. 422.

One sheet, containing two specimens of the plant described and a small plant accidentally mixed with them of *M. pomeridianum*, Linn. The original reference is—

M. sabulosum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 17 (1791).

Thunberg collected it in sandy places in Zwartland and near Saldhana Bay. His specimens are identical with Bolus No. 9004, from open places between Tulbagh Kloof and Pikiniers Kloof: with Stephens No. 7168, between Warmbaths and Modderfontein Farm: and with Stephens and Glover No. 8735, from Het Kruis, which have erroneously been distributed as being *M. pomeridianum*, Linn.

M. scabrum, Thunb., "Fl. Cap.," p. 420.

Three sheets, marked 1, 2, and 3, stated to grow on hills near Capetown.

Sheet 1 contains one very poor specimen of a species that is not determinable, but is not *M. scabrum*, Linn.

Sheets 2 and 3 both contain only specimens of *M. bracteatum*, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 185 (1789).

M. secundum, Thunb., "Mus. Nat. Acad. Upsal. Auct.," Part II, p. 12 (1827), name only, no description.

This is not quoted in Thunb. "Fl. Cap." The specimen is identical with those of Burke No. 468 and is *M. articulatum*, Thunb. (which see).

M. serratum, Thunb., "Fl. Cap.," p. 427.

One sheet, containing one flowerless branch. It is possibly *M. tenuifolium*, Linn., but certainly not *M. serratum*, Linn. No locality is given for it.

M. sessile, Thunb., "Fl. Cap.," p. 419.

One specimen in fruit, collected in the Karroo between the Olifants River and the Bokkeveld Mountains. The original place of publication is—

M. sessile, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 14 (1791).

Evidently a woody shrub or shrublet. The type consists of a straight terete main branch about $1\frac{1}{4}$ line thick at the base, giving off numerous very short opposite leaf-bearing and flowering branchlets 2-5 lines long, including the flowers, and three spreading alternate branches $1-1\frac{1}{2}$ inch long (but are broken at the ends and have been longer) and more than $\frac{1}{2}$ line thick; internodes 2-4 lines long; bark brown. Leaves 1-2 closely placed pairs to each branchlet. $1-1\frac{3}{4}$ line long and about as broad, probably larger when alive, half-globose, rounded at the apex, flat or slightly concave (from shrinkage?) above, very convex on the back, glabrous, smooth, apparently pellucid-dotted. Pedicel terminal and solitary on the short lateral branchlets, 1 line long, not bracteate (as Sonder wrongly states), gradually passing into the obconic calyx-tube, which is $1\frac{1}{2}$ line long. Calyx 5-lobed; lobes $1-1\frac{1}{2}$ line long, ovate, obtuse. Flowers red, *ex* Thunberg, but the specimen is in unripe fruit. Capsule 2-2 $\frac{1}{4}$ lines in diameter, with 5 acute ridges on the flattish top, purplish, *ex* Thunberg.

I have not been able to match this with any specimen at Kew.

Sonder has placed *M. cymbiforme*, Haw., "Obs.," p. 264, as a synonym of *M. sessile*, Thunb., but that species is a totally different plant, with much larger leaves than those of *M. sessile*, Thunb.

M. spathulatum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VII, Append., p. 5 (1791), not of Willdenow.

This species is omitted from all other books, including Thunberg's "Flora Capensis," and no specimen so named exists in his Herbarium. *M. spathulatum*, Willd., which seems to have hitherto been mistaken for Thunberg's plant by Willdenow, De Candolle, Sonder, and Berger (who wrongly quotes the name as "*M. spathulifolium*"), is an annual, and is the same as *M. criniflorum*, Linn. f., which grows near Capetown, whilst *M. spathulatum* of Thunberg is described as a perennial, from the Roggeveld region, and therefore not at all likely to be the same species. The following is a translation of Thunberg's description of *M. spathulatum*.

A stemless perennial. Root thick, fleshy, with fibres, divided above into several little stems. Leaves about 6, radical, approximate, erect, alternately opposite, about as long as a finger-nail ("unguicularia"), connate (at the base), ovate, obtuse, with a

point (apiculate). tapering and smoothish at the basal part, papillate-scabrid at the upper part, slightly concave above, convex on the back, entire, green, with a red margin, the two lower withered. Flower solitary, from the centre of the leaves, radical, pedunculate, white. Peduncle shorter than the leaves, 1-flowered. Calyx 5-lobed; lobes ovate-lanceolate.

It grows on the intermediate Roggeveld, and was out of flower in November.

Although no specimen in Thunberg's Herbarium is named *M. spathulatum*, I find in it an unnamed specimen that so exactly agrees with the original description of that species, that I think there can be no doubt whatever that it is the specimen of *M. spathulatum* from which he made his description, but that he omitted to write the name on the sheet, thus probably causing it to be omitted from his "Flora Capensis." This specimen is marked on the back of the sheet merely as having been collected at the Cape of Good Hope by Thunberg. The following is a description of the specimen:—

A stemless perennial. Root-stock fleshy, about 2 inches long and $\frac{3}{4}$ inch thick at the upper part, where it divides into a few branches 3–4 lines long and 2–4 lines thick, each crowned with a small rosette of leaves; the lower part is abruptly contracted into a more slender portion giving off fibres. Leaves 4–6 in each rosette or growth, opposite, crowded, erect, or the outer spreading and withering; each leaf 4–7 lines long, 4–5 lines broad at the apical part, spathulate-obovate, very obtusely rounded at the apex, with a short apiculus, narrowed into a stout petiole-like part from about the middle to the base, where they are very shortly united, the upper part on both surfaces scabrid-papillate, from being covered with small, crowded, hard tubercles, which when highly magnified are seen to be themselves rough from being covered with very minute, hard, conical papillae; the lower or petiolar part is smoother; they appear to have been flat or perhaps slightly concave above, convex on the back. The specimen is flowerless, but there is a detached capsule, which evidently belongs to the plant, as the calyx-lobes are covered with the same peculiar tubercles, which on the lower part pass into crowded smooth dots, that are probably slightly prominent on the living plant. This capsule is 6 lines in diameter, hemispheric and 5-valved, with 5 calyx-lobes and a piece of a slender pedicel 1 line long, that is not dotted, attached to it. The calyx-lobes are 2–3 lines long, broadly ovate, apparently acute.

This very distinct plant is nearly allied to *M. calcareum*, Marloth, which also has the same peculiar type of tubercles upon its leaves, but they are longer, differently rounded at the apex, and are without the little apiculus present on those of *M. spathulatum*.

M. spinosum, Thunb., "Fl. Cap.," p. 420.

Three sheets, marked α , β , and γ , collected in the Karroo between Olifants River and the Bokkeveld Mountains, and at Hantam.

Sheet α contains a specimen identical with "*M. spinosum*, α ," of Herb. Drège, but it is not *M. spinosum*, Linn., and is undescribed. I therefore describe it as—

M. aculeatum, N.E.Br.

A shrub resembling *M. spinosum* in habit, with 3-forked spiny flowering branches. Branches straight, not intricately intermingled. Internodes 4–12 lines long, with grey or brown bark on the old parts, the younger parts, including the spines, which are 6–9 lines long and nearly or quite half a line thick at their base, marked with elongated immersed dots, glabrous. Leaves opposite, short and stout, shortly united at the base, $1\frac{1}{2}$ – $2\frac{1}{2}$ lines long, $1\frac{1}{2}$ –2 lines broad and 1 line thick on the dried specimens, flat above, slightly keeled beneath, obtuse or apiculate, spreading, each with a young shoot 2–3 lines long in its axil, bearing two pairs of crowded leaves, obscurely gland-dotted. Flowers solitary from the tufts of leaves on the widely spreading spines. Pedicels 1 line or less long. Calyx 5-lobed, about $1\frac{1}{2}$ line in diameter across the short and very broadly obconic (or hemispheric?) tube (ovary); lobes subequal, about 1 line long, ovate, obtuse, three with broad

membranous margins. Corolla probably about 5 lines in diameter, with petals about 2 lines long. Stamens numerous, stigmas 5, erect, 1 line long, stout at the base, tapering to an awn-like point.

Drège's specimen was collected in the Nieuweveld, between Rhenoster Kop and Ganze Fontein, in the Beaufort West Division. The above description, however, is made entirely from Thunberg's specimen.

Sheet β contains two specimens of a very distinct spiny species that I cannot match with any specimen at Kew, but it is too imperfect to describe.

Sheet γ contains a specimen of a species allied to the true *M. spinosum*, Linn., but is apparently distinct. Masson also collected it; his specimens are at the British Museum. In Herbaria and books several species are included under the name *M. spinosum*.

M. splendens, Thunb., "Fl. Cap.," p. 414.

One specimen from Upsala Botanic Garden of *M. splendens*, Linn.

M. stipulaceum, Thunb., "Fl. Cap.," p. 422.

One sheet so named, containing a specimen from Upsala Botanic Garden, which is *M. falcatum*, Linn., and not at all like *M. stipulaceum*, Linn.

M. subincanum, Haw., in "Phil. Mag.," Vol. LXIV, p. 427 (1824).

Two sheets thus named by Sonder, and apparently correctly. The specimens on one sheet match those of Zeyher No. 2604.

M. tenuifolium, Thunb., "Fl. Cap.," p. 421.

Four sheets, marked 1, 2, 3, and 4 (but the 4 is written very like a 2). No indication of locality is given.

Sheet 1 contains two specimens of a new species allied to *M. gracile*, Haw., but is more slender, and is identical with Rogers No. 16826 (from near Clanwilliam), and Stephens No. 5170 (from Boontjes River).

Sheet 2 contains a specimen of *M. polyanthou*, Haw., "Synop.," p. 270 (1812), of which *M. glomeratum*, Salm Dyck, "Mesemb.," § 48, fig. 1 (not of Linnaeus) is a synonym. It is quite distinct from the plant wrongly figured by Salm Dyck as being *M. polyanthou*, Salm Dyck's plant being *M. violaceum*, D. C.

Sheet 3 contains a specimen of *M. productum*, Haw.

Sheet 4 contains two species. The right-hand specimen is *M. productum*, Haw., in "Phil. Mag.," Vol. LXIV, p. 425 (1824). The left-hand specimen may possibly be a weak branch of *M. tenuifolium*, Linn., but is not determinable with any certainty.

M. testiculare, Thunb., "Fl. Cap.," p. 412.

One sheet, containing two specimens in fruit of *M. testiculare*, Ait., "Hort. Kew.," ed. 1, Vol. II, p. 181 (1789); Thunb. in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 6 (1791).

Collected in the Karroo between Olifants River and the Bokkeveld Mountains.

M. tetragonum, Thunb., "Fl. Cap.," p. 426.

One specimen collected at Hantam, the type of **M. tetragonum**, Thunb., "Prodr.," p. 91 (1800). Thunberg's specimen is the same as Zeyher No. 702 and Pearson Nos. 3066, 3391, and 3688, which have incorrectly been distributed under the name of *M. defoliatum*, Haw. That species, however, differs in habit, has less connate leaves and shorter pedicels. Possibly a plant collected by Pearson at Plaatklip, Namaqualand, and distributed under the same name without a number, may be the true *M. defoliatum*, Haw. (*M. clavatum*, Jacq., not of Haw.).

Thunberg describes the stems of *M. tetragonum* as "subtetragonous," but his specimen has terete stems with a slight compression at the nodes, where alone they can be called subtetragonous, and they are in no way different from the stems of the specimens of Zeyher and Pearson.

Sonder in "Fl. Cap.," Vol. II, p. 432, erroneously quotes *M. fasciculatum*, Thunb., as a synonym of *M. tetragonum*, but the two species are totally different, as I have noted under *M. fasciculatum*, Thunb.

N.B.—As Moench, two years later, in his "Methodus Plantas, Supplementum," p. 191 (1802), also published a species under the name of *M. tetragonum*, which has been omitted by modern monographers. I would like to point out to future workers upon this genus, that it is probably a synonym of some species described by Haworth. The description of *M. tetragonum*, Moench, is as follows:—

Stem erect, shrubby, glabrous, smooth greyish; branches tetragonous. Leaves decussate, distinct (i.e. not united at the base), triquetrous, keeled, acute, not dotted, with the margins and keel reddish. Flowers solitary subsessile. Calyx 5-lobed; lobes ovate, three with membranous tips. Corolla reddish, with subulate petals.

M. tortuosum, Thunb., "Fl. Cap.," p. 427.

Two sheets marked α and β . Collected in the Karroo.

Sheet α contains two specimens of *M. tortuosum*, Linn., which agree with those of Pearson No. 3057 (from the valley of Kamsob).

Sheet β contains four pieces of *M. expansum*, Linn.

M. trichotomum, Thunb., "Fl. Cap.," p. 419.

One specimen, collected between the Olifants River and the Bokkeveld Mountains, the type of *M. trichotomum*, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 14 (1791).

This dwarf shrubby species continuously branches in a trichotomous manner from the base, with internodes 5–12 lines long, greyish, not dotted with white. The ultimate branches are only $1\frac{1}{4}$ –2 inches long, not 2–3 inches as stated by Sonder. Leaves opposite, 2–5 lines long and apparently about 1 line broad and 1 line thick, flat above, slightly keeled at the apical part on the back (obsoletely trigonous, *ex* Thunberg), obtuse, firm, smooth (not papillate), green. Flowers terminal, solitary, sessile. Calyx 4-lobed; lobes unequal, the larger pair leaf-like up to 3 lines (or perhaps more) long, and twice as long as the shorter pair, which are ovate, obtuse, membranous, keeled on the back. Corolla with the outer petals spreading, dark purple, the inner short, whitish. Styles 4, very short, erect, purple; stigmas simple, spreading, yellow.

The above is Thunberg's description with particulars from the specimen included. The floral structure cannot be seen without damaging the specimen, as there are only two flowers upon it. I fail to match this species with any specimen at Kew, but Pearson No. 5549 is evidently allied to it, differing by its foliage, white-dotted branches, and pedicellate flowers.

M. tripolium, Thunb., "Fl. Cap.," p. 417.

One specimen from Upsala Botanic Garden, which is *M. lanceum*, Thunb., not *M. tripolium*, Linn. (See under *M. lanceum*.)

M. truncatum, Thunb., "Fl. Cap.," p. 412.

One sheet, containing two specimens, collected among rocks in the Kamanassie Karroo in all probability, as he states that it flowers in January and February, and he was in that region in January, 1774. But Thunberg also states that it grows in the Karroo below the Bokkeveld, and near Hex River. As he only collected in these localities in October and November, 1773 and 1774, the Kamanassie Karroo is the more probable locality for it, and I have no doubt he mistook other species of this group for *M. truncatum*, for many of them are very similar in appearance. This plant is—

Conophytum truncatum, N.E.Br., in the "Gardeners' Chronicle" (1922). Vol. LXXI, p. 261.

Mesembryanthemum truncatum, Thunb., in "Nov. Act. Acad. Leop.-Car. Ephem.," Vol. VIII, Append., p. 5 (1791).

Plant forming dense tufts, composed of closely packed branches $\frac{1}{2}$ –1 $\frac{1}{2}$ inch long and about 3 lines thick, including the numerous wrinkled, slightly shining, brown sheaths of a firm or somewhat parchment-like (not membranous) texture with which they are covered. Growths about 4 lines long and 3–4 lines in diameter across the top, which Thunberg describes as "retuse-truncate." The dried growths are glabrous, smooth, slightly shining, of a tan-brown colour, and conspicuously dotted. The specimens seem to have the remains of either some very damaged flowers or young fruits that are damaged and not in a fit state to examine, and also one perfect capsule $2\frac{1}{2}$ lines in diameter, with 5 valves. Thunberg describes the calyx as 4-lobed, but in this specimen it was doubtless 5-lobed, although the lobes have disappeared. Yet it is not uncommon with species of this genus for flowers on the same plant to vary in the number of the calyx-lobes from 4–5.

This species seems never to have been found by any other collector, and has never been in cultivation.

M. tuberosum, Thunb., "Fl. Cap.," p. 421.

This species is now missing from Thunberg's Herbarium. But as, according to the description, it is quite a different plant from that to which Linnaeus had long before applied the same name. I propose for it the following change of name, and add a translation of Thunberg's description :—

M. stratum, N.E.Br.

M. tuberosum, Thunb., "Prodr.," p. 90 (1800), and "Fl. Cap.," p. 421, not of Linnaeus.

Stem prostrate, 9 inches or more long, filiform, terete, branching, glabrous, greyish. Branches opposite, somewhat secund, decumbent, about 3 inches long, papulose, green. Leaves opposite, united and sheathing at the base, about half as long as a finger-nail, terete-subulate, erect, papulose. Flowers terminating the branchlets, pedunculate, red. Peduncle thickened (upwards?), about as long as the leaves, 1-flowered, papulose. Calyx 5-lobed, papulose: lobes lanceolate, somewhat obtuse, erect. Petals linear, obtuse.

Grows on dry plains near Downes (Daunis), in the Calvinia Division.

Thunberg makes no mention of the plant having a tuberous root-stock or producing tubers, so that his name "*tuberosum*" seems inapplicable, unless he intended it to mean that because the plant was papillate on all green parts it was full of small swellings (from *tuber*, a swelling).

M. umbellatum, Thunb., "Fl. Cap.," p. 414.

There are six sheets bearing this name, marked *a*, *β*, *γ*, *δ*, *ε*, and *ζ*. Thunberg describes three plants as varieties under this name, but I am unable to fit his specimen to the descriptions with certainty.

Sheet *a* contains one specimen of a species I am not able to identify. It is identical with Bolus No. 9009 (from near Clanwilliam).

Sheet *β* contains three specimens: the right-hand specimen, which I have marked "B" on the sheet, is *M. parviflorum*, Jacq., "Hort. Schoenbr.," Vol. III, p. 15, t. 278 (1798). The other two specimens, marked "A," I do not match with any species at Kew.

Sheet *γ* contains two specimens of what is probably an undescribed species, which is the same as Pearson No. 5181 (from the southern slopes of Pikinier's Pass).

Sheets *δ* and *ε* contain specimens that I believe to belong to *M. tuberosum*, Linn.

Sheet *ζ* contains two specimens of a new species allied to *M. arboriforme*, Burch., but has shorter and stouter leaves, and the branches of the cyme diverge at a smaller angle, but it is scarcely in good enough condition to describe from.

Neither of the specimens belong to *M. umbellatum*, Linn., and I am unable to say to which of the localities he gives the various specimens belong. He states that they grew in the Karroo between Olifants River and the Bokkeveld Berg, and in the Karroo beyond Hartequas Kloof.

M. uncinatum, Thunb., "Fl. Cap.," p. 420.

One sheet, containing four specimens, three of which belong to *M. uncinatum*, Linn., "Sp. Pl.," ed. 1, p. 483 (1753); *M. uncinellum*, Haw., "Rev.," p. 125 (1821).

These specimens are identical with Rehmann No. 2844 (from Verkeerde Vley, in the Ceres Division).

The fourth specimen is a scrap of *M. villosum*, L. (*M. Ecklonis*, Salm Dyck, "Mesemb.," § 49, fig. 5), evidently accidentally mixed with the other specimens.

Thunberg collected *M. uncinatum* on the Karroo between the Olifants River and the Bokkeveld Mountains.

M. verruculatum, Thunb., "Fl. Cap.," p. 435.

One sheet, containing three specimens, all belonging to different species, neither of which is *M. verruculatum*, Linn.

Thunberg gives two descriptions, and states that one (description *a*) grows in the Karroo between the Olifants River and the Bokkeveld, and the other (description *β*) at Hantam.

The lower specimen on the sheet, which seems to fit description "*a*," has been marked "*β*" by Sonder, and quoted by him as being *M. veruculoides*, Sond. But it is not the plant described by him under that name, his description having been made from a plant collected by Zeyher. Thunberg's specimen is not determinable.

The middle specimen on the sheet is *M. glaucum*, Linn.

The upper specimen is different from either of the others, and consists of a piece of a stout, prostrate, leafy, purple stem without flowers. It is identical with Wolley Dod's specimens Nos. 2420 and 3663 (from Lion's Head, near Capetown), which I have not identified.

Besides these enumerated above, Thunberg's Herbarium contains four sheets that are unnamed by Thunberg, one of them being named by Sonder. The following is my identification of these :—

M. caninum, Lam., "Encycl.," Vol. II, p. 487 (1786).

One sheet, labelled on the back as from Upsala Botanic Garden.

M. noctiflorum, Linn.

One specimen, labelled "*Caroo infra Bockland.*"

A specimen that is without information on the sheet, may possibly be *M. spectabile*, Haw., "Obs.," p. 385 (1795).

M. villosum, Linn., "Sp. Pl.," ed. 1, p. 483 (1753).

M. Ecklonis, Salm Dyck, "Mesemb.," § 49, fig. 5 (after 1836).

One specimen, stated on the back of the sheet to have been collected at the Cape of Good Hope.

It is correctly named *M. Ecklonis* by Sonder, which so exactly agrees with the original description of *M. villosum* under the name of "*Mesembryanthemum caule folisque pubescentibus*" in Linnaeus. "*Hortus Cliffortianus*," p. 216, as quoted in his "*Species Plantarum*," that I have no doubt whatever of its identity with that plant, although there is no specimen of *M. villosum* in the Linnean Herbarium.

	PAGE		PAGE
<i>Aizoon hispanicum</i> , Linn.....	146, 159	<i>Mesembryanthemum criniforme</i> , Linn. f.,	
<i>microphyllum</i> , Bartl.....	155	Thunb.....	152, 157, 159
<i>Conophytum truncatum</i> , N.E.Br.....	165	<i>crystallinum</i> , Linn.,	
<i>Galenia filiformis</i> , N.E.Br.....	155	Thunb.....	152
<i>herniariaefolia</i> , Fenzl.....	155	<i>crystallophanes</i> , Eckl. &	
<i>Glottiphyllum longum</i> , N.E.Br.....	157	Zey.....	145
<i>Mesembryanthemum acinaciforme</i> , Linn....	145, 156	<i>cymbiforme</i> , Haw.....	162
<i>aculeatum</i> , N.E.Br.....	163	<i>debile</i> , Haw.....	—
<i>Aitonis</i> , Jacq.....	145, 146, 160	<i>decussatum</i> , Thunb... 152,	153
<i>albidum</i> , Linn., Thunb... 145		<i>defoliatum</i> , Haw.....	164
<i>anatomicum</i> , Haw.....	154	<i>deltoides</i> , Linn., Thunb..	153
<i>angulatum</i> , Thunb... 145,	160	<i>denticulatum</i> , Haw.....	153
<i>angulatum</i> var. <i>gracile</i> ,		<i>difforme</i> , Linn.....	153
Sond.....	146	<i>difforme</i> , Thunb.....	153
<i>angulatum</i> var. <i>ovatum</i> ,		<i>digitatum</i> , Ait.....	153
Sond.....	160	<i>digitiforme</i> , Thunb.....	153
<i>apetalum</i> , Linn. f.... 146,	159	<i>dissimile</i> , N.E.Br.....	157
<i>arboriforme</i> , Burch.....	166	<i>dolabriforme</i> , Linn.....	153
<i>articulatum</i> , Thunb.. 146,	147, 162	<i>dolabriforme</i> , Thunb... 153	
<i>auratum</i> , Sond.....	148	<i>echinatum</i> , Ait.....	161
<i>aureum</i> , Linn.....	148	<i>Ecklonis</i> , Salm Dyck... 167	
<i>aureum</i> , Thunb.....	148	<i>edule</i> , Linn., Thunb... 154	
<i>barbatum</i> , Berger.....	148	<i>elongatum</i> , Eckl. & Zeyh.	160
<i>barbatum</i> , Haw.....	148	<i>emarcidum</i> , Thunb... 154	
<i>barbatum</i> , Linn.....	148	<i>emarginatum</i> , Linn.,	
<i>barbatum</i> , Thunb.....	148	Thunb.....	154
<i>bellidiflorum</i> , Thunb... 148		<i>erectum</i> , Haw.....	155
<i>"bellidifolium" (?)</i> ,		<i>erigeriflorum</i> , Jacq.....	150
Herb. Thunb.....	148	<i>exiguum</i> , N.E.Br.....	153
<i>"bicolor."</i> Herb. Thunb.	149	<i>expansum</i> , D. C.....	157
<i>bicolorum</i> , Thunb.....	149	<i>expansum</i> , Linn..... 154,	157, 165
<i>brachiatum</i> , Ait.....	152	<i>expansum</i> , Thunb.....	154
<i>bracteatum</i> , Ait..... 149,	162	<i>expersum</i> , N.E.Br.....	158
<i>brevifolium</i> , Ait..... 149,	150, 156	<i>falcatum</i> , Linn.....	164
<i>brevifolium</i> , Haw., and		<i>falcatum</i> , Thunb.....	154
var. β	149	<i>fasciculatum</i> , Thunb.. 154,	165
<i>brevifolium</i> , Sond.....	156	<i>fastigiatum</i> , Haw.....	155
<i>calamiforme</i> , L.....	149	<i>fastigiatum</i> , Thunb.....	155
<i>calcareum</i> , Marl.....	163	<i>felinum</i> , Hill.....	161
<i>calycinum</i> , Haw.....	156	<i>filicaule</i> , Haw.....	152
<i>candens</i> , Haw.....	156	<i>filiforme</i> , Thunb.....	155
<i>caninum</i> , Lam.....	167	<i>floribundum</i> , Haw.....	156
<i>capense</i> , Houtt. ex Thunb.	161	<i>forficatum</i> , Thunb... 145,	156
<i>capillare</i> , Linn. f.... 149,	150	<i>geniculiflorum</i> , Linn.,	
<i>capillare</i> , Thunb... 149,	156	Thunb..... 153,	156
<i>ciliatum</i> , Ait., Thunb... 151		<i>glaucum</i> , Linn., Thunb... 156,	167
<i>clavatum</i> , Haw.....	157	<i>gromeratum</i> , Salm Dyck.	—
<i>clavatum</i> , Jacq.....	164	<i>gracile</i> , Haw.....	164
<i>coccineum</i> , Haw.....	149	<i>gramineum</i> , Haw.....	157
<i>collinum</i> , Sond.....	158	<i>gravidicaule</i> , Haw.....	147
<i>copticum</i> , Jacq.....	146	<i>haeckelianum</i> , Berger... 160	
<i>copticum</i> , Linn.....	159	<i>hispidum</i> , Linn.....	157
<i>corallinum</i> , Thunb.....	151	<i>hispidum</i> , Thunb.....	156
<i>cordifolium</i> , Linn. f.,		<i>hispidum (?)</i> , Hb. Thunb.	156
Thunb.....	152	<i>intonsum</i> , Haw.....	148
<i>crassifolium</i> , Linn... 154,	158	<i>juncum</i> , Haw.....	147
<i>crassifolium</i> , Thunb... 152		<i>lucrum</i> , Haw.....	154

	PAGE
<i>Mesembryanthemum laeve</i> , Thunb.....	157
.. <i>lanceum</i> , Thunb.....	157, 159
	165
.. <i>lateriflorum</i> , D. C.....	—
.. <i>limpidum</i> , Ait.....	152
.. <i>lineare</i> , Thunb.....	157
.. <i>linguaeforme</i> , Thunb.....	157
.. <i>linguiforme</i> , Linn.....	157
.. <i>longum</i> , Haw.....	157
.. <i>loreum</i> , Linn.....	158
.. <i>loreum</i> , Thunb.....	158
.. <i>micans</i> , Linn., Thunb.....	156, 158
.. <i>moniliforme</i> , Thunb.....	158
.. <i>murinum</i> , Haw.....	161
.. <i>mustellinum</i> , Haw.....	161
.. <i>mutabile</i> , Haw.....	149
.. <i>namaquense</i> , Sond.....	153
.. <i>noctiflorum</i> , Linn., Thunb.....	158,
	167
.. <i>nodiflorum</i> , Linn., Thunb.....	159
.. <i>ovatum</i> , Thunb.....	157, 159
	160
.. <i>papuliferum</i> , D. C.....	155
.. <i>papulosum</i> , Linn. f....	146, 160
.. <i>parviflorum</i> , Jacq.....	166
.. <i>pinnatifidum</i> , Linn. f.,	
Thunb.....	160
.. <i>pinnatum</i> , Thunb.....	160
.. <i>plenifolium</i> , N.E.Br....	155
.. <i>polyanthon</i> , Haw.....	164
.. <i>polyanthon</i> , Salm Dyck.....	164
.. <i>pomeridianum</i> , Linn.,	
Thunb.....	152, 161
.. <i>praecultum</i> , N.E.Br.....	156
.. <i>productum</i> , Haw.....	164
.. <i>pruinatum</i> , Thunb.....	161
.. <i>pugioniforme</i> , Houtt.....	152, 161
.. <i>pugioniforme</i> , Linn.....	161
.. <i>pugioniforme</i> , Thunb.....	161
.. <i>pyropaeum</i> , Haw.....	152, 157
.. <i>reptans</i> , Ait.....	152
.. <i>ringens</i> , Thunb.....	161
.. <i>roseum</i> , Willd.....	156
.. <i>rostratum</i> , Linn.....	153, 161
.. <i>sabulosum</i> , Thunb.....	161
.. <i>scabrum</i> , Linn.....	162

	PAGE
<i>Mesembryanthemum scabrum</i> , Thunb.....	162
.. <i>secundum</i> , Thunb.....	147, 162
.. <i>serratum</i> , Linn.....	162
.. <i>serratum</i> , Thunb.....	162
.. <i>sessile</i> , Thunb.....	150, 162
.. <i>spathulatum</i> , Thunb.....	162
.. <i>spathulatum</i> , Willd....	152, 162
.. <i>spathulifolium</i> , Berger....	—
.. <i>spectabile</i> , Haw.....	167
.. <i>spinosum</i> , Linn.....	163, 164
.. <i>spinosum</i> , Thunb.....	163
.. <i>splendens</i> , Linn., Thunb.....	164
.. <i>stelligerum</i> , Haw.....	148
.. <i>stipulaceum</i> , Linn.....	164
.. <i>stipulaceum</i> , Thunb.....	164
.. <i>stratum</i> , N.E.Br.....	166
.. <i>subglobosum</i> , Haw.....	149
.. <i>subincanum</i> , Haw.....	164
.. <i>tenuifolium</i> , Linn.....	162, 164
.. <i>tenuifolium</i> , Thunb.....	164
.. <i>testiculare</i> , Ait., Thunb.....	164
.. <i>tetragonum</i> , Moench.....	165
.. <i>tetragonum</i> , Thunb.....	154, 164
.. <i>Thunbergii</i> , Haw.....	157
.. <i>tortuosum</i> , Linn., Thunb.....	165
.. <i>trichotomum</i> , Thunb.....	165
.. <i>tricolor</i> , Sims.....	157
.. <i>tricolor</i> , Willd.....	152, 157
.. <i>tripolium</i> , Linn.....	154, 165
.. <i>tripolium</i> , Thunb.....	157, 165
.. <i>truncatum</i> , Thunb.....	165, 166
.. <i>tuberosum</i> , Linn.....	166
.. <i>tuberosum</i> , Thunb.....	166
.. <i>umbellatum</i> , Linn.....	167
.. <i>umbellatum</i> , Thunb.....	166
.. <i>uncinatum</i> , Linn., Thunb.....	167
.. <i>uncinellum</i> , Haw.....	—
.. <i>varians</i> , Haw.....	157
.. <i>verruculatum</i> , Linn.....	167
.. <i>verruculatum</i> , Thunb.....	167
.. <i>verruculoides</i> , Sond.....	167
.. <i>villosum</i> , Linn.....	167
.. <i>violaceum</i> , D. C.....	164
<i>Trichocyclus</i> , N.E.Br.....	151
.. <i>ciliatus</i> , N.E.Br.....	151

SOUTH AFRICAN GRAMINEAE.

A NEW GENUS AND SEVEN NEW SPECIES.

By SYDNEY M. STENT.

Mosdenia. Genus novo. *Tragus* similis sed glumae emuricatae, subaequales spiculæ solitariae. Folia eciiliata.

Mosdenia waterbergensis, Stent. (Pl. I.)

Gramen perenne, basin decumbente stolonifere; stolones cataphyllibus brevibus dense imbricatis obtecti. *Innovationes* intravaginales, vaginis brevis et foliis distichis. *Culmi* ascendentes 60–90 cm. longi, simplices, vel ad nodes inferiores ramosi; internodii breviter exserti. *Foliorum vaginae* teres, arctae, striatae, glabrae, laeves, laminis pallidiores; *ligulae* brevis 0·5–0·3 mm. membranaceae; *laminae* lineares ad 12 cm. longae. 5 mm. latae, glaucescentes, glabrae. *Spica* cylindrica, densa, spiculis solitarius nodos tomentosos rachis ferentibus; rachis striata, scabris. *Spiculae* 3·75 mm. longae, uniflorae, anguste lineares, subfalcatae. glabrae, praeter basem crassum minute-pilosum; *glumae* subaequales, inferior paulo longior et latiore, navicularis, falcata, rigide membranacea; carina virida prominente. superior idem sed angustior et strictior; *valva* 3 mm. longa, hyalina, 1–3 nervosa; *palea* subaequale. *Stamina* 3. *Styli* distincti, breves; *lodiculæ* truncatae, ovarium subaequant.

Perennial prostrate ascending, *root-stock* stoloniferous, stolons covered with short, dry, densely imbricate cataphylls. *Innovation shoots* intravaginal with short sheaths and distichously-arranged leaves. *Culms* ascending 60–90 cm. high, simple or branched from the lower nodes, internodes shortly exserted, leafy towards the base. *Leaf sheaths* terete, tight, striate, glabrous, paler than the blades, smooth; *ligule* short 0·5–0·3 mm. membranous; *blades* linear, tapering to an acute or sub-acute point, up to 12 cm. long $\times$ 5 mm. broad, bluish-green, glabrous, smooth or scaberulous on the margins, without a distinct midrib, rounded or slightly eared at the base. *Inflorescence* a single dense spike, the sessile spikelets arranged singly on minutely and densely tomentose notches, in whorls or spirals round a common ribbed and scabrid axis, from which they fall entire. *Spikelets* 3·75 mm. long, 1-flowered, glabrous except at the minutely hairy thickened base, narrow linear, subfalcate; *glumes* subequal to equal, lower slightly longer and broader than the upper, boat-shaped, curved, rigidly membranous, with a prominent green scabrid keel; upper like the lower but narrower and straighter; *valve* about 3 mm. long, delicately hyaline, 1-nerved or sub 3-nerved; *pale* nearly equalling the valve. *Stamens* 3. *Ovary* ovate; *style* short, distinct; *lodicules* truncate, about as large as the ovary.

On sand-veld, Mosdene, Naboomspruit, Waterberg, Transvaal, Galpin M. 447; Rooikop, Springbok Flats, Pole Evans.

This grass was first noticed by Mr. Galpin on his farm Mosdene, and later was collected by Dr. Pole Evans on the farm Rooikop. The latter reports the grass as growing plentifully but being, on account of its short herbage, of little agricultural value.



Mosdenia Waterbergensis, Stent.

A.—Spikelet. B.—Lower glume. C.—Upper glume. D.—Valve. E.—Palea.
F.—Pistil and stamens. G.—Grain.

Eragrostis valida, Stent.

Gramen perenne, dense caespitosum: rhizoma obliqua. *Culmi* erecti, glabri, validi, 120–150 cm. longi, internodiis inclusis: nodis paucis. *Foliorum vaginæ* longae, arctae imbricatae, glabrescentes, inferiores 26 cm. longae, dense hirsutae, plerumque purpureae; *laminae* lineares longe tenuiterque, acutatae, ad 60 cm. longae, 7 mm. latae, glabrae, sed post ligulos pilis longis mollibus tectis, planae, rigidae, supra asperulae, infra, laevae; *ligulae* ad marginem dense ciliatum redactae. *Panicula* patula vel contracta ad 30 cm. longa, rhachibus sulcatis minute scaberulis; *rami* subverticillati vel fasciculati, prope basin divise, *ramuli* breves spiculis pedicellis brevibus-appressis. *Spiculae* griseo-olivaceae, lineares, ad 6·25 mm. longae, 2 mm. latae, 5–7 floribus; rhachilla subpersistens tandem disarticulata; *glumae* subaequales, acutae, carinis scabris, superior paulo longior; *valvae* obtusae circa 2 mm. longae; *paleae* truncatae, carinis scabris; *antherae* 3 mm. longae; *caryopsis* oblonga, 0·3 mm. longa, fusca, dorsa concava.

Perennial densely tufted on an oblique rhizome. *Culms* erect, glabrous, simple, stout, 4–5 feet high, and sheathed for almost their entire length, few noded. *Sheaths* long, the lower ones up to 26 cm. long, tight, imbricating, softly hairy with tubercle based hairs or glabrescent; *blades* linear, tapering to a fine point, up to 60 cm. long × 7 mm. broad, glabrous except for the long soft hairs at the base behind the ligule, smooth below, finely scaberulous above, finely ribbed, flat and rather rigid: *ligule* a short densely ciliate rim. *Panicle* open or contracted up to 30 cm. long; *rachis* sulcate, minutely scaberulous, with glandular swellings in the axils of the branches; *branches* subverticillate or fascicled, divided from near the base. *Spikelets* on short pedicels, more or less appressed to the short branchlets, dark-greyish or olive-green, up to 6·25 mm. long × 2 mm. broad, linear, 5–7-flowered, rachilla sub-persistent, ultimately breaking up; *glumes* subequal, acute, scabrid on the keel, lower 2 mm. long, upper slightly longer; *valves* obtuse, 2 mm. or slightly longer, side nerves reaching beyond the middle: *pale* truncate, with broad flaps and scabrid keels; *anthers* 3 mm. long; *grain* oblong, quadrangular, 0·3 mm. long, brown, concave or grooved on the back.


Maritzburg, Natal, *Angus* 2 (*Govt. Herb.* 13862).

Near to *E. curvata* var. *valida*, but a stouter, coarser grass, with long, tight sheaths that enclose the culms, the lower ones very hairy and long, usually of a purplish colour.

Mr. Angus says of this grass: "A big, coarse variety that grows very erect, has a dark purple stem, which is covered thickly with hairs, and is, if anything, better relished by cattle than No. 3 (*E. curvata*) . . . apparently good feeding, as cattle get into good condition whilst feeding on them."

Schizachyrium inclusum, Stent. (Pl. II.)

Gramen annuum. *Culmi* erecti, simplices, circa 75 cm. longi, 5-nodi, glabri, nodis inferioribus brevissimis teretibus vel subcompressis. *Foliorum vaginæ* laxae, carinae glabrae laeves; *ligulae* membranaceae, ad 0·75 mm. longae, minute fimbriatae; *laminae* lineares, base paulo constricti, glabrae, sed post ligulam pilis longis paucis tectis, 6–10 cm. longae, 2·3 cm. latae, longe attenuatae, apicibus obtuses vel subacutis, planae, vel complicatae, plerumque rubescentes. *Panicula* spuria longa, angusta, exigua; *rami* graciles; *spathae* 3·5 cm. longae, angustae rubescentes laminas minutas apicibus gerentes; *pedunculi* communes graciles, clavati. *Racemi* 2-nati, in spathis inclusi, 2·5 cm. longi, ad circa 6 articuli; *articuli* 3·5 mm. longi, aliquantum concavi, cuneati, apici cupuliforme dentato, 1 mm. longo, breviter ciliati, alioqui glabri; *pedicelli* articulis aequales, complanati, lineares, breviter et parce ciliati, apicibus 2-dentatis. *Spicula sessilis* inter articalum pedicellumque a latere admodum compressa, 4 mm. longa, callo 1 mm. longo, barbato; *gluma inferior* chartacea, marginibus hyalinis, dorso convexa, apicem versus bi-carinata et asperula, nervebus intra-carinis 4–5; *gluma superior* cymbiforma acuta, 1-nerves, breviter mucronata, inferne subchartacea, apicem versus membranacea, vel fere hyalina; *valva inferior* linear oblonga, hyalina, vacua, 2–3 mm.



Schizachyrium inclusum, Stent.

A.—One joint of raceme with sessile and pedicelled spikelets.
B.—Reduced pedicelled spikelet and pedicel.

SESSILE SPIKELET. C.—Lower glume. D.—Upper glume. E.—Lower valve. F.—Upper valve. G.—Grain.

longa; *valva superior* admodum biloba, segmentis linearibus obtusis minute ciliatis; *palea* desunt; *caryopsis* linear, acuta, 3 mm. longa. *Spicula pedicellata* gluma brevissima redacta, mucrone 3.5 mm. longo.

Annual. *Culms* erect, simple, 75 cm. high, about 5-noded, glabrous, lower internodes very short, terete, or sub-compressed. *Leaf sheaths* loose, keeled glabrous, smooth; *ligules* membranous, up to 0.75 mm. long, minutely fimbriate; *blades* linear, slightly constricted at the junction of the sheath, scaberulous on the margins, glabrous except for a few long, white, tubercled based hairs at the base behind the ligule, 2-3 mm. broad, 6-10 cm. long, gradually tapering to an obtuse or sub-acute point, flat or folded, usually turning reddish. *False panicle* long, narrow, and scanty. *Spathes* on sub-capillary branches about 2 cm. long, slender scarious, rusty red towards the tips, up to 3.5 cm. long and 2 mm. broad folded, tipped with a minute undeveloped leaf. *Racemes* very slender on short upward clavate peduncles, entirely included or only partially exerted from the spathes, 2.5 cm. long and about 6-jointed, joints 3.5 mm. long, widening upwards, somewhat concave, with a toothed cupuliform appendage 1 mm. long, shortly and sparsely ciliate, otherwise glabrous. *Pedicles* as long as the joints, flattened, linear, 2-toothed at apex, shortly and sparsely ciliate on the margins. *Sessile spikelets* linear, laterally compressed, convex on the back, 4 mm. long, bearded callus 1 mm. long; *glumes* equal, lower chartaceous with thinly membranous margins, rounded on the back, 2-keeled and scaberulous towards the 2-toothed tip, sparsely ciliate on the margins, intra-carinal nerves 4-5; upper boat-shaped, acute, 1-nerved, nerve extending into a short mucro, thinly chartaceous in the lower part becoming thinly membranous to almost hyaline upwards; lower floret reduced to a linear oblong, hyaline, nerveless valve 2.3-3 mm. long; upper floret 8. valve deeply bifid to below the middle, segments linear, obtuse, minutely ciliate, pale wanting, grain linear, acute, 3 mm. long. *Pedicelled spikelets* reduced to one very small glume, terminating in a mucro 3.5 mm. long.

Enterprise, Rhodesia, Eyles 1646.

This species is founded on one specimen sent by Mr. Eyles. The specimen is a good one and quite complete, but it is obvious that more material of the same species might show trifling variations in measurements, etc., and possibly the culms may be branched.

Enneapogon pretoriensis, Stent. (Pl. III.)

Gramen perenne. *Culmi* erecti, simplices, 30-60 cm. longi, glabri vel internodi inferiores hirsuti. *Foliorum vaginæ* supra arctæ, glabræ; *laminæ* angustæ, lineares, plerumque convolutæ, 18 cm. longæ, 2.5 mm. latæ. *Panicula* angusta, ovata; *rami* solitarii longiores ad 15 mm. longi, internodi rachis inferiores ad 14 mm. longi, superiores deminuenti. *Spiculæ* 5 mm. longæ; *glumæ* æquales vel subæquales, breve pilosæ, apicibus truncatis fulvis; *valvæ* inferioræ dorsis vilosæ, ventris glabræ, sed, lobis basin filis patentibus fimbriatæ; *aristæ* plumosæ, 3-3.5 mm. longæ; *paleæ* 4 mm. longæ, hirsutæ.

Perennial. *Culms* fascicled on a short oblique rhizome, erect, simple, wiry and slender, 30-60 cm. high, glabrous or often shortly hairy at and below the lower nodes, 1-2-noded. *Sheaths* tight except the basal ones, glabrous, finely ribbed; *blades* narrow, linear, convolute or flat, finely pubescent above, up to 18 cm. long $\times$ 2.5 mm. broad, finely ribbed, with 1-2 lateral nerves on either side the midrib. *Panicle* contracted, ovate; *branches* scattered, lower about 14 mm. apart, upper increasingly nearer to each other. *Spikelets* crowded, sessile or shortly pedicelled, about 5 mm. long; *glumes* equal to subequal, shortly hairy, greyish, rust coloured at the truncate tips; lower rather broader and 7-9-nerved; upper narrow, 5-nerved, nerves mostly evanescent above; lower valve 1 line long (excluding awns), villous on back from base to near the middle, glabrous on inner surface except for ring of patent hairs at base of awns; awns plumose to beyond middle, 3-3.5 mm. long, subequal, tips bare and scabrid; pale 4 mm. long, shortly and sparsely hairy on the back, villous on the flaps, and ciliate on the keel; anthers 2 mm. long.



Enneapogon Pretoriensis, Stent.

A.—Spikelet. B.—Lower glume. C.—Upper glume. D.—Valve. E.—Palea.
F.—Pistil and stamens with lodicules.

Pretoria, Wonderboom, *Burtt-Davy* 8905; northern slopes of Magaliesberg, *Fouché* 1. (Of little economic value.)

Near to *E. scoparius*, from which it differs chiefly in the laxer and more distantly branched panicle and broader leaves.

Eragrostis Burttidavii, Stent.

Gramen perenne, dense caespitosum, rhizomate repente pluribus dense confertis brevibus innovationibus. *Culmi* geniculati, 8–40 cm. longi, glabri, 1–2 nodi, internodia exserta. *Foliorum vaginæ* glabrae, tenuiter sulcatae, inferior laxae, breves, latae. *Laminae* glabrae, lineares, culmorum, 7 cm. longae, 2 mm. latae, innovationum angustiores et breviores, planae vel convolutae, apicem versus flexuosae. *Panícula* ovata, ad 10 cm. longa, 7 cm. lata; *rami* solitarii vel 2-nati, gracili, flexuosi; *ramulosi* 2–4 spiculis. *Spiculae* 4 mm. longae, 2 mm. latae, 6–9 florae; *gluma* inferior 0.75 mm. longa, obtusa, superior 1.25 mm. longa, subobtusae; *valvae* 1.5 mm. longae, obtusae, scaberrulae, violaceae, apicibus flavibus; *palea* valvam aequans, truncata, marginibus late inflexis.

Perennial, densely caespitose, on a creeping rhizome with many densely crowded, short innovation shoots. *Culms* geniculate, 8–40 cm. high, glabrous, 1–2-noded, internodes exserted. *Sheaths* glabrous, finely ribbed, tight except the lower ones, which are broader, stout and loose; *ligules* a fringe of hairs. *Leaf blades* glabrous, linear, tapering to a fine point, those of the culm 2 mm. $\times$ 7 cm., those of the innovation shoots narrower and shorter, finely ribbed, midrib scarcely broader than the lateral veins, flat or convolute and flexuous in the upper part. *Panicle* ovate, up to 10 $\times$ 7 cm.; *branches* solitary or or 2-nate, undivided from 2–14 mm. above the base; *rachis* scaberulous or smooth, flexuous; *branchlets* 2–4 spikulate; *pedicels* 2–4 mm. long. *Spikelets* linear, lanceolate, about 4 $\times$ 2 mm., 6–9-flowered; *glumes* unequal, lower 0.75 mm. long, obtusae, upper 1.25 mm. subobtusae; *valves* 1.5 mm. long, obtusae, scaberulous, purple, with light yellowish tips; side nerves fine, extending to beyond the middle; *pale* equalling the valve, truncate, scaberulous, flaps broad; *anthers* 3, up to 1 mm. long.


Very near to *E. bicolor*, Nees; but with a shorter and, in comparison, broader panicle and rather broader spikelets.

Bloemhof, Transvaal, *Burtt-Davy* 10506; Great Karasberg, *Pearson in Bolus Herb.* 8480.

Eragrostis Galpinii, Stent. (Pl. IV.)

Gramen perenne. *Culmi* fasciculati, rhizomate repente, erecti, vel geniculato-ascendentes, glabri, ad 90 cm. longi, pluri nodi, nodi inferiores brevi; culmorum bases et rhizomata cataphyllibus brevibus et scariosibus obtectae. *Foliorum vaginæ* laxae, striatae, inferiores carinatae et compressae; *laminae* angustae, lineares, ad 15 cm. longae, 5 mm. latae, apicibus angustis abtusis, planae, vel complicatae, rigidae, glabrae. *Panícula* late oblonga vel pyramidalis, patula, laxa ad 20 cm. lata, 20 cm. longa; *rami* solitarii, inferiores distantes ramulos 1–3 cm. e basi gerentes. *Spiculae* fere secundae ad 23 mm. longae, 2.5 mm. latae, circa 10–30 florae; *rhachilla* flexuosa, primo persistente; *glumae* subaequales, inferiores 1.5 mm., superiores, 2 mm. longae, minute asperulae; *valvae* purpureae, subacutae, omnino minute scabridae; *palea* acuta, hyalina inter carinas rigide-ciliatas; styli distincti; lodiculi truncati; *antherae* 1.3 mm. longae.

Perennial. *Culms* fasciated on a creeping rhizome, erect or geniculate-ascending, simple or branched, glabrous, slender, up to 3 feet high, many-noded, internodes included, or shortly exserted, or the upper most sometimes long exserted, lower short; bases of culms and rhizomes covered with short, scarios scales. *Sheaths* loose, striate, the lower ones sub-compressed and keeled, glabrous except at the often long-bearded mouth; *ligule* a ciliate membranous rim; *blades* narrow, linear, up to 5 mm. broad $\times$ 15 cm. long, tapering to a narrow, rounded point, flat or folded, rather firm, glabrous, smooth below except towards the tips, finely scaberulous above. *Panicle* broadly oblong or pyramidal, spreading, lax up to 20 cm. broad $\times$ 20 cm. long, branches solitary, lower ones rather distant, spreading,



Stella Lowry.

Eragrostis Galpinii, Stent.

A.—Branchlet showing flexuous persistent rachillae. B.—Spikelet. C.—Lower glume.
D.—Upper glume. E.—Valve. F.—Pale. G.—Pistil and stamens with lodicules. H.—Grain.

unbranched for from 1-3 cm. from the base, up to 11 cm. long; *branchlets* spreading, the lowest pair of the lowest branches usually close together sub 2-nate, the upper ones gradually shorter and solitary, less divided and with fewer spikelets and often second and scabrid; quite glabrous, without or with few hairs in the axils. *Spikelets* linear, up to 23 mm. long $\times$ 2.5 mm. wide, and 10-30-flowered, rachilla at first persistent; *glumes* subequal, lower 1.5 mm. long, upper 2 mm. long, 1-nerved, minutely scabrid on nerve; *valves* over 2 mm., purplish, sub-acute, 3-nerved, side nerves long, minutely scabrid all over; *palea* acute; rigidly ciliate on the keels, scabrid, nearly equalling the valve, hyaline between the rigidly ciliate keels; *ovary* oval; *style* distinct; *lodicules* truncate. *Stamens* 3; *anthers* 1.3 mm. long; *grain* ovoid, rounded; embryo small.

Mosdene, Naboomspruit, Waterberg, *Galpin* 456; Warmbaths, *Burt-Davy* 1149; near Nylstroom, *Burt-Davy* 2034; Letaba River, Zoutpansberg, *Junod, Govt. Herb.*, 4673.

Near to *E. gangetica*, from which it differs in the widely spreading panicle, longer branchlets, and fewer spikelets.

***Tristachya pedicellata*, Stent.**

Gramen perenne, caespitosum, rhizomate obliquo. *Culmi* validi, ad 120 cm. longi, glabri, laeves, 1-3 nodi. *Foliorum vaginæ* aliquantum arctæ, striatæ, glabræ, sed inferiores villosulæ; *laminæ* lineares, planæ, 15-23 cm. longæ, 6-8 mm. latæ, glabræ, rigidae, supra asperulæ, infra laeves. *Panicula* contracta, nodi inferiores pilosi; *rami* ad 5 cm. longi, saepe multo minus, superiores saepe nihilo redacti. *Gluma*, inferior, rigide, 3-nervosa, 13 mm. longæ, 4 mm. lata, glabra, laeves, ovata, subacuta; superior, 24 mm. longæ, 4.5 mm. lata, apice obtusa. *Valva* inferior, 7-nervæ, 20 mm. longæ, glabra, obtusa, laevis; *antheræ* 8 mm. longæ; *lodiculæ* 1.3 mm. longæ, 0.8 mm. latæ; *palea* brevior. *Valva* superior, ad 12 mm. longæ, 7-9 nervæ, mucrones 2 loborum lateralium breves; arista media valida, ad 6.5 cm. longæ; *palea* apice acuta.

Compactly caespitose on oblique rhizoms. *Culms* stout, up to 120 cm., glabrous, smooth, 1-3-noded, internodes exerted. *Leaf sheaths* somewhat tight, finely ribbed and glabrous, except the lowest, which are loose and finely and appressedly hairy between the ribs; *blades* linear, tapering to an acute point, 15-23 cm. long $\times$ 6-8 mm. wide, glabrous, rigid, flat, scabrid above and on the cartilaginous margins, smooth below. *Panicle* contracted, scanty; rachis smooth, with oblique, membranous and hairy scales around the lower nodes; *branches* up to 5 cm. long, often much shorter, the upper so much reduced that the triplets of spikelets appear to be sessile on the rachis. *Pedicels* distinct, 2.4 mm. long. *Lower glume* strongly 3- sub 5-nerved, with a few irregular, oblique, transverse veins, 13 mm. long $\times$ 4 mm. wide (flattened out), glabrous, smooth, ovate, subacute; *upper* 3-nerved, 24 mm. long $\times$ 4.5 mm. wide (flattened), narrowing to an obtuse apex, a few irregular oblique veins connecting the nerves in the upper part, glabrous and smooth. *Lower valve* 7-nerved, 3 nerves percurrent, 20 mm. long, glabrous, smooth, obtuse; *anthers* 8 mm. long; *lodicules* 1.3 mm. long $\times$ 0.8 mm. broad; *palea* 14 mm. long, with scabrid keels and bifid tips, lobes acute. Upper floret 12 mm. long, with long-bearded callus 4 mm.; *valve* 12 mm. long, 7- sub 9-nerved in the upper part, hairy, bifid, awned from between the lobes; awn 6.5 cm.; lobes including short mucro 4 mm.; *palea* with acute scabrid keels, very narrow between the keels, 9 mm. long; *stamens* 3; *style* distinct; *ovary* trigonous; *lodicules* lanceolate, 2 mm. long; *caryops* long, narrow, cylindrical, up to 7 mm. long, grooved on the face; *embryo* 2 mm. long.

Near to *Tristachya rehmanni*, from which it differs chiefly in the pedicelled spikelets, the short bristles of the upper valve and in the broader leaves.

Waterberg (Transvaal): Warmbaths, *National Herbarium, Burt-Davy* 1144; Naboomspruit, *Galpin* 428.

A FUNGUS OF ECONOMIC IMPORTANCE ON THE AVOCADO

(*Persea Americana*).

By E. M. DOIDGE.

COMPARATIVELY little is known about the fungous diseases to which the Avocado is subject. In Florida it is said to suffer from certain leaf and fruit spot diseases caused by a *Colletotrichum* sp., and the young growth is attacked by a scab fungus said to be identical with *Cladosporium citri*, but none of these fungi is recorded as attacking the twigs and larger limbs.

During 1921 it was reported from Louis Trichardt that certain Avocado trees were in a very bad state, and that spraying with bordeaux mixture had proved quite ineffectual. The twigs which were sent for examination were covered with minute pustules, the fruiting bodies of an ascomycete, and as none of the diseased wood had been excised previous to spraying it was not surprising that the inroads of the fungus had not been checked. As a consequence the trees had become so badly cankered that it was impossible to remove all the diseased bark without seriously affecting the vitality of the trees, and at this stage it was impossible to arrest the progress of the malady.

SYMPTOMS.

The effect of the fungus on the twigs and limbs of the Avocado is very similar to that produced by *Physalospora cydoniae* on the limbs and twigs of apple trees.

In the earlier stages of infection the bark becomes sunken and discoloured; the diseased area increases in size, and gradually girdle the twig or branch, so causing the death of the terminal portion. Branches, 2-3 inches in diameter, are killed in this way, and extensive cankers are produced on the larger limbs and the trunk. The bark of the cankered areas is dark-coloured and sunken, the boundary between the healthy and diseased tissue being marked by a raised, reddish-brown line. As the bark is destroyed it dries out and cracks horizontally, eventually falling away from the wood, which has become dead and discoloured. Numerous fruiting pustules of the pathogen may be observed scattered over the diseased areas of the bark.

The fruits on the diseased trees are affected, showing a corky growth at the lower end, which may possibly be due to the same fungus (but this fact has not been established, as it was not possible to obtain trees in bearing for inoculation purposes). The affected area is irregular in form and varies in size. On fruits examined it was roughly 10×5 cm. and 4×5 cm. The surface is very much roughened, verona brown (Ridgway) in colour, and deeply cracked, after the fashion of tortoise-shell.

This affection of the fruit is apparently identical with one which occurs in Florida and is mentioned in a paper by H. E. Stevens, published in the "Proceedings of the Florida State Horticultural Society" for 1918. His description is as follows: "Another common type of injury frequently noted on the fruits is referred to as anthracnose by some of the growers. This type of injury is very similar to melanose of citrus fruits in general appearance. It is superficial, and appears in the form of dark reddish-brown, caked masses on the surface of affected fruits. The markings are hard, compact, and the surface is cracked or broken. The injury may cover only a part or the whole surface of the fruit. It makes an unsightly fruit, but apparently does not affect the quality. The disease is apparently caused by a fungus, perhaps a *Gloeosporium* or a closely related species."

MORPHOLOGY.

Perithecia have been found on cankered areas of the twigs of the Avocado pear (*Persea Americana*). They are usually scattered, standing separate from one another. Sometimes, however, they occur in small groups of two to four, and occasionally they have been found so closely crowded as to form a pseudo-stroma, but no true stroma has been observed. They are buried in the cortical tissues, but at maturity the papillate ostiole breaks through to the surface and protrudes slightly.

The form of a typical perithecium is shown in fig. 1, which is a camera-lucida drawing from a medial longitudinal section. The form is subglobose or broadly ellipsoid, 130–190 μ in the horizontal diameter and 100–120 μ in the vertical diameter. The average dimensions are about 170 $\mu \times 100 \mu$.




Fig. 1.

Section through a perithecium of *Physalospora perseae*.

The wall is differentiated into two layers: the thickness of the outer layer is variable being thinner or almost absent at the base, and thickest at the sides near the base, where it is 16–24 μ thick. The pseudo-cells are brown, polygonal, 4–5 μ in diameter, or somewhat flattened. Within this there is an inner layer of delicate, colourless, compressed hyphae of much more uniform thickness.

The papilla is usually central, but may be oblique: it is 80–100 μ long and 30–50 μ in diameter, and its walls show two distinct layers similar to those observed in the sides of the perithecium; the ostiole appears in section as a narrow passage between the walls of the papilla. The asci interspersed with paraphyses fill the cavity of the perithecium, and there are numerous thread-like paraphyses lining the neck and the inside of the papilla.

The asci are abundant, usually clavate, sometimes tending to the cylindrical or curved, $80-100\ \mu \times 18-23\ \mu$, eight-spored; they are slightly thickened round the apex, but no canal or apical pore could be observed. The asci do not stain blue with iodine; they disappear rapidly at maturity, so that one rarely finds an entire ascus containing mature spores; the wall of the ascus seems to break down as soon as the spores are mature.

The ascospores are more or less distichous, hyaline, ellipsoid or subfusoid, tapering abruptly at each end to a blunt apex, $20-21\ \mu \times 8-10\ \mu$. There are two to three large oil drops in each spore which stain pink with Guéguen's triple stain, the rest of the spore stains a deep blue.




Fig. 2.

Asci and spores of *Physalospora persae*.




Fig. 3.

Section through two pycnidia.

Pycnidia are found on some of the stem cankers in the place of perithecia or are interspersed with them. The pycnidia are single or in small groups: some are spherical or sub-spherical with a distinct papilla, that is to say, they are similar to the perithecia; others are conical, spherical or elongated, without a papilla (fig. 3). They are formed under the epidermis or in the cortical tissues, and are erumpent at maturity. The horizontal diameter is $120-130\ \mu$, vertically they measure $70-187\ \mu$, and there is a double wall, the two layers being similar to those described in the perithecia. From the inner wall arises

the spore-bearing layer, which occupies almost all the space not taken up by the ostiole. The cells lining the cavity arch inward, and elongate to form a stalk, from the tip of which the pycnospore is abstricted.

The pycnospores are ellipsoid and colourless, and do not turn brown at any time; they are $18-20\ \mu \times 5-6\ \mu$ in dimensions.

Physalospora Perseae, nov. sp.

Peritheciis sparsis v. gregariis, tectis, ostiolo papillato erumpente, atris, ellipsoideis v. subglobosis, $130-190\ \mu \times 100-120\ \mu$; ascis clavatis, paraphysatis, $80-100\ \mu \times 18-23\ \mu$, octosporis: paraphysibus filiformibus, numerosis, hyalinis: sporidiis distichis, continuis, hyalinis, ellipsoideis v. subfusiformis utrinque obtusis, $20-21\ \mu \times 8-10\ \mu$. Pycnidii subglobosis v. ellipsoides, $120-130\ \mu \times 70-187\ \mu$; sporulis hyalinis ellipsoideis, $18-20\ \mu \times 5-6\ \mu$.

Hab. in foliis *Perseae americanae*, Louis Trichardt, Transvaal, 25.4.21, leg. J. H. Cronwright.

When the spores have been discharged from the pycnidia or perithecia, one frequently finds the cavity occupied by another fungus, which is probably a saprophyte or secondary parasite. This organism produces spherical pycnidia about $90\ \mu$ in diameter, and filled with small, narrow, ellipsoid, fuliginous, two-celled spores.

The Fungus in Culture.

The fungus was isolated a number of times from diseased twigs by plating (1) ascospores and (2) pycnidia in agar, and (3) from the mycelium present in small pieces of diseased wood. In each case the same fungus was obtained, and the cultural characters were identical. It grows readily on most of the culture media in common use, but as the fruiting bodies develop slowly it was found advisable to use conical flasks rather than petri dishes for culture work, as the latter dried out too rapidly.

Cultures on prune agar, oatmeal agar, and maize-meal agar were kept in the greenhouse in a bright diffused light. The mycelium is at first cottony, but after a few days the submerged hyphae are green or blue-green for some time, after which the growth is dark or nearly black. The outer ends of the aerial hyphae maintain their original cottony appearance and become greyish in colour. Cultures in active growth often show distinct zoning. Thus far the fungus *Physalospora Perseae* closely resembles in its cultural characters *Ph. Cydoniae*, which is parasitic in apple twigs,* but the formation of pycnidia is very different from that observed in cultures of the latter fungus.

The submerged hyphae, as seen by transmitted light, are olive-green or brownish in colour, and are about $3\ \mu$ thick; the aerial hyphae are more slender and hyaline. After three or four weeks on oatmeal agar small bodies consisting of tangled masses of hyphae appear on the surface of the culture; these increase in size until they form tussocks, 3-6 mm. in diameter and about 3 mm. in height; these are dark-coloured bodies, but with a thin covering of the cottony aerial mycelium. After about six weeks the top of those bodies becomes covered with hard, black pustules, from which seven to eight weeks old cultures masses of conidia ooze out.

A cross-section through one of the sclerotial bodies first observed in a three weeks old culture shows that it consists of tangled masses of hyphae similar in form to the submerged hyphae of the ordinary mycelium. Later there is a differentiation with a sterile base, consisting of interlaced dark-coloured hyphae, brown or olive-green, by transmitted light, and an upper portion of more closely woven, lighter-coloured hyphae in which a number of cavities are developed. These are irregularly arranged in one or more layers, ellipsoid or flask-shaped—usually papillate, $250-320\ \mu \times 130-200\ \mu$. The inside of the cavity is lined with a layer of compressed hyaline hyphae, which eventually arch inwards and form the conidiophores. The tip of the conidiophore swells up and the mature conidium is cut off.

* Hesler, L. R.: Black Rot, Leaf Spot, and Canker of Pomaceous Fruits. Cornell Univ. Agric. Exp. Sta. Bull. 379. Aug. 1916.

On maize-meal agar and prune agar slight differences are observed. On maize-meal agar the pycnidial cushions are up to 8 mm. in diameter; on prune agar the form is somewhat different, the pycnidial cushions are columnar, 5-6 mm. high and 3-5 mm. thick. They have a sterile base, which is usually rather smaller in diameter than the rounded head in which the pycnidial cavities are found.

Sugar agar appears to be an unsuitable medium; there is no aerial mycelium, and only a few small sterile sclerotia are formed. Potato agar is also an unfavourable medium. On solution N. agar only a few small sub-spherical cushions are formed, which are 2-3 mm. in diameter, and show some internal differentiation, but none were observed in which mature pycnidial chambers were formed.

The masses of conidia oozing out of the pycnidia are pale orange-yellow (Ridgway). Individual conidia are hyaline, thin-walled, with granular contents, and containing a few oil drops. The majority are ellipsoid in form, a few are sub-fusoid or sub-pyriform; they vary from $16.5-23.5 \mu$ in length and from $6-7.5 \mu$ in breadth, the greater number measuring $18-20 \mu \times 6.5-6.7 \mu$.



Fig. 4.

Germinating conidia after 6 hours at 25°C . in hanging drop.

A number of hanging drop cultures in Ward tubes, each containing a small number of conidia, were put into an incubator at 25°C . After six hours a considerable number had germinated (fig. 4). The form and character of the spore remained unchanged, and a single germ tube, $2.5-3 \mu$ thick, colourless and non-septate, had been produced at or near one pole.

After twenty-four hours possibly 20 per cent. of the conidia remained unchanged and had not germinated. The remainder had lost their granular appearance and had become slightly broader and more broadly rounded at the ends, the majority being $6.7-7.5 \mu$ broad. They were still quite hyaline, but had become septate: most commonly with 2-transverse septa, but a few were 1-septate, and there was an occasional one which was 3-septate. Those which were germinating had, with a few exceptions, produced two germ tubes, one from each terminal cell (fig. 5).

The conidia, which had germinated in six hours, had developed considerably, and had produced elongated hyphae which were septate and branched.

After forty-two hours there was no further change, except in the increased length of the germ tubes and the increased number of spores with two and three septations. At no time was any suggestion of a change of colour observable in the conidia.

The ascus stage was not found in culture, but, as stated elsewhere, its connection with the conidial stage has been traced.

Inoculation Experiments.

Several seedling Avocados in the greenhouse were inoculated with conidia from cultures on oatmeal agar and with pieces of mycelium. In each case a small incision was made in the bark, the infective material inserted, and the branch at the point of inoculation was kept moist for about forty-eight hours.




Fig. 5.
Conidia after 24 hours at 25° C.

The plants which were inoculated with conidia soon showed signs of infection. The tissues around the incision became discoloured and sunken; in some cases the affected area slowly increased in size for a few weeks until it was about 1 cm. in diameter, and then its progress was arrested. Two trees became completely girdled, the whole stem was invaded and killed, and after six months typical conidia were produced. The pycnidia and conidia on these artificially infected trees were identical with those found on twigs in the orchard.

PATHOLOGICAL HISTOLOGY.

The fungus appears to be a wound parasite. In the cortical tissues of infected twigs there is a rather coarse brownish mycelium about $3.5\ \mu$ thick. The cell walls are discoloured, but there is no very prominent deposit in any of the cells. At certain spots there are masses of densely woven hyphae in the epidermal and hypodermal cells, which rupture the cuticle,




Fig. 6.

Section through twig 6 months after inoculation, showing mycelium in cells of cortex.




Fig. 7.

Section through xylem region of same twig, showing mycelium confined to cells of medullary rays.

and are the early stages in the development of the pycnidia and perithecia. The mycelium is intracellular, it penetrates through the cortical cells to the cambium, and invades the xylem region, where it is apparently confined to the cells of the medullary rays. In a large number of sections examined no hyphae were observed in the xylem vessels.

PREVENTIVE MEASURES.

In its early stages the disease could probably be arrested by surgical methods and spraying with bordeaux mixture. As in dealing with other cankers all diseased twigs should be removed, cankers on larger limbs excised, and all wounds painted with some disinfectant and protective substance. In its later stages it is difficult and almost impossible to arrest the progress of the disease.



SPECIES OF ELEPHANTORRHIZA IN THE SOUTH AFRICAN HERBARIA.

By E. P. PHILLIPS, M.A., D.Sc., F.L.S.

The genus *Elephantorrhiza* is a small genus belonging to the *Leguminosae*, and was founded by Bentham in 1842 on the common Transvaal plant, *E. Burchellii*. Four years later he described another species also from the Transvaal, *E. Burkei*. Since then Schinz and Harms each described a species found in South Africa, and two years ago Mr. Burtt-Davy described a species from the Transvaal. An examination of the material in the South African Herbaria has brought to light two additional species which hitherto have been confused with *E. Burchellii*.

Two of the seven species found in South Africa have an acaulescent habit, viz., *E. Burchellii* and *E. obliqua*.

The shape and venation of the pinnules differ in all the species, and afford a good character whereby the species may be distinguished. The reader is referred to the accompanying plate illustrating the pinnules of the seven species described.


Elephantorrhiza Burchellii, commonly known as "Eland's Boontjes," is an extremely common plant on the high veld in spring, and in many parts of the Transvaal is the dominant species on veld which has been burnt. It is an acaulescent plant with enormous roots, which extend to a considerable depth into the ground. Young plants have been excavated by the writer with roots which penetrate to a depth of over 6 feet. The roots of this species have been used for tanning, but other plants have now superseded it. Mr. C. Howlett, of the Division of Botany, carried out a series of dyeing experiments with a 30 per cent. solution of the roots. Wool was treated with a 1 per cent. of the following mordants, and gave the following results (the colours mentioned are those of Ridgway's Colour Standards and Nomenclature):—

1. Ammonium molybdate.....Orange rufous. (Plate II.)
2. Ammonium vanadate.....Plumbeous black. (Plate LII.)
3. Chromium chloride.....Naphthalene yellow. (Plate XVI.)
4. Ferrous sulphate.....Quaker drab. (Plate LI.)
5. Nickel sulphate.....Salmon colour. (Plate XIV.)
6. Stannous chloride.....Empire yellow. (Plate IV.)
7. Sodium tungstate.....Salmon colour. (Plate XLV.)
8. Uranium nitrate.....Rufous. (Plate XIV.)
9. Zinc sulphate.....Salmon colour. (Plate XIV.)
10. Cobalt acetate.....Bittersweet pink. (Plate II.)

The above colours are, however, not permanent, and since these experiments were conducted some three years ago some of the colours in the samples of wool have changed.

The seeds of this species show an interesting and uncommon type of germination, which has been described by Miss J. Hofmeyr in the "South African Journal of Natural History," Vol. III, No. 1.

The writer is indebted to the Curators of the various Herbaria for lending him the material in their respective collections, and to Mr. J. Hutchinson, F.L.S., of the Kew Herbarium, for comparing some of the specimens with material in the Kew and British Museum Herbaria.



Del. Stella Gower.

- Fig. 1.—*E. obliqua*, Benth., Tavy, var. *Glabra*, Poill.
 Fig. 2.—*E. Burchellii*, Benth.
 Fig. 3.—*E. pubescens*, Phill.
 Fig. 4.—*E. Burkei*, Benth.
 Fig. 5.—*E. rangeri* Phill.
 Fig. 6.—*E. Woodii*, Phill.
 Fig. 7.—*E. suffruticosa*, Schinz.

KEY TO SPECIES.

Acaulescent plants with leaves and inflorescence arising from large and thickened root-stocks.

Pinnules ovate, prominently oblique at the base..... 1. *obliqua*.

Pinnules linear, not or scarcely oblique at the base..... 2. *Burchellii*.

Shrubs or small trees with a distinct aerial stem.

Branches pubescent.

Inflorescence about 1 cm. in diameter; calyx lobes acute, as long as the tube; pinnules glabrous.....

Woodii var. *pubescens*

Inflorescence about 7 mm. in diameter; calyx lobes obtuse, half as long as the tube; pinnules pubescent.....

3. *pubescens*.

Branches glabrous.

Midrib more or less in the middle of the pinnule; pinnules equal at the base.....

4. *Burkei*.

Midrib approaching one margin of the pinnule; pinnules oblique or manifestly unequal at the base.

Pinnules with a distinct lateral vein from the base and parallel with the margin.....

5. *Rangeri*.

Pinnules with a midrib only.

Leaves 5-10-jugate; branches fluted..... 6. *Woodii*.

Leaves about 20-jugate; branches not fluted..... 7. *suffruticosa*.

1. **E. obliqua**, Burt-Davy, in "Kew Bulletin" (1920).

An acaulescent plant. *Stem* pubescent. *Leaves* bipinnate, 9-20 cm. long, 4-6 jugate; petiole and rachis pubescent; pinnae 3-8 cm. long; pinnules 0.6-1.5 cm. long, 2-6 mm. broad, ovate, very distinctly mucronate at the apex, very unequal at the base, glabrous, with the midrib prominent beneath, approaching one margin of the pinnule and excurrent into the mucro; lateral veins faint. *Inflorescence* axillary, 4-6 mm. long, cylindric. *Flowers* pedicelled. *Pedicels* about 1 mm. long, glabrous, articulated at the middle. *Calyx-tube* 0.75-1.5 mm. long, campanulate, glabrous; lobes 0.5 mm. long or less, ovate, obtuse, or subacute (sometimes one lobe minutely apiculate). *Petals* sometimes slightly connate at the base, 3.5-4 mm. long, 1 mm. broad, oblong-linear, cucullate, and with an inflexed mucro at the apex, glabrous. *Filaments* connate at the base, 6 mm. long, linear, glabrous; anthers 1 mm. long, elliptic, with a caducous stalked gland at the apex. *Ovary* sessile, 5 mm. long, linear in outline, glabrous; style 7 mm. long, cylindric; stigma simple, opening by an apical pore. *Fruit* not seen.

TRANSVAAL.—Wakkerstroom District: Volksrust. *Jenkins in Herb. Transvaal Mus.* 9296.

Var. **glabra**, Phillips (Pl. V, fig. 1; Pl. VI).

Differs from the type in being perfectly glabrous and in the veining of the pinnules being very distinct.

TRANSVAAL.—Middelburg District: Middelburg, *Jenkins in Herb. Transvaal Mus.* 9128; Botsabelo, *Eiselen in National Herb.* 1229.

2. **E. Burchellii**, Bth., in Hook. "Journ. Bot.," IV (1842) 344 (Pl. V, fig. 2; Pl. VI).

An acaulescent plant. *Stem* glabrous. *Leaves* bipinnate, 5.5-22 cm. long, 5-13-jugate; petiole and rachis glabrous; pinnae 2.5-11 cm. long; pinnules 0.3-1.4 cm. long, 0.25-2 mm. broad, subulate, linear to linear-oblong, acute, slightly narrowed and equal at the base, glabrous, with the midrib distinct and approaching one margin of the leaf. *Inflorescence* arising from the woody portion of stem, also axillary, 4-11 mm. long, cylindric. *Flowers* pedicelled. *Pedicels* 0.5-1.5 mm. long, usually articulated at the base, sometimes at the middle. *Calyx-tube* 0.75-1.25 mm. long, campanulate, glabrous; lobes 0.5-1 mm. long, ovate, acute or obtuse. *Petals* free or slightly connate at the base, 3-4 mm. long, 1 mm. broad, oblong or oblong-lanceolate, acute or subacute, sometimes cucullate at the apex, glabrous. *Filaments* connate at the base, 5.5-6.5 mm. long, linear, glabrous; anthers 0.75-1 mm. long, suborbicular or elliptic, with a caducous stalked gland at the apex. *Ovary* sessile, 3 mm. long, linear in outline, glabrous; style 4-5 mm. long, cylindric; stigma simple, opening by a terminal pore. *Fruit* 8-20.5 cm. long, 2.6-3.8 cm. broad; valves flat or convex.—*Hurv. in "Fl. Cap.,"* II, 277.

SOUTH AFRICA.—Without precise locality, *Mrs. Barber* 55; *Zeyher* 127.

CAPE PROVINCE.—Queenstown District: Zwatkei, December, *E. & Z.* 1693; near Mission Station at Shiloh, 2000 ft., December, *Baur* 379, and in *Herb. Norm. Austro-Afric.* 964; plains round Queenstown, 3600 ft., October, *Galpin* 1917. Kimberley District: Kimberley, *Marloth* 852.

BASUTOLAND.—Leribe District: Near Mission Station at Leribe, *Mrs. Dieterlen* 46, and in *Herb. Musei Austro-Afric.* 7256.

ORANGE FREE STATE.—Ladybrand District: Ladybrand, November, *Rogers in Herb. Transvaal Mus.* 5057; between Senekal and Bethlehem, c. 5000 ft., December, *Flanagan* 1903. Harrismith District: Bester's Vlei, near Harrismith, 5500 ft., December, *Bolus* 8157. Bethlehem District: Bethlehem, *Phillips* 3186. Bloemfontein District: P.O. De Burg, Bloemfontein, February, *Cyrus in Govt. Herb.* 8794. Thabanchu District: Caledon and Vaal Rivers, December, *Zeyher*; Pniel, Vaal River, 4000–5000 ft., November, *Bolus* 2075.

TRANSSVAAL.—Potchefstroom District: Potchefstroom, May, *Burtt-Davy in Govt. Herb.* 52. Heidelberg District: Vereeniging, September, *Leslie in Herb. Transvaal Mus.* 5673; Heidelberg, November, *Miss Leendertz in Herb. Transvaal Mus.* 8094. Bloemhof District: "Kaffraria," Christiana, March, *Native Aaron in Govt. Herb.* 14432. Marico District: Zeerust, January, *Miss Leendertz in Herb. Transvaal Mus.* 11433. Witwatersrand District: Jeppetown Ridge, Johannesburg, c. 6000 ft., October, *Gillfillan* 44A, and in *Herb. Galpin* 6006; Florida, near Johannesburg, December, *Mrs. Hutton* 621. Pretoria District: Groenkloof, near Pretoria, *Phillips* 3051; Pretoria Kopjes, October, *Miss Leendertz* 341, and in *Herb. Transvaal Mus.* 8428, and in *Herb. Austro-Afric.* 1077, October, *Burtt-Davy* 2294; Gezina, near Pretoria, November, *Miss Leendertz* 434, and in *Herb. Transvaal Mus.* 8429. Rustenburg District: Rustenburg, 4500 ft., *Miss Nation* 225; Beestkraal, December, *Jenkins in Herb. Transvaal Mus.* 6963. Waterberg District: Near Nylstroom, November, *Burtt-Davy* 2017; Springbok Flats, 3000 ft., October, *Burtt-Davy in Govt. Herb.* 7066; Pietpotgietersrust, October, *Miss Leendertz in Herb. Transvaal Mus.* 7383; Moorddrift, October, *Miss Leendertz in Herb. Transvaal Mus.* 7364. Wakkerstroom District: Mooifontein, 5500 ft., October, *Schlechter* 3567, and in *Herb. Transvaal Mus.* 5399. Barberton District: Hillsides at Barberton, 2300–2800 ft., September–October, *Galpin* 562; Barberton, November, *Miss Williams in Herb. Transvaal Mus.* 7637, 3000 ft., December, *Thorncroft*, 666, and in *Natal Herb.* 15089.

SWAZILAND.—Without precise locality, *Miss Stewart in Herb. Transvaal Mus.* 8857. Batlapin County: Without precise locality, *Nelson in Herb. Transvaal Mus.* 11740.

KALAHARI.—Without precise locality, *Schinz*. 276. Vryburg District: Armoed's Vlake, November, *Burtt-Davy in Govt. Herb.* 10963.


NATAL.—Mount Moreland, 500 ft., May, Wood 2607; Nonoti, 100–200 ft., October, *Natal Herb.* 14202.

3. *E. pubescens*, Phillips, sp. nov. (Pl. V, fig. 3).

Rami pubescentes. *Folia* 2–10-jugate; foliola 5–8·5 mm. longa, 1 mm. lata, oblongo-linear, apice mucronata, basi obliqua, pubescentia. *Inflorescentia* 7 cm. longa, 7 mm. lata, cylindrica. *Pedicellus* 0·5 mm. longus, glaber. *Tubus calyci* 0·5 mm. longus, campanulatus; lobi 0·25 mm. longi, ovati, apice obtusi.

RHODESIA.—Six miles below Kafue Bridge, *Rogers* 8659.

Branches striate, pubescent. *Leaves* bipinnate, 4–17 cm. long, 2–10-jugate; petiole and rachis pubescent; pinnae 5–7 cm. long; pinnules 5–8·5 mm. long, 1 mm. broad, oblong-linear, shortly mucronate, unequal at the base, pubescent more especially on the under surface, with the midrib distinct and approaching one margin of the pinnule. *Inflorescence* axillary, 7 cm. long, about 7 mm. in diameter, narrowly cylindric. *Flowers*



Elephantorrhiza Burchellii, Benth.

Elephantorrhiza obliqua, Burt-Davy,
var. *glabra*, Phillips.



[Photo]

Field of *Elephantorrhiza Burchellii*, Benth., at
Groenkloof, near Pretoria.

[E. P. Phillips.

pedicelled. *Pedicels* 0.5 mm. long, glabrous. *Calyx-tube* 0.5 mm. long, campanulate, glabrous; lobes 0.25 mm. long, ovate, obtuse, not ribbed. *Corolla*, with a tube, 0.5 mm. long; lobes 2 mm. long, 0.75 mm. broad, oblong, obtuse, slightly inflexed at the apex. *Filaments* connate at the base, 3.5 mm. long, linear, acute at the apex, glabrous; anthers scarcely 0.5 mm. long, subglobose, with a minute caducous gland at the apex. *Ovary* shortly stalked, 1.5 mm. long, linear in outline, glabrous; style 2 mm. long, cylindric; stigma simple.

This species is readily distinguished from the former by the pubescent pinnules and the short, rounded calyx lobes, which are only half as long as the calyx-tube.

4. **E. Burkei**, Benth., in Hook., "Journ. Bot.," V (1846) 81 (Pl. V, fig. 4).

A branched shrub 3-5 ft. high, or small tree 8-10 ft. high. *Stem* and branches glabrous. *Leaves* bipinnate, 7-23 cm. long, 4-6-jugate; petiole and rachis glabrous; pinnæ 3-10 cm. long; pinnules 0.7-1.6 cm. long, 1-3 mm. broad, oblong or linear-oblong, obtuse, or shortly and bluntly mucronate at the apex, sometimes emarginate, slightly narrowed and equal at the base, with the midrib distinct beneath and situated in the middle of the pinnule. *Inflorescence* axillary, 5-10 cm. long, cylindric. *Flowers* pedicelled; pedicels 0.75-2 mm. long, articulated at the middle, more rarely at the base. *Calyx-tube* 1-1.5 mm. long, campanulate, glabrous; lobes 0.5-1 mm. long, ovate, subacute or obtuse, usually (rarely without) with very minute but distinct projections in the sinuses or on the margins of the lobes. *Petals* 2.5-4 mm. long, 1.25-1.5 mm. broad, lanceolate, elliptic-lanceolate or elliptic, cucullate and inflexed at the apex, glabrous. *Filaments* connate at the base, 4-6.5 mm. long, linear, glabrous; anthers 0.75-1 mm. long, elliptic or almost quadrate in outline, with a stalked caducous gland at the apex. *Ovary* sessile, 2-3 mm. long, linear in outline, glabrous; style 2.5-4 mm. long, cylindric; stigma simple, opening by a terminal pore. *Fruit* 7-21 cm. long, 2.8-3.5 cm. broad; valves flat.—*Haro*, in "Fl. Cap.," II, 279; *E. rubescens*, Gibbs in "Journ. Linn. Soc. Bot., XXXVI, 441 (1906).

SOUTH AFRICA.—Without locality (probably Portuguese East Africa), *Junod* 435. Transvaal.—Pretoria District: Kopjes about six miles east of Pretoria, *Phillips* 3040; Wonderboom Poort, near Pretoria, October, *Miss Leendertz* 314, and in *Herb. Transvaal Mus.* 8430; near the "Wonderboom," December, *Burt-Davy* 2663, *Miss Leendertz* 433, and in *Herb. Transvaal Mus.* 8431; Magaliesburg, October, *Zeyher* 560, 321. Waterberg District: Warmbaths, January, *Bolus* 11862; Pietpotgietersrust, September-January, *Miss Leendertz* in *Herb. Transvaal Mus.* 4757, 5581. Bechuanaland.—Mafeking, *Appleyard*.

RHODESIA.—South Rhodesia: Matlopo Hills, 4500 ft., September, *Rogers* 5335, *Galpin* 7081; Zambesi Gorge, 2900 ft., September, *Galpin* 7062.

5. **E. Rangeri**, Harms, in Engl., "Bot. Jahrb.," XLIX, 420 (1913) (Pl. V, fig. 5).

A shrub 4 m. (?) high. *Branches* glabrous. *Leaves* bipinnate, 10-19 cm. long, 5-7-jugate; petiole and rachis glabrous; pinnæ 3-6.5 cm. long; pinnules 4-9 mm. long, 0.5-1.75 cm. broad, linear-oblong, shortly and bluntly mucronate, somewhat narrowed at the base and hardly unequal, with the midrib distinct approaching one margin, and a lateral vein arising from the base and approaching the opposite margin, glabrous. *Inflorescence* axillary, 5.5-6.5 cm. long, cylindric. *Flowers* pedicelled. *Pedicels* 0.75 mm. long, articulated at the base. *Calyx-tube* 1 mm. long, campanulate, glabrous; lobes 0.5 mm. long, broad, oblong-lanceolate, obtuse or subacute, glabrous. *Filaments* connate at the base, 5 mm. long, linear, glabrous; anthers 0.75 mm. long, almost quadrate in outline, with a caducous stalked gland at the apex. *Ovary* sessile, 2 mm. long, linear in outline, glabrous; style 2.5 mm. long, cylindric; stigma simple, opening by a terminal pore. *Fruit* not seen.

SOUTH-WEST AFRICA.—Hohearte (?), December, *Dinter* 2264, and in *Herb. Muses. Austro-Afric.* 4485.

6. **E. Woodii**, Phillips, sp. nov. (Pl. V, fig. 6.)

Rami glabri, striati. *Folia* 5–10-jugata; foliola 2·5–4 mm. longa, 1 mm. lata, oblongo-linear, apice acuta, mucronata, base obliqua, glabra. *Inflorescentia* 8·5–11 cm. longa, cylindrica. *Tubus calyci* 0·75–1·25 mm. longus, campanulatus, glaber; lobi 0·25–5 mm. longi, ovati, apice acuti.

NATAL.—Weenen County: Pieters, near Colenso, 3000–4000 ft., *Wood* 7958, and in *Natal-Gout. Herb.* 8768.

Branches glabrous, fluted. *Leaves* bipinnate, 6–15 cm. long, 5–10-jugate; petiole and rachis glabrous; pinnae 1·8–6 cm. long; pinnules 2·5–4 mm. long, 1 mm. broad, oblong-linear, acute, mucronate, very unequal-sided at the base, with the midrib indistinct or evident and then approaching one margin of the leaf, glabrous. *Inflorescence* axillary, 8·5–11 cm. long, cylindric. *Flowers* pedicelled. *Pedicels* 1·25–1·5 mm. long, articulated at the base. *Calyx-tube* 0·75–1·25 mm. long, campanulate, glabrous; lobes 0·25–5 mm. long, ovate, acute. *Petals* slightly connate at the base, 3·5–4·5 mm. long, 1·25 mm. broad, lanceolate, acute. *Filaments* connate at the base, 3·5–5 mm. long, cylindric, glabrous; anthers 1 mm. long, elliptic, with a caducous stalked gland at the apex. *Ovary* sessile, 2·5 mm. long, linear in outline; style 4·5–5·5 mm. long, cylindric, glabrous; stigma simple, opening by a terminal pore. *Fruit* (only one seen), 9 cm. long, 3·2 cm. broad, subfalcate; valves flat.

Var. **pubescens**, Phillips.

Rami pubescentes.

NATAL.—Near Little Tugela, *Wood* 2867.

This differs from the species in the pubescent branches and in being sparsely pubescent on the petiole and rachis. The calyx-lobes have a distinct rib and are membranous between the ribs. The variety and the species require further examination when further material comes to hand, as it is probable that the former may be raised to specific rank.

E. suffruticosa, Schinz., in "Mem. Herb. Boiss.," No. 1, 117 (Pl. V, fig. 7).

A bush 2–4 m. high. *Branches* smooth, glabrous, with dark bark. *Leaves* bipinnate, 16–24 cm. long, about 20-jugate; petiole and rachis glabrous; pinnae 2–7 mm. long; pinnules 3·5–5 mm. broad, linear, acute or subacute, manifestly unequal-sided at the base, with a distinct midrib approaching one margin of the pinnule, glabrous. *Inflorescence* up to 10 cm. long, cylindric. *Flowers* pedicelled. *Pedicels* 0·25–0·75 mm. long, articulated at the base or apex. *Calyx-tube* 1 mm. long, campanulate, glabrous; lobes 0·5 mm. long, ovate, obtuse. *Petals* 3 mm. long, 1 mm. broad, lanceolate, acute or subacute. *Filaments* connate at the base, 5 mm. long, cylindric, glabrous; anthers 0·75–1 mm. long, elliptic, with a stalked caducous gland at the apex. *Ovary* sessile, 2·5 mm. long, linear in outline, glabrous; style 2·5–3·5 mm. long, cylindric, glabrous; stigma simple opening by an apical gland. *Fruit* 8–26 cm. long, 1·7–2·5 cm. broad, linear, very rarely oblong, valves flat.

RHODESIA.—Umtali District: Odanzi River Valley, *Teague* 291.

SOUTH-WEST AFRICA.—Okahandja, 1300 m., November, *Dinter* 314, February, *Dinter*, 2781; Bull's Mouth Pass on steep slopes, bush 10–15 ft. high, December, *Pearson* in *Percy Sladen Memorial Exped.* 8919, and in *Herb. Musei Austro-Afric.* 11972; Awas Mountains, Pass between Haris and Aub, January, *Pearson* in *Percy Sladen Memorial Exped.* 9658.

CONTENTS.

	PAGE
SOUTH AFRICAN ASCOMYCETES IN THE NATIONAL HERBARIUM ...	195
SOUTH AFRICAN GRAMINEAE	222
INDEX TO VOLUME I	305

SOUTH AFRICAN ASCOMYCETES

IN THE

NATIONAL HERBARIUM.

By ETHEL M. DOIDGE.

PART III.

A LARGE proportion of the fungi described in this paper were collected recently in the George, Knysna, and Humansdorp Districts, where the forests appear to be very rich in species. For assistance in the determination of the host plants I have to thank Mr. Phillips, the Forest Research Officer at Deepwalls, Knysna, and Mr. Fourcade, as well as the members of the staff of the National Herbarium.

The recent monograph by Fitzpatrick on the *Coryneliaceae* rendered necessary a revision of the fungi of this group, and all the South African collections in the Herbarium have been examined and the nomenclature revised accordingly.

I am indebted to Dr. Sydow for the determination of a number of *Phyllachoras* on grasses and for descriptions of a number of new species, and to Dr. Butler, of the Imperial Bureau of Mycology, for a number of references in literature not obtainable locally.

86. *Hysterographium Fraxini* (Pers), De Not., var. *Oleastri*, Desm.

Syll. Fung. II, p. 776.

On twigs of *Olea verrucosa*, Wellington, Cape Province, November, 1910, Doidge [990].

Perithecia scattered on entirely or partially decorticated twigs, superficial, straight, or occasionally slightly curved, ellipsoid, hard, black, rounded at the ends, 0.75–1.5 mm. long, 0.5–0.7 mm. broad; longitudinal fissure smooth, with swollen inflexed lips; asci clavate, rounded at the apex, briefly pedicellate, $170\text{--}210 \times 34\text{--}45 \mu$, eight-spored, paraphysate; paraphyses filiform, flexuous; spores distichous, ellipsoid-oblong, very slightly constricted at the medial septum, muriform with 7–9 transverse and 2–3 longitudinal septa, fuscous, $34\text{--}36 \times 16\text{--}18 \mu$.

87. *Hysterographium spinicolum*, sp. nov.

On thorns of *Acacia spirocarpoides*, Weenen, Natal, 30.7.12 [2542].

Perithecia scattered, oriented with their length parallel with the long axis of the thorn, linear, 0.5–1.5 mm. long, 0.3–0.4 mm. broad, acutely rounded at both ends, black, carbonaceous, lips at first closely connivent, finally open in the middle, forming an elliptic aperture exposing the reddish-brown disk; asci paraphysate, linear-oblong, eight-spored, pedicellate, $130\text{--}140 \times 13\text{--}16.5 \mu$; paraphyses filiform, numerous; spores monostichous or subdistichous, slightly constricted at the medial septum, golden-brown, oblong to ellipsoid, muriform, with 5–6 transverse septa and 1–2 longitudinal septa, $20\text{--}22 \times 8\text{--}10 \mu$.

Hysterographium spinicolum, Doidge, sp. nov.

Perithecia sparsa, lineares, $0.5\text{--}1.5 \times 0.3\text{--}0.4$ mm., utrinque acutiuscula, atra, carbonacea, longitudinaliter rimosa, dein aperta; asci paraphysati, oblongo-lineares, octospori, stipitati, $130\text{--}140 \times 13\text{--}16.5 \mu$; paraphysibus numerosis, linearibus; sporae monostichae v. subdistichae, medio leniter constrictae, luteo-brunneae, oblongae v. ellipticae, muriformae, transverse 5–6 septatae, in longitudinem 1–2 septatae, $20\text{--}22 \times 8\text{--}10 \mu$.

Hab. in spinis *Acaciae spirocarpoides*, Weenen, Natal, 30.7.12 [2542].

88. *Hysterographium Acaciae*, n. sp.

On bark of *Acacia giraffae*, Smitskraal, Boshof District, Orange Free State, 22.9.11, Burtt-Davy [1904].

Perithecia gregarious, at first almost spherical, later somewhat elliptic, up to 0.5 mm. diameter, black, carbonaceous, lips closely connivent; asci oblong or clavate, paraphysate, eight-spored, somewhat thickened round the apex, briefly pedicellate, $100-120 \times 18-21 \mu$; paraphyses filiform; spores distichous, ellipsoid, not constricted, fuscous, muriform, with 6-8 transverse and 1-2 longitudinal septa, $27-30 \times 10 \mu$.

Hysterographium Acaciae, Doidge, n. sp.

Perithecia gregaria, elliptica v. rotundata, atra, carbonacea, 0.5 mm. diametro, labiis arcte conniventibus; asci oblongi v. clavati, paraphysati, octospori, apice parum incrassati, breve stipitati, $100-120 \times 18-21 \mu$; paraphysibus linearibus; sporae distichae ellipticae, non constrictae, fuscae, muriformae, transverse 6-8 septatae, in longitudinem 1-2 septatae, $27-30 \times 10 \mu$.

Hab. in corticis *Acaciae giraffae*, Boshof District, 22.9.11, leg. Burtt-Davy [1904].

89. *Cyclothea Bosciae*, n. sp.

On leaves of *Boscia caffra*, Ebb and Flow, Wilderness, George District, 10.5.23, Doidge [17125].

Forms on the leaves numerous, subcircular, brown spots 2.5 mm. diameter, most conspicuous on the under surface, less sharply defined, but visible on the upper surface and showing concentric zoning: old spots are thin and somewhat translucent. Intramatrical hypostroma very copious, consisting of closely interwoven hyaline hyphae, and penetrating right through the mesophyll. Hyphal strands from the hypostroma grow through the numerous stomata in the lower epidermis and produce the thinly crustaceous, dull-black, superficial ascostromata. Loculi linear, irregular in form and size, curved or bent, closely crowded, and dehiscing by longitudinal slits. Hypothecium thin, colourless; outer wall of loculi radial in structure, consisting of hyphae about 3.5μ thick. Asci paraphysate, not staining blue with iodine, sessile, saccate or cylindrical, eight-spored, not thickened round the apex, $33.3-46.6 \times 10-13.3 \mu$; paraphyses not numerous, filiform, septate. Spores two-celled, hyaline, subdistichous, $11.6-13.3 \times 8.3 \mu$, upper cell slightly broader and more broadly rounded than lower.

Cyclothea Bosciae, Doidge, n. sp.

Stromata superficialia, 2-5 mm. diametro, hypophylla ex hypostromate incolore, subepidermali, copiose per stromata numerosa prorumpente oriunda; loculis linearibus, gregariis, irregularibus: hypothecio tenue incolore; asci paraphysati, saccati v. cylindracei, octospori, $33.3-46.6 \times 10-13.3 \mu$; sporae 1-septatae, hyalinae, $11.6-13.3 \times 8.3 \mu$, loculo supero parum latiore.

Hab. in foliis *Bosciae caffrae*, Wilderness, George District, 10.5.23, leg. Doidge [17125].

90. *Macowaniella myrsinicola*, n. sp.

On leaves of *Myrsine melanophleos*, Greytown, Natal, 16.2.22, Doidge [15529]; Montagu Pass Road, George District, 9.5.23, Doidge [17105].

Conidial stage only: Claridge, Natal, 31.5.15, Doidge [8995]; East London, 24.11.17, Doidge [10908].

Stromata epiphyllous, round, superficial, up to 2 mm. diameter, crowded towards the midrib of the leaf, often in contact with one another, but the outline of each circular stroma is always evident and they do not become merged into a continuous mass. Loculi elliptic, straight, or branched and irregular in form, scattered or in small groups, and sometimes arranged more or less in a circle, but not with their axes on radiating lines, $330-480 \times 175-320 \mu$, $50-70 \mu$ high, dehiscing by a longitudinal slit. Hypothecium delicate, pseudocellular: covering membrane radial in structure. Asci paraphysate, ovate to clavate, eight-spored, thickened round the apex, briefly pedicellate, not staining blue with

iodine, $40-44 \times 20-23 \mu$. Spores fuscous, distichous or subtristichous, two-celled, ellipsoid, rounded at both ends, $17-20 \times 7.5-8.5 \mu$, cells subequal, or upper cell somewhat shorter and broader than the lower. Hypostroma epidermal and subepidermal: there is a distinct layer of colourless mycelium between the epidermis and the palisade cells, with hyphae extending downwards into the latter, but the hypostroma is much less strongly developed than in *Macowaniella congesta*. Superficial mycelium fuscous; hyphae $3-3.5 \mu$ thick, tortuous, branching irregularly and anastomosing, bearing small, dark-brown, one-celled hemispherical hyphopodia, $6-6.5 \times 3 \mu$. Conidiophores erect, dark-brown, $13-16 \times 3.5 \mu$, bearing at their tips conidia similar in form to those of *Asterodothis solaris*: conidia four-celled, brown, cylindrical or club-shaped, straight or slightly curved, somewhat constricted at the septa, $37-44 \times 6.5-7 \mu$.

Macowaniella myrsinicola, Doidge, n. sp.

Stromata ascophora epiphylla, rotundata, usque 2 mm. diametro, hypostromate epidermale et subepidermale oriunda, pluries affixa; loculi elliptici, recti v. furcati, sparsi v. subgregarii, $330-480 \times 175-320 \mu$, $50-70 \mu$ alti, hypothecio tenue; asci paraphysati, ovati v. clavati, octospori, apice incrassati, breviter stipitati, $40-44 \times 20-23 \mu$; sporae fuscae, distichae v. subtristichae, ellipticae, utrinque rotundatae, loculis subaequalibus v. loculo supero parum breviores et latiores, $17-20 \times 7.5-8.5 \mu$; mycelium epiphyllum fuscum, ex hyphis $3-3.5 \mu$ latis, tortuosis, ramosis compositum; hyphopodia parva, brunnea, continua, hemisphaerica, $6-6.5 \times 3 \mu$; conidiis 3-septatis, brunneis, cylindraceis v. clavatis, $37-44 \times 6.5-7 \mu$.

Hab. in foliis *Myrsine melanophleas*, Greytown, Natal, 16.2.22, leg. Doidge [15529].

91. *Polyrhizon Celastris*, n. sp.

On leaves of *Celastrus acuminatus*, Deepwalls, Knysna District, 13.5.23, Doidge [17207].

Epiphyllous, on yellow-brown leaf spots, up to 5 mm. diameter; opposite the centre of each collective stroma there is a yellowish spot visible on the lower side of the leaf; each collective stroma is composed of a large number of single stromata developing in concentric rings. Single stromata crowded, spherical in outline, $300-600 \mu$ in diameter, irregular in height, $30-50 \mu$: outer wall opaque, radiating in structure, consisting at the margin of brown hyphae, $4-5 \mu$ thick. Each stroma has only a few loculi, arranged round a central sterile column which is connected with the dark epidermal hypostroma; the peripheral part of the stroma lies free on the surface of the leaf. Hypothecium pseudocellular, pale fuscous. Asci paraphysate, eight-spored (material rather old and asci not well seen); spores dark-brown, opaque, almost black, two-celled, constricted, smooth, $26-28.5 \times 11.5-13.5 \mu$, upper cell somewhat broader.

Polyrhizon Celastris, Doidge, n. sp.

Stromata epiphylla, orbicularia, usque 5 mm. diametro, e pluribus ascomatibus circinantibus concreta; ascomata partialia pede centrali epidermide innata v. marginem radiatim ex hyphis $4-5 \mu$ crasses contexta, $300-600 \mu$ diametro, $30-50 \mu$ alta, loculi pauci, immersi: asci octospori, paraphysati; sporae atro-brunneae, opacae, 1-septatae, constrictae, leves, $26-28.5 \times 11.5-13.5 \mu$, loculo supero paullo latiore.

Hab. in foliis *Celastris acuminati*, Deepwalls, Knysna District, 13.5.23, leg. Doidge [17207].

92. *Hysterostoma Faureae*, n. sp.

On leaves of *Faurea McNaughtonii*, Gouna Forest, Knysna District, 23.5.23, Phillips [17293].

Stromata epiphyllous, less frequently hypophyllous, numerous, dull-black, minute, $300-400 \mu$ diameter, elevated in the centre into a slight papilla, which eventually breaks away and forms an irregularly circular opening. The stroma is surrounded by a fringe of radiating hyphae, which are brown, $4-6 \mu$ thick, frequently septate, often adhering by their

lateral walls and forming strands of 2–3 hyphae, and bearing a few hemispherical hyphopodia. The radiating hyphae often give rise to smaller secondary stromata, and thus form groups which are up to 2 mm. diameter.

The loculi are solitary or few in each stroma, round or irregular in outline, 70–90 μ high. The covering membrane is composed of radiating hyphae 3–3.5 μ thick. The hypothecium is thin, hyaline, and composed of delicate hyphae: it is connected at many points with the subcuticular hypostroma. Hypostroma hyaline, lying between the cuticle and the epidermis and causing disorganization and discoloration of the contents of the epidermal cells. Asci paraphysate, eight-spored, clavate or narrow ovate, slightly thickened round the apex, 45–50 $\times$ 20–23 μ , not staining blue with iodine. Spores olivaceous-brown, distichous, two-celled, constricted, 20–23 $\times$ 8.5–10 μ ; they appear to be surrounded by a mucilaginous envelope when immature; upper cell broader and more broadly rounded than the lower.

Hysterostoma Faureae, Doidge, n. sp.

Stromata epiphylla, sparsa, atra, minuta, 300–400 μ diametro, ex hypostromate sub, cuticulari oriunda, pluries affixa, periphice in hyphas radiantibus dissoluta. Hyphae brunneae, 4–6 μ crassae, hyphopodiis paucis hemisphericis. Loculi pauci vel solitarii, 70–90 μ alti, hypothecio tenui. Asci paraphysati, octospori, clavati v. ovati, apice lenitee incrassati, 45–50 $\times$ 20–23 μ . Sporae distichae, 1-septatae, constrictae, brunneae 20–23 $\times$ 8.5–10 μ , loculo supero latiore.

93. *Palawaniella Dovyalidis*, n. sp.

On leaves of *Dovyalis rhamnoides*, Ebb and Flow, Wilderness, George District, 15.5.23, Doidge [17117]; Bathurst District, 14.7.19, Doidge [12347].

Ascigerous stromata, epiphyllous, dull-black, opaque, carbonaceous, attached to the leaf at many points, superficial, minute, 190 μ diameter, or up to 360 $\times$ 200 μ , forming groups up to 5 mm. diameter, developing centrifugally, outer wall very opaque, radiating in structure, composed of hyphae about 5 μ thick. Hypostroma well developed in the epidermal cells under each ascostroma, forming dense knots of hyphae in the epidermal cells, which are hyaline for the most part, but fuscous where they are connected by numerous delicate processes with the superficial hypothecium. These knots of hyphae in the epidermal cells are connected by delicate subepidermal strands, which also penetrate between the palisade cells. Loculi 1–2 in each stroma, opening by a round pore or by stellate fissures in the outer wall. Loculi flat, hypothecium thin. Asci four-spored, paraphysate, clavate, sessile or with a short foot, thickened round the apex, 33–40 $\times$ 13.5–16.5 μ . Spores pale fuscous, two-celled, not constricted, clavate, 15–18 $\times$ 5–6.3 μ , upper cell much longer than the lower, lower cell not more than 3.5 μ long. Pycnidia similar to the ascostromata, conidia hyaline, subspherical to ellipsoid, 6.5–13.5 $\times$ 5–6.5 μ .

Palawaniella Dovyalidis, Doidge, n. sp.

Stromatis ascigeris epiphyllis, tota longitudine matrice adnatis, superficialibus, atris, opacis, carbonaceis, sparsis, minutis, rotundatis v. ovalibus, 190 μ diametro v. 360 $\times$ 200 μ , saepe plus minus confluentibus et tunc usque 5 mm. diametro, in medio crassis, marginem v. tenuioribus, radiatim ex hyphis 5 μ crassis contextis; hypostromate epidermide bene, subepidermide parum evoluto; loculis in stromatibus singulis v. duobus, tandem poro rotundo v. stellatim disrumpentibus; hypothecio tenui; ascis 4-sporis, paraphysatis, clavatis, sessilibus v. breve stipitatis, apice incrassatis 33–40 $\times$ 13.5–16.5 μ ; sporis fuscis, clavatis, 1-septatis, haud constrictis, 15–18 $\times$ 5–6.3 μ , loculo supero longiore, infero ca. 3.5 μ longo.

Hab. in foliis *Dovyalidis rhamnoides*, Wilderness, George District, 15.5.23, leg. Doidge [17117].

94. *Asterinella Pterocelastri*, n. sp.

On leaves of *Pterocelastrus variabilis*, var. *tricuspidatus*, Storms River, Humansdorp District, 15.5.23, Doidge [17190]; Deepwalls, Knysna District, 13.5.23, Doidge [17222].

Hypophyllous, forming thin irregularly circular, sooty blotches up to 1 cm. diameter, often confluent and covering larger areas of the leaf; mycelium copious, reticulate, composed of pale fuscous hyphae 3–4 μ thick; hyphae straight or subtorulose, without hyphopodia, branching and anastomosing profusely; thyriothechia fuscous, only slightly darker than the mycelium, flattened-hemispherical, 150–180 μ diameter, composed of radiating hyphae about 3 μ thick, cells 3–3.5 μ long in the centre, longer near the periphery, margin fimbriate, the radiating hyphae running out and anastomosing with the mycelial hyphae; asci aparaphysate, eight-spored, ovate, thin-walled, not staining blue with iodine, 30–37 $\times$ 20–25 μ ; spores conglobate, ellipsoid, two-celled, slightly constricted, pale fuscous, thin-walled, delicate, smooth, 17–20 $\times$ 6.5–8.5 μ , upper cell slightly broader and more broadly rounded than the lower.

Asterinella Pterocelastri, Doidge, n. sp.

Hypophylla, plagulas primitus, orbiculares, usque 1 cm. diametro, sed mox confluyendo majores et irregulares formans; mycelium ex hyphis fusciculis, 3–4 μ latis, rectis v. subtorulosis, ramosis formatum; hyphopodia desunt; thyriothechia orbicularia, 150–180 μ diametro, radiatim ex hyphis 3 μ crassis contexta, fimbriata; asci aparaphysati, octospori, ovati, 30–37 $\times$ 20–25 μ ; sporae conglobatae, ellipsoideae, 1-septatae, leves leniter constrictae, fuscae, 17–20 $\times$ 6.5–8.5 μ , cellula superiore leniter latiore.

Hab. in foliis *Pterocelastri variabilis*, Storms River, Humansdorp District, 15.5.23, leg. Doidge [17190].

95. *Asteromyxa inconspicua*, n. sp.

On leaves of *Chilianthus arboreus*, Van Stadens Pass, Cape Province, 19.5.23, Doidge [17252].

Forming minute, very inconspicuous, black spots on the upper surface of the leaves. Mycelium radiating, composed of fuscous, slender hyphae, without hyphopodia, 3–5 μ thick, septate, branching irregularly and anastomosing, with a tendency to adhere to adjoining hyphae by the lateral walls, and so form radiating strands, 2–4 hyphae in breadth; often producing at the tips of the lateral branches large, oval, brown conidia 16–20 $\times$ 11–13 μ . Thyriothechia few, 1–3 in centre of mycelium, circular, at first flat, shield-shaped, radiating in structure, composed of radiating hyphae about 3 μ thick, later becoming almost hemispherical and more irregular in outline, 160–270 μ diameter. Asci fugaceous, eight-spored, ovate, aparaphysate, sessile, about 30–35 $\times$ 22–24 μ ; interior of thyriothechium filled at maturity with dirty-brown mucilaginous matter in which the groups of spores are embedded, the asci having disappeared. The swelling of this mucilage forces apart the radiating cells of the covering membrane, which disintegrates. Spores elliptic, two-celled, pale fuscous, somewhat constricted, upper cell slightly broader, 17–22 $\times$ 6–8.5 μ .

Asteromyxa, Theiss et Syd. (Ann. Myc., XV, 1917, p. 419), is a monotypic genus, which bears among the ahyphopodiate genera the same relation to *Asterinella* as *Englerulaster* does to *Asterina* among the hyphopodiate genera. It is based on *A. hirtula* (= *Dimeriella hirtula*, Speg) as type, and this species has a number of setae on the thyriothechium. *Asteromyxa inconspicua* has all the characters of the genus *Asteromyxa* except the setae, and is therefore assigned to that genus. The diagnosis of the genus *Asteromyxa* must be broadened to include species whose thyriothechia have no setae, or else a new genus must be established for these.

Asteromyxa inconspicua, Doidge, n. sp.

Epiphylla, plagulas minutas parum perspicuas formans: mycelium laxè ramosum ex hyphis tenuibus fuscis, 3–5 μ crassis compositum; hyphopodia nulla; thyriothechia pauca in quaque plagula, hemisphaerica, 160–270 μ diametro, radiatim ex hyphis ca. 3 μ crassis contexta, mox mucosè dissoluta; asci evanescentes, octospori, ovati, aparaphysati, 30–35 $\times$ 22–24 μ ; sporae ellipticae, 1-septatae, leniter constrictae fuscae leves, 17–22 $\times$ 6–8.5 μ .

Hab. in foliis *Chilianthi arborei*, Van Stadens Pass, Cape Province, 19.5.23, leg. Doidge [17252].

96. *Parasterina reticulata*, n. sp.

On leaves of *Celastrus buxifolius*, Durban, Natal, 1897, Medley Wood (Wood Nos. 6452 and 6458) [9500 and 9518]; East London, 24.11.17, Doidge [10913].

On *Celastrus nemorosus*, Knysna, Cape Province, 3.6.12, Pienaar [2429]; Umgeni, Natal, 16.7.15, Medley Wood [9023]; Van Stadens Pass, Cape Province, 9.3.16, Bottomley [9558]; Alice, Cape Province, 18.11.17, Doidge [10974]; Langholm, Bathurst District, 14.7.18, Doidge [12370]; Howiesons Poort, near Grahamstown, 12.7.19, Doidge [12386]; Van Stadens Pass, 19.5.22, Doidge [17262].

On *Celastrus ruber*, Kentani, Cape Province, 10.4.15, Pegler (Pegler No. 2336) [9069].

On *Celastrus Harveyanus*, Henley, Natal, 24.5.15, Doidge [9000]; Claridge, Natal, 31.5.15, Doidge [8997].



Fig. 1.

Parasterina reticulata.

On *Celastrus pyracanthus*, Pirie Forest, Kingwilliamstown, 8.7.19, Doidge [12330].

On *Celastrus Senegalensis*, Nelspruit, Transvaal, 18.10.13, Van der Bijl [7076]; Rikatli, Portuguese East Africa, September, 1918, Junod [11725].

On *Elaeodendron croceum*, Woodville Forest, George District, 11.11.17, Doidge [10941]; Kentani, Cape Province, 17.4.12, Pegler [2253].

On *Pleurostylia capensis*, Kentani, Cape Province, 20.7.12, Pegler [2532].

On *Putterlickia pyracantha*, East London, 19.7.19, Doidge [12407].

On *Putterlickia verrucosa*, Winkle Spruit, Natal, 28.5.15, Doidge [9007]; Umgeni, Natal, 16.7.15, Medley Wood [9033]; Kentani, Cape Province, 15.12.14, Pegler. (Pegler No. 1967) [8783].

Amphigenous, forming thin, black growths up to 1 cm. diameter, frequently numerous, especially on the upper surface, and coalescing to form a continuous growth covering the leaf surface. Mycelium radiating, composed of brown, fairly straight hyphae about $3.5\ \mu$ thick, the primary hyphae radiating like the spokes of a wheel and giving rise to numerous branches which anastomose and form a continuous open network. The form of the network varies slightly in different specimens: in some the radiating hyphae are more marked, in some the meshes of the network are irregularly angular, in others they tend to be curved. Hyphopodia not very numerous, sometimes rare, most easily seen on the younger hyphae, unilateral or alternate, continuous, hemispherical or obliquely flattened, sometimes sublobed, $3.5-4\ \mu$ high, $8-13\ \mu$ broad. Thyriothecia scattered, fairly numerous more crowded towards the centre of the mycelium, flattened-hemispherical, $220-250\ \mu$ diameter, dehiscing by stellate fissures, the central part later falling away and exposing the asci, dark-brown, opaque, radiating in structure, composed of radiating hyphae about $3\ \mu$ thick; margin fimbriate, radiating hyphae running out and anastomosing with the mycelial hyphae. Asci numerous, paraphysate, eight-spored, ovate or broadly ellipsoid, slightly thickened round the apex, sessile, $40-43 \times 23.5-26\ \mu$. Paraphyses filiform, flexuous; fairly numerous, slightly exceeding the asci. Spores distichous, two-celled, brown ellipsoid, constricted at the septum, smooth, rounded at both ends, $20-23.5 \times 8-9\ \mu$, loculi ellipsoid, upper loculus slightly broader.

Parasterina reticulata, Doidge, n. sp.

Amphigena, plagulas tennes, orbiculares, usque 1 cm. latas, dein confluentes et plus minus effusas formans; mycelium reticulatum ex hyphis longiusculis, ramosis radiantibus, brunneis, ca. $3.5\ \mu$ crassis compositum; hyphopodia haud numerosa, alternantia v. solitaria, sessilia, hemisphaerica v. rarius sublobata, $3.5-4\ \mu$ alta, $8-13\ \mu$ lata; thyriothecia sparsa, sat numerosa, rotundata $220-250\ \mu$ diametro, prinitus stellatim dehiscencia per aetatem late aperta, radiatim ex hyphis, $3\ \mu$ crassis contexta, fimbriata; asci numerosi paraphysati, octospori, ovati v. late elliptici, sessiles $40-43 \times 23.5-26\ \mu$, paraphysibus, filiformis, flexuosis, sat numerosis; spores distichae, 1-septatae, brunneae, ellipticae, constrictae. laeves, utrinque rotundatae, $20-23.5 \times 8-9\ \mu$, loculo supero parum majore.

Hab. in foliis *Celastris Senegalensis*, Rikatli, Portuguese East Africa, 1918, leg. Junod [11725].

97. *Parasterina laxa*, Doidge.

Syn. *Parasterina brachystoma* (Rehm) Th., var. *laxa* (Trans. Roy. Soc. S. Africa VII, p. 245, 1920).

On leaves of *Grumilea capensis*, Woodbush, Zoutpansberg District, 3.8.11, Doidge [1758].

On *Oxyanthus Gervardi*, Berea, Durban, 28.1.18, Van der Bijl [11366].

On *Plectronia obovata*, Deepwalls, Knysna District, 13.5.23, Doidge [17226].

Epiphyllous, forming black, carbonaceous spots up to 10 mm. diameter; mycelium radiating, more or less reticulate, formed of straight, brown hyphae $5-6.5\ \mu$ thick, frequently septate, cells $16-24\ \mu$ long, branching irregularly and anastomosing; hyphopodia alternate or unilateral, continuous, sessile, briefly cylindrical, clavate, hemispherical or pyriform straight or subuncinate, $6.5-13.5\ \mu$ high, $6-13.5\ \mu$ broad; thyriothecia hemispherical, scattered, $200-350\ \mu$ diameter, composed of radiating hyphae about $3\ \mu$ thick; dehiscence at first stellate, later the whole central portion falls away, leaving only the margin with its fimbriate hyphae running out into the mycelium; asci eight-spored, subspherical or broadly ovate, sessile, thickened round the apex, staining golden-brown with iodine, $53-65 \times 40-50\ \mu$; spores conglobate, constricted, brown, 1-septate, minutely verrucose at maturity, $30-33 \times 15-16.5\ \mu$, cells subequal or upper cell broader by $2-3\ \mu$, each cell almost spherical or ovate.

98. *Asterina gerbericola*, n. sp.

On leaves of *Gerbera cordata*, Storms River, Humansdorp District, 15.5.23, Doidge [17175].

Epiphyllous, forming small black spots up to 5 mm. diameter; mycelium delicate, formed of sinuous, brown hyphae 3–3.5 μ thick, branching irregularly and anastomosing, cells 24–27 μ long; hyphopodia distant, not very numerous, alternate, one-celled, sessile, erect or decumbent, variously lobed, lobes rounded, frequently there are three lobes which are themselves sublobed, 6–10 μ high and 8–13 μ broad; thyriothecia numerous, developing at the tips of secondary hyphae, at first fan-shaped, later flattened-hemispherical, 120–130 μ diameter, composed of radiating hyphae 2–2.5 μ thick, margin fimbriate, dehiscence stellate; asci paraphysate, eight-spored, broadly ovate or subspherical, 20–23 $\times$ 23–30 μ , not staining blue with iodine; spores conglobate, two-celled, slightly constricted, dark-brown, opaque, cylindrical, broadly rounded at both ends, verrucose at maturity, 13.5–16.5 $\times$ 7–8.3 μ , cells subequal.

This species closely resembles *Asterina undulata*, from which it differs chiefly in the size and form of the hyphopodia, better developed mycelium, and longer verrucose spores.



Fig. 2.

Asterina gerbericola.

Asterina gerbericola, Doidge, n. sp.

Plagulas epiphyllas, usque 5 mm. diametro atras formans; mycelium tenue, ex hyphis sinuosis, brunneis, ramosis 3–3.5 μ latis, septatis (articulis 24–27 μ longis) compositum; hyphopodia haud numerosa, alternantia, sessilia, varie lobata, 6–10 μ alta, 8–13 μ lata; thyriothecia numerosa in summis hyphis secundariis oriunda, orbicularia 120–130 μ diametro, radiatim ex hyphis 2–2.5 μ crassis contexta, ambitu fimbriata, stellatim dehiscencia; asci paraphysati, octospori, ovato-globosi 20–23 $\times$ 30 μ ; sporae con-

globatae, 1-septatae, cylindraceae, leniter constrictae, utrinque late rotundatae, brunneae, opacae, verrucosae, $13.5-16.5 \times 7-8.3 \mu$, loculis fere aequalibus.

Hab. in foliis *Gerberae cordatae*, Storms River, Humansdorp District, leg. Doidge [17175].

Asterina undulata affinis.

99. ***Asterina Streptocarpi***, n. sp.

On leaves of *Streptocarpus Rexii*, Deepwalls, Knysna District, 13.5.23, Doidge [17223].

Epiphyllous, forming minute black spots up to 3 mm. diameter; mycelium undulate, formed of flexuous hyphae $3.5-5 \mu$ thick, irregularly branched cells $16-20 \mu$ long; hyphopodia one-celled, alternate, fairly numerous, usually erect, plurilobate, with 2-5 (usually 3) main lobes, which are again sublobed, $6-10 \mu$ high, $8-13.5 \mu$ broad;



Fig. 3.

Asterina streptocarpi.

thyriothecia developed at the tips of lateral branches, flattened-hemispherical, reddish-brown, $100-150 \mu$ diameter, dehiscence stellate, formed of radiating hyphae $2.5-3 \mu$ thick (cells $5-6 \mu$ long); margin at first smooth, later fimbriate, a number of flexuous hyphae running out and anastomosing with the mycelial hyphae; asci paraphysate, four-spored, sessile, not staining blue with iodine, globose or broadly ovate, $20-24 \times 23-24 \mu$; spores conglobate, two-celled, brown, smooth, constricted, cylindrical, broadly rounded at both ends, $16-18 \times 7-8.5 \mu$, cells equal, or upper cell slightly broader.

This species is closely related to *A. undulata*, *A. gerbericola*, and *A. dilabens*, Syd.; like the two former, it has thyriothecia, which originate at the tips of lateral branches. It is nearest *A. gerbericola*, from which it differs chiefly in its smooth spores, coarser, undulate hyphae, and more numerous hyphopodia.

Asterina Streptocarpi, Doidge, n. sp.

Plagulas epiphyllas, atras, minutas, usque 3 mm. diametro formans; mycelium undulatum, ex hyphis flexuosis, ramosis, 3.5–5 μ latis, septatis (articulis 16–20 μ longis) compositum; hyphopodia continua, modice copiosa, alternantia, plurilobata, 6–10 μ alta, 8–13.5 μ lata; thyriothecia in summis hyphis secundariis oriunda, orbicularia, 100–150 μ diametro, brunnea, radiatim ex hyphis, 2.5–3 μ crassis (cellulis 5–6 μ longis) contexta; ad ambitum hyphis flexuosis fimbriata, stellatim dehiscencia; asci aparaphysati, 4-sporei, ovato-globosi, 20–24 $\times$ 23–24 μ ; sporae conglobatae, 1-septatae, constrictae, leves, cylindraceae, utrinque late rotundatae, 16–18 $\times$ 7–8.5 μ , cellulis aequalibus v. superiore leniter latiore.

Hab. in foliis *Streptocarpi Rexii*, Deepwalls, Knysna District, 13.5.23, leg. Doidge [17223].

A. gerbericola affinis.

100. *Asterina celtidicola*, P. Henn., var. *microspora*, Doidge.

Trans. Roy. Soc. S. Africa, VIII, p. 260, 1920.

On leaves of *Kiggelaria africana*, Keurkloof Forest, George District, 9.5.23, Doidge [17111]; Deepwalls, Knysna District, 13.5.23, Doidge [17221]; Storms River, Humansdorp District, 15.5.23, Doidge [17191].

On stems of *Capparis citrifolia*, Ebb and Flow, Wilderness, George District, 10.5.23, Doidge [17123].

This variety was originally collected on *Maerua pedunculosa* and *Oncoba Kraussiana* in Natal.

101. *Amazonia Goniomae*, n. sp.

On leaves of *Gonioma Kamassi*, Deepwalls, Knysna District, 13.5.15, Doidge [17209]; Storms River, Humansdorp District, 15.5.23, Doidge [17230]; Keurkloof Forest, George District, 9.5.23, Doidge [17109].

Amphigenous, forming irregularly circular, dull-black spots up to 7 mm. diameter; mycelium dense, radiating, consisting of meliola-like hyphae 6.5–9 μ thick, frequently septate, cells 20–27 μ long, with numerous usually opposite, less frequently unilateral branches; capitate hyphopodia, two-celled, stipitate, straight or curved, clavate-cylindrical, sometimes sublobed, 20–27 $\times$ 8.6–10 μ , stalk cell short, cylindrical; thyriothecia scattered, 160–200 μ diameter, flattened, shield-shaped, circular, formed of radiating hyphae about 6 μ thick, margin very briefly fimbriate; asci aparaphysate, evanescent, two-spored; spores flattened-cylindrical, 4-septate, slightly constricted at the septa, brown, broadly rounded at both ends, 40–47 $\times$ 15–16.5 $\times$ 10–12 μ .

Amazonia Goniomae, Doidge, n. sp.

Amphigena, plagulas atras irregulares v. orbiculares, usque 7 mm. diametro formans; mycelium densum, ex hyphis, brunneis, 6.5–9 μ crassis, septatis, ramosis compositum, cellulis 20–27 μ longis, ramis numerosis, oppositis v. rarius unilateralibus, hyphopodia capitata alternantia, numerosa, recta v. curvata, stipitata, 20–27 $\times$ 8.6–10 μ , cellula superiore clavato-cylindracea interdum sublobata; thyriothecia sparsa orbicularia, 160–200 μ diametro, radiatim ex hyphis, ca. 6 μ crassis contexta, ambitu hyphis similibus fimbriata; asci aparaphysati bisporei; sporae oblongae, 4-septatae, leniter constrictae, brunneae, utrinque late rotundatae, 40–47 $\times$ 15–16.5 μ .

Hab. in foliis *Goniomae Kamassi*, Deepwalls, Knysna District, 13.5.23, leg. Doidge [17209].

102. *Morenoella Phillipsii*, n. sp.

On leaves and stems of *Ocotea bullata*, Deepwalls, Knysna District, 13.5.23, Doidge [17205]; Montagu Pass Road, George District, 9.5.23, Doidge [17127].

Forming thin, spreading black growths on the under-side of the leaves, and densely clothing the stems and petioles of young seedlings. Hyphae fuscous, undulating, $3-3.5\ \mu$ thick, branching irregular, septa obscure, hyphopodia alternate or unilateral, continuous, briefly cylindrical, straight or subuncinate, $6-15 \times 3.5-5\ \mu$, sometimes anastomosing with another hyphopodium or with a branchlet to form a circle or oval; thyriothezia numerous, scattered, elliptic to linear, occasionally almost circular, straight, curved or bent at an angle, $240-600 \times 90-160\ \mu$, radiating in structure, formed of hyphae about $3\ \mu$ thick, raggedly fimbriate at the margin, dehiscing by a longitudinal slit, which later becomes widely distended, exposing the asci: asci aparaphysate, eight-spored, ovate, numerous, thickened



Fig. 4.

Morenoella Phillipsii.

round apex, sessile or with a short foot, not staining blue with iodine, $26-33.5 \times 18-20\ \mu$; spores tristichous or conglobate, fuscous, 1-septate, ellipsoid, rounded at both ends, very deeply constricted, so that the cells very often fall apart before the spore germinates, $13.5-15 \times 5-6.3\ \mu$, each cell ellipsoid.

My attention was drawn to this fungus by Mr. Phillips, the Assistant Officer for Sylvicultural Research, who is stationed at Deepwalls, in the Knysna District, and is conducting investigations into the regeneration of the indigenous forest. The stems of young *Ocotea bullata* seedlings are frequently densely clothed with the mycelium of *Morenoella Phillipsii*, which completely covers the stems for 10, 20, or 30 cm. It is not

improbable that the fungus has a serious effect on the growth of stinkwood seedlings.

In several stem sections I thought that I detected the presence of a subcuticular hypostroma, but failed to establish any connection between this and the thyriothechia; there was no trace of this in leaf sections. I have therefore assigned the fungus to the genus *Morenoella*.

Morenoella Phillipsii, Doidge, n. sp.

Hypophylla et caulicola, plagulas tenues, effusas hypophyllas formans atque caules late et dense obtegens; mycelium ex hyphis fuscis, undulatis, 3-3.5 μ latis, ramosis, obscure septatis compositum; hyphopodia alternantia, breve cylindracea, recta v. subuncinata, 6-15 $\times$ 3.5-5 μ ; thyriothechia numerosa, sparsa, elliptica v. linearia, recta, curvata vel geniculata, 240-600 $\times$ 90-160 μ , radiatim ex hyphis ca. 3 μ crassis composita, ambitu fimbriata, primum rima angusta longitudinali deinde late aperta; asci paraphysati, octospori, ovati, numerosi, apice incrassati, 26-33.5 $\times$ 18-20 μ ; sporae tristichae v. conglobatae, fuscae, medio septatae et valde constrictae, ellipsoideae, 13.5-15 $\times$ 5-6.3 μ .

Hab. in foliis *Ocoteae bullatae*, Deepwells, Knysna District, 13.5.23, leg. Doidge [17205].

103. *Phragmothyriella parenchymatica*, Doidge.

Syn. *Zukalia parenchymatica*, Doidge (Trans. Roy. Soc. S. Africa, VIII, p. 114, 1920).

On leaves of *Xymalos monospora*, Kentani, 3.3.15, Pegler [8864].

Thyriothechia epiphyllous, scattered, 300-400 μ diameter and about 170 μ high, fuscous, irregularly parenchymatous in texture, paler towards the margin, with a single polyscous hymenium; asci numerous, ellipsoid-ovate, briefly pedicellate, 50-60 $\times$ 20-30 μ ; spores obliquely distichous or conglobate hyaline or yellowish, 5-7-septate, ellipsoid, sometimes slightly curved, 30-40 $\times$ 8-10 μ ; mycelium pale yellowish-brown, consisting of hyphae, 3.5-5 μ thick; these are copiously branched and anastomosing, forming in the neighbourhood of the perithecia a continuous pseudo-parenchymatous layer, so that the covering wall of the thyriothecium merges gradually into the mycelium.

104. *Ascostratum insigne*, Syd.

Ann. Myc., X, 1912, pp. 41-42, fig. 2.

On rind of *Euphorbia* sp., Amanzimtoti, Natal, 10.7.11, Doidge [1660], 20.5.13, Doidge [6625].

In each case closely associated with *Pleomassaria gigantea*, Syd.

Stromata scattered, erumpent, later superficial, round or irregular in form, rather flat, $\frac{1}{2}$ - $\frac{2}{3}$ mm. diameter, the entire length of the base being attached to the host; outer wall dark-brown, almost black; inner part of stroma composed of light-coloured plectenchyma, and containing numerous irregularly scattered loculi; loculi each with a single ascus; asci ovate-globose, slightly thickened round the apex, eight-spored, 60-70 $\times$ 40-55 μ ; spores oblong or broadly fusiform, straight or often asymmetrical, obtuse at both ends, 7-9-septate, not constricted, hyaline, 45-58 $\times$ 16-18 μ .

105. *Irene implicata*, n. sp.

On leaves of *Chilianthus arboreus*, Van Stadens Pass, 19.5.23, Doidge [17251].

Mycelium hypophyllous, forming rather indefinite round, grey-black spots, 2-4 mm. diameter. Hyphae pale fuscous, rather thin-walled, subtorulose, 5-6.6 μ thick, cells variable in length, branching irregularly anastomosing and forming a tangled network amongst the lepidote scales on the under side of the leaf. Capitate hyphopodia usually alternate or unilateral, sometimes opposite, stipitate, 20-40 $\times$ 10-20 μ ; stipe cell usually cylindrical or constricted at base, 3.5-16 μ in length; terminal cell very variable in form and dimensions, club-shaped, truncate, or variously and fantastically lobed. Mucronate hyphopodia, solitary, opposite or in whorls of 3 or 4, slender lageniform, straight or curved, 16-27 $\times$ 5-6.5 μ . Setae none. Perithecia crowded together in the centre of the mycelium globose, somewhat

flattened, verrucose, carbonaceous, 240–290 μ diameter, collapsing at maturity; outer wall composed of rounded or conical cells, which are irregularly polygonal at the base and about 10 μ in diameter. Asci two-spored, evanescent. Spores brown, 4-septate, elliptic, constricted at the septa, straight or slightly curved, tapering somewhat toward the rounded ends, 50–54 μ long; medial cell longer than the others and 20–23.5 μ broad; terminal cells are not more than 13–14 μ broad.

Irene implicata, Doidge, n. sp.

Hypophylla, plagulas parum perspicuas, griseolas, orbiculares, 2–4 mm. latis formans; mycelium laxè ramosum, implicatum ex hyphis fuscis, subtorulosis, 5–6.6 μ crassis compositum; hyphopodia capitata alternantia v. raro opposita, 20–40 $\times$ 10–20 μ , cellula inferiore cylindrica v. basim constricta, 3.5–16 μ longa, superiore irregularia, clavata




Fig. 5.

Irene implicata: mycelium with capitate and mucronate niphopodia, and spores.

truncata v. varie lobata; hyphopodia mucronata plerumque opposita, angusta, lageniformia, recta v. curvata, 16–27 $\times$ 5–6.5 μ ; perithecia in centro plagularum pauca conferta, atra, carbonacea, verrucosa, 240–290 μ , in sicco collapsa; asci bispori: sporae brunneae, 4-septatae, ellipticae ad septa constrictae, rectae v. subcurvatae; utrinque attenuatae, 50–54 μ longae; cellula media 20–23.5 μ lata, cellulis extimis 13–14 μ latis.

Hab. in foliis *Chilianthi arborescentis*, Van Stadens Pass, 19.5.23, leg. Doidge [17251].

106. *Irene Peglerae*, Doidge.

Trans. Roy. Soc., V, p. 730, 1916.

On *Halleria lucida*, Keurkloof Forest, George District, 9.5.23, Doidge [17116]; Storms River, Humansdorp District, 15.5.23, Doidge [17182]; Deepwalls, Knysna District, 13.5.23, Doidge [17200].

This species was originally collected at Kentani on *Anastrabe integerrima*; it appears to be very common on *Halleria lucida* in the coast forest belt.

107 *Meliola comata*, Doidge.

Trans. Roy. Soc. S. Africa, VIII, p. 111, 1920.

On leaves of *Pyrenacantha scandens*, Woodville Forest, George District, 11.11.17, Doidge [11020]; Storms River, Humansdorp District, 15.5.23, Doidge [17192]; Deepwalls, Knysna District, 13.5.23, Doidge [17208].

The host of *M. comata* was previously (loc. cit.) incorrectly determined as (?) *Ipomoea* sp.

108. *Meliola Evansii*.

Trans. Roy. Soc. S. Africa, VIII, p. 112, 1920.

On leaves of *Celastrus acuminatus*, Deepwalls, Knysna District, 13.5.23, Doidge [17206].

On *Elaeodendron croceum*, Deepwalls, Knysna District, 13.5.23 [17216].

M. Evansii has previously been recorded on *Mystroxydon* at Mossel Bay and *Celastrineae* undetermined from the Zoutpansberg.




Fig. 6.

Meliola Knowltoniae.

109. *Meliola Knowltoniae*, n. sp.

On leaves of *Knowltonia vesicatoria*, Storms River, Humansdorp District, 15.5.23, Doidge [17177].

Epiphyllous, forming rather thin, black, carbonaceous spots up to 5 mm. diameter: mycelium consisting of brown, opaque, rather sinuous hyphae 9–11 μ thick (usually 10 μ): cells for the most part 35–40 μ long, branches opposite: capitate hyphopodia alternate, brown, opaque, usually one to each cell, stipitate, 27–34 $\times$ 13.5–17 μ , majority at an angle of about forty-five degrees with hypha, but sometimes more widely divergent, stipe cell cylindrical, 6.5–10 μ long, terminal cell club-shaped, sublobed or subuncinate, curving towards or away from the hyphae, but usually the latter; mucronate hyphopodia fairly numerous near centre of mycelium, opposite, not lighter in colour than the capitate hyphopodia, 16–17 $\times$ 8–10 μ , ampulliform with rather a short neck, 3.5 μ thick; mycelial setae numerous in the neighbourhood of the perithecia, less numerous elsewhere, simple, straight or somewhat flexuous, 400–500 μ long, 10 μ thick at base, the majority tapering very gradually to the blunt and somewhat less opaque apex, which is 2–3 μ thick, a few of the setae are less tapering and have a blunt apex up to 5 μ thick; perithecia in groups in the centre of the mycelium, spherical, black, carbonaceous, slightly rough, 160–200 μ diameter; asci two-spored, evanescent; spores 4-septate, cylindrical-compressed, broadly rounded at both ends, slightly constricted at the septa, brown, 42–46 $\times$ 16.5–20 $\times$ 13.5 μ .

Meliola Knowltoniae, Doidge, n. sp.

Epiphylla, plagulas atras, usque 5 mm. diametro formans; mycelium ex hyphis, brunneis, opacis, subsinuosis, 9–11 μ crassis, septatis, ramosis, compositis, cellulis plerumque, 35–40 μ longis, ramis oppositis; hyphopodia capitata alternantia stipitata 27–34 $\times$ 13.5–17 μ , cellula superiore, clavata, sublobata v. subuncinata; hyphopodia mucronata sat numerosa, opposita, 16–17 $\times$ 8–10 μ , ampullacea; setae mycelicae rectae v. subflexuosae, simplices 400–500 μ longae, basi 10 μ crassae, apicem obtusum v. sensim attenuatae; perithecia gregaria, globosa, atra, carbonacea, 160–200 μ diametro; asci bispori; sporae 4-septatae, cylindratae, utrinque late rotundatae, leniter constrictae. 42–46 $\times$ 16.5–20 μ .

Hab. in foliis *Knowltoniae vesicatoriae*, Storms River, Humansdorp District, 15.5.23, leg. Doidge [17177].

110. *Meliola Toddaliae*, Doidge.

Trans. Roy. Soc. S. Africa, V, p. 732, 1916.

On leaves of *Fagara capensis*, Thun., Pirie Forest, 8.7.19, Doidge [12275]; Winters Kloof, Natal, 13.9.19, Doidge [12440].

On *Fagara Davyi*, Gouna Road, Low Forest, February, 1922, Keet [15531].

M. Toddaliae was originally described as occurring on *Toddalia lanceolata*, and is commonly found on this host.

111. *Perisporium irenicolum*, n. sp.

On *Irene Peglerae* on leaves of *Halleria lucida*, Deepwalls, Knysna District, 13.5.23, Doidge [17201].

Mycelium pale fuscous, hyphae very delicate, 1–2 μ thick, but well developed, forming a dense tangle between the hyphae of the *Irene*. Associated with this mycelium, and probably connected with it, are very numerous conidiophores and conidia of a *Helminthosporium* sp. Stevens (*Bot. Gazette*, 64, 1918, pp. 228–229) noticed similar conidia in connection with *Perisporium Meliolae*. Perithecia subspherical, rather taller than broad, smooth, 190–230 μ diameter, wall membranous, formed of irregular, polygonal cells about 6 μ diameter. Asci numerous, fascicled, eight-spored, clavate or cylindrical, sessile or with a short foot, 70–100 $\times$ 18–20 μ , thickened round the apex, staining dull-purple with iodine, paraphysate. Spores distichous, fuscous, clavate, 5-septate when mature, slightly constricted at the medial septum, broadly rounded above, tapering to the lower end, 42–44 μ long, 13.5–16 μ thick at the broadest point, narrower and obtuse at the lower end.

The genus *Perisporium* was discarded by Theissen and Sydow (*Ann. Myc.*, 1917, p. 448), and the genus *Meliolina* established for species similar to *Meliola*, but with

ahyphopodiate mycelium. The present species does not agree well with the species of the genus *Meliolina* in that it has eight-spored fasciculate asci, but it is closely related to *Perisporium Meliolae*, Stevens (loc. cit.); it is therefore provisionally assigned to the genus *Perisporium*.

(?) *Perisporium irenicolum*, Doidge, n. sp.

Amphigenium in mycelio *Irene Peglerae* parasiticum; mycelio tenue, bene evoluto, hyphas fuscas, 1-2 μ cr. composito; perithecia subsphaerica, laeves, membranacea, 190-230 μ diametro; asci numerosi, paraphysati, fasciculati, octospori, clavati v. cylindranei, sessiles v. breviter pedicellati, 70-100 $\times$ 18-20 μ , apice incrassati; sporae distichae, fuscae, clavatae, 5-septatae, medio leniter constrictae, 42-44 $\times$ 13.5-16 μ .

Hab. in foliis *Halleriae lucidae*, Deepwells, Knysna District, 13.5.23, leg. Doidge [17201].

112. *Phaeostigme circumsedens*, n. sp.

Parasitic on mycelium of (?) *Asterina* or (?) *Clypeolella* sp., on leaves of *Mikania capensis*, Storms River, Humansdorp District, 15.5.23, Doidge [17163].

Mycelium delicate, pale fuscous or olivaceous, *without hyphopodia*, composed of hyphae 2-3 μ thick, which branch and anastomose irregularly and closely invest the hyphae of the host. Perithecia scattered, spherical, somewhat papillate, but without a typical ostiole, subverrucose, brown, 100-120 μ diameter, wall consisting of irregular, polygonal cells, 6-10 μ diameter; asci eight-spored, paraphysate, narrow cylindrical or obclavate, thin-walled, rounded but not thickened at the apex, 40-50 $\times$ 13-16 μ ; paraphyses filiform; spores distichous, two-celled, constricted, smooth, fuscous, cylindrical or broadly ellipsoid, broadly rounded at both ends, 11-13 $\times$ 5-6.3 μ , cells subequal and almost spherical, or upper cell slightly broader and more broadly rounded than the lower.

Phaeostigme circumsedens, Doidge, n. sp.

Mycelium in (?) *Asterina* sp. v. (?) *Clypeolella* sp. parasiticum, mycelium (?) *Asterinae* arete circumsedens, ex hyphis tenuibus, fuscidulis v. olivaceis, 2-3 μ latis, dense intricatis, ramosis et anastomosantibus, septatis compositum; perithecia sparsa, globosa, ostiolo typico parentia, 100-120 μ diametro, parenchymatica e cellulis, 6-10 μ diametro, composita; asci paraphysati, fasciculati, anguste cylindranei v. obclavatae, 40-50 $\times$ 13-16 μ ; sporae distichae, 1-septatae, leves, fuscae, oblongae v. late ellipsoideae, utrinque late rotundatae, 11-13 $\times$ 5-6.3 μ , loculis subaequalibus v. loculo supero paullo latiore.

Hab. in foliis *Mikaniae capensis*, Storms River, Humansdorp District, 15.5.23, leg. Doidge [17163].

113. *Chaetostigmella capensis*, Doidge.

Syn. *Phaeodimeriella capensis*, Doidge (Trans. Roy. Soc. S. Africa, V, p. 719, 1917, Plate LVIII, fig. 8).

On leaves of *Apodytes dimidiata*, Knysna, Cape Province, 3.6.12, Pienaar [2426].

The genus *Chaetostigmella* has been established by Sydow (Ann. Myc., 1917, p. 199) for the species with paraphyses, formerly assigned to the genus *Phaeodimeriella*. The mycelium of *Ph. capensis* was incorrectly described (loc. cit.) as having hyphopodia. The hyphopodiate mycelium is that of an *Asterina*, on which the *Chaetostigmella* is parasitic. *Ch. capensis* has abundant hyphae, 2-3 μ thick, which are at first fuscous, then brown, closely investing the hyphae of the host, branching profusely and forming a dense network between the hyphae of the *Asterina*.

114. *Chaetostigmella asterinicola*, Doidge.

Syn. *Phaeodimeriella asterinicola*, Doidge (Trans. Roy. Soc. S. Africa, VIII, p. 115, 1920).

This species, having paraphysate asci, must also be assigned to the genus *Chaetostigmella*.

115. *Phaeodimeriella plumbea*, n. sp.

Parasitic on *Asterina gerbericola*, Storms River, Humansdorp District, 15.5.23, Doidge [17176].

Mycelium well developed, closely investing the hyphae of the *Asterina*, and forming an open network between the hyphae of the host; mycelium greyish-green, younger hyphae almost hyaline, very slender, 1–2 μ thick, branching irregularly; perithecia numerous, black, globose, 80–100 μ diameter, with a crown of 6–12 setae; setae straight, brown, 3.5–5 μ thick at the base, tapering to an acute apex, 55–70 μ long; asci numerous, paraphysate, fasciculate, eight-spored, ellipsoid, sessile, 36.5–43 $\times$ 10–13.3 μ ; spores distichous, brown, narrow-ellipsoid, 13.3–15 $\times$ 3.3–3.5 μ , upper cell somewhat broadened just above the septum, giving the effect of a slight constriction at the septum.

Phaeodimeriella plumbea, Doidge, n. sp.

Mycelium hyphas *Asterinae* dense amplexans, ex hyphis plumbeis v. viridulis, 1–2 μ crassis, tenuibus, dense reticulatis, ramosis et anastomosantibus, septatis compositum; perithecia superficialia, atra, 80–100 μ diametro, setis 6–12 coronatis; setis rectis, brunneis, basim 3.5–5 μ crassis, ad apicem acutem attenuatis, 55–70 μ longis; asci numerosi, fasciculatis, paraphysati, octospori, ellipsoidei, sessiles, 36.5 $\times$ 43 $\times$ 10–13.3 μ ; sporae distichae, anguste ellipsoideae, 1-septatae, brunneae, leniter constrictae, locula supero latiore, 13.3–15 $\times$ 3.3–3.5 μ .

Hab. in foliis *Gerberae cordatae*, Storms River, Humansdorp District, 15.5.23, leg. Doidge [17176].

116. *Rizalia confusa*, n. sp.

On leaves of *Olea capensis*, Storms River, 15.5.23, Doidge [17172]; Assegai Bosch, 16.5.23, Doidge [17233].

On *Olea Pegleri*, Kentani, 6.5.15, Pegler (Pegler No. 2316) [9072] (immature).

Hypophyllous, forming very numerous, minute olivaceous or rusty-brown spots; these soon become confluent and cover large areas of the leaf-surface. Mycelium variable: there is a delicate mycelium composed of fine, pale fuscous hyphae, barely 1 μ thick, which branches and anastomoses to form an irregular network and produces conidia and erect, stouter hyphae. The conidia are pale fuscous, 4–7-septate, narrow clavate, 30–40 μ long, 3.5–5 μ broad at the rounded tip, tapering to a thickness of 1 μ at the base. The erect hyphae [(?) setae] are coarser, dark-brown, septate, obtuse, 3–3.5 μ thick and 60–100 μ long. The fine mycelium apparently gives rise to the coarser, darker mycelium, which produces the perithecia. This consists of abundant, straight, olivaceous or brown hyphae, 3–3.5 μ thick, without hyphopodia, branching and anastomosing irregularly and with some semi-erect branches. The perithecia are numerous, black, spherical with a rough wall, 80–100 μ diameter, bearing a few short, more or less erect hyphae, which are irregular in size and number. The lower half of the perithecium is surrounded by a tangle of hyphae. Asci few in each perithecium, eight-spored, sessile, paraphysate, ovate or broadly clavate, thin-walled, somewhat thickened round the apex, 40–50 $\times$ 23–24 μ . Spores parallel, two-celled, narrow-cylindrical, not constricted, hyaline, not tapering to the rounded ends, 23–24.5 $\times$ 6.5–8 μ , cells equal or subequal.

Rizalia confusa, Doidge, n. sp.

Hypophylla; mycelium bene evolutum, ex hyphis olivaceis, 3–3.5 μ crassis, ramosis, anastomosantibus, ahyphopodiatis compositum, setis erectis v. suberectis nonnullis; perithecia numerosa, basi mycelio suffulta, extus setis paucis, atris, irregulariter obsita, globosa, carbonacea, 80–100 μ diametro; asci pauci in quoque perithecio, octospori, paraphysati, sessiles, ovati v. late clavate, apice leniter incrassati, 40–50 $\times$ 23–24 μ , sporae parallelae, medio septatae, cylindratae, haud constrictae, hyalinae, utrinque rotundatae et haud attenuatae, 23–24.5 $\times$ 6.5–8 μ .

Hab. in foliis *Oleae capensis*, Storms River, Humansdorp District, 15.5.23, leg. Doidge [17172].

117. *Theissenula Woodiana*, Doidge.

Syn. *Zukalia Woodiana*, Doidge (Trans. Roy. Soc. S. Africa, VIII, p. 114).

On leaves of *Celastrus alatus*, Winkle Spruit, Natal, 20.5.15, Doidge [9008]; Durban, 22.7.15, Medley Wood [9061].

The genus *Zukalia*, which was originally described as a "*Meliola hyalophragmia*," has been merged by Theissen and Sydow, in their recent revision of the Perisporiales, in the genus *Chaetothyrium* (Ann. Myc., XV, p. 477). An examination of the fungi classified under the genus *Zukalia* in the cryptogamic section of the National Herbarium shows that neither *Z. Woodiana* nor *Z. parenchymatica* can be placed in the genus *Chaetothyrium*; *Z. Woodiana* very closely resembles the type species of *Theissenula*, from which it differs chiefly in the occasional presence of 5-septate spores.

Hypophyllous, forming rusty-brown spots 10–15 mm. diameter; these are frequently so numerous as to coalesce and form continuous irregular brown patches. Mycelium fuliginous, tangled at the centre of the spots, radiating towards the circumference. Hyphae not hyphopodiate, about 3 μ thick, copiously branched. Conidia not observed. Perithecia numerous, crowded in irregular groups, superficial, globose, 95–115 μ diameter; perithecial wall at first parenchymatous in appearance. Asci paraphysate, several in each perithecium, eight-spored, ellipsoid or ovate, briefly pedicellate, 30–40 $\times$ 13–17 μ . Spores hyaline or yellowish, subdistichous or parallel in the ascus, narrow ellipsoid or subclavate; at first subequally 1-septate, later 3-septate or occasionally 5-septate, 18–24 $\times$ 4–5 μ . The spore readily falls apart at the medial septum.

118. *Parenglerula Macowaniana* (Thüm), v. Höhn.

Frag. zur Myk., X, No. 525, in Sitzungsber. der Kais. Akad. d. Wiss. in Wien: Math. Nat., Kl. CXIX, 1, p. 465, 73, 1910.

Theissen, F., Mykologische Abhandlungen. Verk. Zool.-bot. Gesellsch. Wien, LXVI, pp. 347–350, 1916.

Syn. *Meliola Macowaniana*, Thüm, Flora, LX, p. 204, 1877; *Asterina Macowaniana*, Kalch et Cke., Grevillea, VII, p. 57, 1878, IX, p. 33, 1880, with diagnosis; Syll. Fung., I, p. 41; Exsicc. Thumen. Myc. Ilino., 568; Rehm. Ascom. 395.

On leaves of *Celastrus buxifolius*, South Africa.

On leaves of *Celastrus buxifolius*, Ashburton, Natal, 22.4.16, Doidge [9700]; Bedford-Cape Province, 20.11.17, Doidge [10892]; Belmont Valley, Grahamstown, 15.11.17, Doidge [10949]; Alice, Cape Province, 18.11.17, Doidge [10973]; Grahamstown, 4.6.19, Britten [14213].

On *Celastrus polyacanthus*, Assegai Bosch, Humansdorp District, 16.5.23, Doidge [17246].

On *Elaeodendron capense*, Van Stadens Pass, Cape Province, 13.11.17, Doidge [10890].

On *Cassine capensis*, Van Stadens Pass, Cape Province, 13.4.17, Doidge [10885].

On *Scolopia Mundii*, Assegai Bosch, Humansdorp District, 16.5.23, Doidge [17245].

I have not had an opportunity of examining the original specimens, but from v. Höhnel's description there can be no doubt that the fungi mentioned above must be assigned to this species. The description is taken from that of v. Höhnel and Theissen; an examination of several recent collections entirely confirms their observations.

Usually epiphyllous, forming black circular spots 2–3 mm. diameter, which are usually so numerous as to become confluent, and large irregular areas are then covered by the mycelium. In the specimen on *Scolopia Mundii* [17245] the mycelium forms larger spots, up to 7 mm. diameter, and the leaf tissues underneath are discoloured to a yellowish-brown. No such discoloration was observed on the *Celastrus* leaves.

Mycelium consisting of stout, dark-brown, septate undulating, coarse-walled hyphae, 6–8 μ thick, branching profusely, spreading in an irregularly radial direction and furnished with numerous hyphopodia. Hyphopodia alternate or unilateral, unicellular, hemi-

spherical or irregularly oval, 6.5–14 μ high and 6–15 μ broad, having a light-coloured circular pore in the upper portion.

Perithecia produced in great profusion, densely aggregated in the central part of the mycelium, black, coarse, irregularly globular, very rough, almost tubercular, 50–120 μ in diameter, and without a trace of an ostiole. They are poised on a slightly narrowed base about 40 μ wide, from which the hyphae of the subiculum radiate. The perithecial wall disappears at maturity, falling apart into separate cellular filaments, which surround the asci and envelope them after the fashion of paraphyses. The asci and these filaments are embedded in a stiff mass of mucilage, staining a dingy-violet with iodine, which secretes an insoluble, dark-brown mass, which completely conceals the internal structure. The thin, coarse skin thus produced disintegrates, in consequence of the swelling of the mucilaginous content, into minute, irregular fragments often resembling cells, and thus simulates a dark-brown perithecial membrane.

The mature perithecia contain 1–10 oval, thick-walled asci, four- to eight-spored, broadly rounded above, constricted at base, 57–65 $\times$ 40–50 μ . The cellular filaments representing the remains of the perithecial membrane dissolved by mucilaginous histolysis are two- to four-septate, grey-brown to dirty-yellow, smooth-walled, not constricted at the septa, obtuse, usually incurved, 20–45 $\times$ 5–7 μ . They form at first a closed perithecial sheath, but are soon forced asunder, giving way to the outward pressure of the mucilage, which then forms a hardening coarse crust over the original membrane.

Spores bicellular, elongated ovoid, rounded at both ends, thick-walled, at first covered with a thin mucilaginous sheath, remaining hyaline for a long time, then turning dark-brown, 25–32 $\times$ 13.5–15 μ . The transverse wall is situated below the centre, one cell being 15.5–18 μ and the other 12–15 μ long; the position of the wall is therefore more variable than is indicated by v. Höhnelt or Theissen. The upper cell is 12–15 μ broad, and the lower 8.5–13 μ .

119. *Capnodium citricolum*, McAlp.

Proc. Linn. Soc. of New South Wales, 1896, Part IV, p. 491.

Syll. Fung., XIV, p. 476.

On twigs of *Citrus* sp., Karreekloof, Lydenburg District, 24.7.19 [12313].

Effuse, black, crustaceous, covering the entire surface of the affected parts; some of the hyphae creeping, greenish or hyaline, copiously branched, septate, 6–8.5 μ thick, ascending branches short, simple, septate, bearing conidia which are hyaline or pale-greenish, continuous, uni- or bisepate, spherical, oval or elliptical, slightly constricted, 7.5–24 $\times$ 4–11 μ , sometimes moniliform; other hyphae are dark-coloured, greenish-brown or almost black, very frequently septate, constricted, slightly or copiously branched, rigid, 9.5–11 μ thick, bearing brown conidia, which are mostly elliptic, uniseptate, 7.5–16 $\times$ 5.5–8.5 μ . Perithecia and pycnidia are intermingled, opaque, black, oblong, oval, flask-shaped or club-shaped, 112–250 $\times$ 52–112 μ . Asci cylindrical to clavate, sessile, rounded at the apex, four-, six- or eight-spored, 70–80 $\times$ 19–20 μ . Spores brown, oblong, sometimes subfusoid, typically obtuse at both ends, constricted near the middle, 5–6-septate, often with longitudinal or oblique septa, distichous, sometimes tristichous, about 21–24 $\times$ 8.5–9.5 μ . Paraphyses hyaline, often granulose, elongated-clavate, 9.5 μ broad at the apex, equalling the asci.

The description quoted above is that of McAlpine; this specimen agrees with it very well, and is the only South African specimen of *Capnodium* in the National Herbarium on which perithecia have been developed.

120. *Trichothyrium elegans*, n. sp.

On mycelium of (?) *Asterina* sp. on *Cunonia capensis*, Montagu Pass Road, George District, 9.5.23, Doidge [17129]; Storms River, Humansdorp District, 15.5.23, Doidge [17186]; Deepwalls, Knysna District, 13.5.23, Doidge [17220].

Epiphyllous, parasitic on the mycelium of (?) *Asterina* sp. or (?) *Clypeolella* sp., which is so heavily parasitized that no thyriothecia are formed, and consequently cannot be determined accurately; mycelium not forming a continuous pellicle, but closely investing the hyphae of the host, formed of delicate, pale fuscous hyphae about $3\ \mu$ thick, branching and anastomosing to form an open network between the hyphae of the host; hyphopodia none: perithecia radiate in structure, orbicular, $160\text{--}170\ \mu$ diameter, flat, shield-shaped, with a round central ostiole $15\text{--}25\ \mu$ diameter, radiating hyphae $2.5\text{--}3\ \mu$ thick; asci ovate-oblong or oblong, sessile, rounded and somewhat thickened at the apex, apophysate, $57\text{--}65 \times 16\text{--}17.5\ \mu$; spores distichous or tristichous, clavate, 1-septate, not or very slightly constricted, hyaline, $20\text{--}23 \times 6.5\text{--}8\ \mu$; upper cell shorter, broader, and more broadly rounded than the lower, which tapers toward the lower end.




Fig. 7.

Trichothyrium elegans.

Trichothyrium elegans, Doidge, n. sp.

Epiphyllum, in mycelio (?) *Asterinae* sp. parasiticum; mycelio hyphas *Asterinae* arcte circumsedente sed haud pelliculam continuum efformante, ex hyphis obscure fuscis, $3\ \mu$ latis, ramosis et anastomosantibus composito; hyphopodiis nullis; perithecia radiatim oriunda, orbicularia, $160\text{--}170\ \mu$ diametro, brunnea, dimidiato-scutata, centro ostiolo rotundo, $15\text{--}25\ \mu$ lato praedita, radiatim ex hyphis $2.5\text{--}3\ \mu$ latis composita; asci ovato-oblongi v. oblongi, sessiles, apice rotundati et leniter incrassati, non paraphysati, $57\text{--}65 \times 16\text{--}17.5\ \mu$;

spora distichae v. tristichae, clavatae, 1-septatae, haud v. vix constrictae, hyalinae, $20-23 \times 6.5-8 \mu$, cellula superiore brevior v. latior.

Hab. in foliis *Canoniae capensis*, Montagu Pass Road, George District, 9.5.23, leg. Doidge [17129].

121. *Trichothyrium robustum*, n. sp.

Parasitic on mycelium of *Meliola comata* on *Pyrenacantha scandens*, Deepwalls, Knysna District, 13.5.23, Doidge [17208A].

Mycelium pale fuscous, ribbon-like, forming a continuous pellicle, completely clothing the hyphae and hyphopodia of the *Meliola*, and extending for a short distance on either side; ribbon-like thallus following the course of the hyphae of the host and branching where its hyphae branch, somewhat lobed at the margin, the lobes corresponding in position with the hyphopodia, $40-70 \mu$ wide, formed of hyphae $2-3 \mu$ thick, which are prosenchymatous in appearance, running longitudinally in the centre, and spreading out fan-wise near the margin at an angle of about forty-five degrees. Perithecia very numerous,




Fig. 8.

Trichothyrium robustum.

gregarious, flattened, round, produced on the edges of the ribbon-like thallus, $120-160 \mu$ diameter, with a round, central pore, $13-17 \mu$ diameter, formed of radiating hyphae about 3μ thick, cells almost cubical; asci numerous, fasciculate, clavate or elliptic, sessile, eight-spored, $30-40 \times 13-16.6 \mu$, paraphysate; paraphyses not very numerous, hyaline, filiform; spores distichous, hyaline, 1-septate, clavate, slightly constricted at septum, $16-17 \times 5-6.5 \mu$; upper cell shorter and broader, broadly rounded or somewhat acute at the apex; lower cell tapering to the base.

Trichothyrium robustum, Doidge, n. sp.

Mycelium fuscum, mycelio hyphas *Meliolae* parasiticum vitis alternatim ramosis. $40-70 \mu$ latis lobatis, lobis apice rotundatis, pinnatim ex hyphis, $2-3 \mu$ latis prosenchymaticis; perithecia ad marginem thalli numerosa, gregarie rotundata-depressa.

120–160 μ diametro, ostiolo rotundo, 13–17 μ dato praedita; asci numerosi, fasciculati clavati v. elliptici, sessiles, octospori, paraphysati, 30–40 $\times$ 13–16.6 μ ; paraphysibus parum numerosis hyalinis, filiformibus: sporae distichae, hyalinae, 1-septatae, clavatae, leniter constrictae, 16–17 $\times$ 5–6.5 μ , cellula superiore latiore et brevior.

Hab. in foliis *Pyrenacanthae scandentis*, Deepwalls, Knysna District, 13.5.23, leg. Doidge [17208A].

Trichothyrium dubiosum affinis.

Actinopeltella, Doidge, nov. gen.

Omnia ut in *Actinopeltis* sed sporae hyalinae 1-septatae.




Fig. 9.

Actinopeltella nitida.

122. Actinopeltella nitida, n. sp.

Parasitic on the mycelium of *Asterina Streptocarpi* on leaves of *Streptocarpus Rexii*, Deepwalls, Knysna District, 13.5.23, Doidge [17225].

Subiculum absent; perithecia scattered, fuscous, brown, lenticular-turbinate, 100–120 μ diameter, about 50 μ high, composed of radiating hyphae 2.5–3 μ thick, ostiolo irregularly round, 16–20 μ diameter, surrounded by a single crown of 6–10 stout, dark-brown setae, spreading horizontally, up to 40 μ long, 3.5–4 μ thick at base, apex acute; asci aparaphysate, delicate, narrow-obclavate, sessile, not thickened round the apex, eight-spored, 40–50 $\times$ 10–13.5 μ ; spores distichous, hyaline, ellipsoid or subclavate, two-celled, not constricted, rounded at the ends, 10–13 $\times$ 3–4.5 μ , cells subequal, or upper cell slightly broader.

Actinopeltella nitida, Doidge, n. sp.

Subiculum nullum; perithecia sparsa, fusco-brunnea, rotundato depressa, 100–120 μ diametro, radiatum ex hyphis, 2.5–3 μ crassis contexta, ostiolo rotundo 16–20 μ diametro praedita, setis brunneis 6–10, usque 40 μ longis, acutis coronata; asci aparaphysati, anguste obclavati, octospori, 40–50 $\times$ 10–13.5 μ ; sporae distichae, hyalinae, ellipsoidae v. subclavatae, 10–13 $\times$ 3–4.5 μ , cellulis aequalibus v. superno paulo latiore.

Hab. in foliis *Streptocarpi Rexii*, Deepwalls, Knysna District, 13.5.23, leg. Doidge [17225].

123. Tripospora tripos (Cke.), Lind.

Engl. and Prantl., Die Nat. Pflanzenfamilien, 1, p. 413, 1897.

Fitzpatrick, Monograph of the Coryneliaceae, Mycologia, 12, pp. 232–233, figs. 22–25, 44, 45, 1920.

Syn. *Corynelia tripos* (Cke.), Grevillea, VIII, p. 34, 1879; *Tripospora Cookei* (Cke.), Sacc. in Berl. et Vogl. Additamenta Syll. Fung., p. 194, 1886.

On *Podocarpus elongata*, Hk., Somerset East, Cape Province, leg. MacOwan, Rabh. Wint. Fung. Europaei 3150 [3490]; Pirie Forest, Kingwilliamstown, 10.1.14, District Forest Officer [7355]; near Uitenhage, June, 1915, Paterson [9047]; Kingwilliamstown, 19.6.14, District Forest Officer [7816]; Van Stadens Pass, Cape Province, 13.11.17, Doidge [10867]; Maritzburg, 1.2.21, Sim [14274].

Stromata bearing a compact cluster of perithecia, 0.5–1.5 × 0.5 mm., not irregularly scattered, arranged definitely in rows and becoming confluent; these rows reaching sometimes a length of 10 mm., but usually shorter, several rows frequently formed on the surface of one leaf; perithecia occurring on the stroma in a compact cluster of 2–16 (usually 4–8), and on isolated stromata radiating towards all sides, so oriented when the stromata form a row that they point to the left and right, the appearance of the fungus thus becoming regular. Young perithecium definitely flask-shaped, with a roughened, spherical to ovoid, ascigerous basal portion and a long cylindrical, glabrous, shiny neck, which is rounded at the tip and blunt. The neck of the perithecium in early stages provided with a canal and marked at the apex with a minute umbilicus, but closed; in later stages the apex of the beak flattened to form a slightly convex disk, the diameter of which equals that of the ascus-bearing portion of the perithecium; this disk becoming fimbriate-lacerate, and assuming a reddish-brown, fuzzy appearance, finally definitely dehiscent, the margin recurving, exposing the lighter-coloured inner wall of the neck of the perithecium, and resulting in the formation of a broad, funnel-shaped cavity, the centre of which is usually filled with a black mass of spores. Immature asci, 30–35 × 40–60 μ ; long-stalked, eight-spored, ovate evanescent. Ascospores very characteristic, unicellular, star-shaped, consisting of four (rarely five) conical, sharp-pointed projections, radiating from a rounded central portion, hyaline, when young becoming dark-brown, at maturity opaque and nearly black, thick-walled, 22–34 μ diameter (measured from tip to tip of adjacent projections).

124. *Corynelia fruticicola* (Pat.), v. Höhn.

Sitzber, Kais. Akad. Wiss. Wien., 120, p. 450, 1911.

Fitzpatrick, Monograph of the Coryneliaceae, Mycologia, 12, pp. 240–241, figs. 20, 21, 1920.

Syn. *Corynelia carpophila*, Syd., Engl. Bot. Jahrb., 45, p. 264, 1920.

On fruit of *Myrsine melanophleos*, Woodbush, Zoutpansberg District, 5.8.11, Doidge [1736]; Kentani, Cape Province, 5.10.12, May, 1913, and 3.6.14, Pegler [5617, 6621, and 7795]; Tugela Valley, near Mont aux Sources, Natal, 15.5.20, Doidge [14121]; Duncairn, near Maritzburg, 13.7.21, Doidge [14865]; Greytown, Natal, 16.2.22, Doidge [15410].

Stromata fruticulous, pulvinate, black; usually rounded, often laterally confluent to form an extensive effuse, black crust, which partially or completely envelops the fruit: individual stromata 1 mm. or less in diameter, covered by a crowded cluster of perithecia, frequently showing many stages of development on a single stroma; perithecium resembling closely that of *Tripospora tripos*, definitely flask-shaped, with a spherical to ovoid, ascigerous, basal portion, and a long, narrow-cylindrical neck, rounded and blunt at the apex. The neck of the perithecium provided with a canal and in early stages marked at the apex with a minute umbilicus, but closed: in later stages the apex broadened to form a wide, slightly convex disk, the diameter of which equals or exceeds that of the basal ascigerous portion: this disk becoming fimbriate-lacerate, and assuming a reddish-brown, fuzzy appearance, finally dehiscent, the margin recurving, forming a funnel-shaped opening, and exposing the lighter-coloured inner wall. Asci ovate to clavate, long-stalked, thin-walled, evanescent, eight-spored, paraphysate, 11–14 × 20–25 μ . Ascospores when young hyaline, smooth, and by mutual pressure polyhedral; at maturity spherical, brown, thick-walled, echinulate, unicellular, 6.5–10.5 μ in diameter. Pycnidia seated on the stroma among the perithecia more or less globose, black; pycnosporos hyaline, elongated to allantoid, 4–6 × 1 μ .

125. *Corynelia uberata*, Fr.

Ex Acharius Systema Mycologicum, 2, p. 535, 1822.

Fitzpatrick, Monograph of the Coryneliaceae, Mycologia, 12, pp. 247–251, figs. 13–18, 1920.

Syn. *Corynelia clavata* (L.), Sacc. in R. Pirotta, Osservazioni sopra alcuni funghi : Nuovo Giornale Botanico Italiano, 21, pp. 312–317, 1889.

On *Podocarpus latifolia*, Endl., Grahamstown, 13.7.1907, on fruit [493]; Natal, Medley Wood (Wood No. 3203) [11206]; Woodbush, Zoutpansberg District, 4.8.11, Doidge [1770]; St. James, Cape Province, 22.12.12, Pole Evans [5572]; Grahamstown, Burt-Davy [5580]; Kingwilliamstown, 19.6.14, Sun. [7815]; Keiskama Hoek, 26.3.15, District Forest Officer [8970]; Umtata, Cape Province, 13.5.16, Pegler [9739]; Deepwalls, Knysna District, 13.5.23, Doidge [17204].

On *Podocarpus elongata*, Hk., near Somerset East, MacOwan, Rabh. Wint. Fung. Eur. 3149 [3489]; Woodbush, Transvaal, March, 1910, Gray [885]; Berg River, near Wellington, Cape Province, 8.11.10, Doidge [1743]; Pirie Forest, Kingwilliamstown, 10.1.14, District Forest Officer [7354]; Kingwilliamstown, 7.2.14 (on fruits), District Forest Officer [8258]; Keiskama Hoek, 26.3.15, District Forest Officer [8971]; Keiskama Hoek, 30.5.15 (on fruits), Keet [14141]; Storms River, Humansdorp District, 15.5.23, Doidge [17165].

On *Podocarpus falcata*, Eshowe, Zululand, 30.1.12, Pole Evans [2027]; Kentani, Cape Province, July, 1913, Pegler [6900]; Ngome, Cape Province, 21.3.14, Foster [7410].

Stromata scattered, infrequently confluent, more or less definitely circular, 1–3 mm. in diameter, chiefly hypophyllous, but often amphigenous, caulicolous and fructicolous, not usually erumpent before the limitation of the perithecia, in section homogeneous and black or dark-brown, externally dull-black and minutely roughened, bearing finally a crowded cluster of 20–50 or more perithecia. Perithecia first making their appearance through the ruptured epidermis as hemispherical protuberances on the stroma, the ascigerous cavity being partially buried and pure white within. The young perithecium cartilaginous soon attaining a characteristically conical form, the apex smooth and shining, the base rough and dull, later protruding farther and developing a cylindrical neck, which becomes clavate by the pronounced enlargement of the tip. Perithecium at maturity approximately 1 mm. in length, brittle, somewhat dumb-bell shaped, usually bent in the narrow middle portion and appearing consequently inequilateral, this being especially pronounced in those individuals borne at the margin of the stroma. The swollen apex of the perithecium at maturity flattened and deeply cleft by one to several transversely running furrows, the resulting ridges usually breaking into scales, giving the apex of the perithecium a pronouncedly shaggy appearance, finally dehiscient along the line of the middle furrow by a wide and deep slit, the two lips pulling apart and usually recurving, thus exposing the ashy to brown inner wall. Asci ovate to clavate, long-stalked, thin-walled, evanescent, paraphysate, eight-spored, $20\text{--}26 \times 34\text{--}44 \mu$. Ascospores when young hyaline, smooth, and by mutual pressure polyhedral; at maturity spherical, brown, thick-walled, echinulate, unicellular, $9\text{--}14 \mu$ (mostly 12μ) in diameter. Pycnidia sometimes developed on the stroma; pycnospores elongated, $5\text{--}7 \times 2 \mu$.

126. *Calonectria capensis*, n. sp.

Parasitic on *Irene Podocarpi* on leaves of *Podocarpus elongata*, Storms River, Humansdorp District, 15.5.23, Doidge [17167].

Mycelium arachnoid, closely investing the mycelium of the *Irene* and forming a network between the hyphae of the host, dirty-white or yellowish to reddish, $1\text{--}3 \mu$ thick, creeping, densely and intricately branched and anastomosing, usually co-extensive with the *Irene* mycelium. Perithecia few, scattered, globose, $130\text{--}200 \mu$ diameter, membrane golden-yellow, cells irregular-polygonal, $7\text{--}10 \mu$ diameter. Perithecial setae rather numerous, similar in colour to perithecial membrane, erect or suberect, straight or curved, not tapering, apex blunt, septate, $100\text{--}130 \mu$ long, $5\text{--}6 \mu$ thick. Asci numerous, fasciculate, eight-spored, narrow-clavate or cylindrical, thin-walled, rounded at the apex, tapering to the base,

83–100 $\times$ 13–16 μ . Paraphyses numerous, filiform. Spores 5–6-septate when mature, hyaline, fusoid-clavate, straight or curved, occasionally subsigmoid, tapering more gradually to the lower end, both ends obtuse, not constricted at the septa, 50–54 $\times$ 6–6.5 μ .

Calonectria capensis, Doidge, n. sp.

Mycelium in mycelio *Irene Podocarpi* parasiticum, arachnoideum, ex hyphis pallide flavidulis v. rubris, 1–3 μ latis, ramosis compositum; perithecia pauca, sparsa, globosa, 130–200 μ diametro, flava, contextu parenchymatico e cellulis polygonis, 7–10 μ diametro, pilis rectis v. curvatis, obtusis, septatis, 100–130 μ longis, 5–6 μ latis; asci fasciculati, octospori, angusti-clavati v. cylindracei, apice rotundati, 83–100 $\times$ 13–16 μ , paraphysibus numerosis filiformis; sporae hyalinae, fusoidae-clavatae, rectae v. curvatae, utrinque attenuatae sed obtusae, haud constrictae, 50–54 $\times$ 6–6.5 μ .

Hab. in foliis *Podocarpi elongatae*, Storms River, Humansdorp District, 15.5.23, leg. Doidge [17167].

127. *Phragmosperma Marattiae* (P. Henn.), Th. et Syd.

Ann. Myc., XIV, pp. 450–451 (461).

Syn. *Micropellis Marattiae*, P. Henn., Hedwigia, 34, 1895; Sacc., Syll. Fung., XI, p. 382.

On leaves of *Marattia fraxinea*, Winters Kloof, Natal, 13.7.11, Doidge [1632].

This fungus is taken by Theissen and Sydow (loc. cit.) as the type species of the genus *Phragmosperma* of the group *eu-Montagnelleae*. The genus is characterized by single perithecia, which are half-immersed in the leaf-tissues, without a true stroma and without an ostiole, the upper wall eventually breaking down. Asci eight-spored, paraphysate; spores hyaline, with several transverse walls.

Phragmosperma Marattiae (P. Henn.), Th. et Syd., has single perithecia, which are closely crowded and rhomboid, forming groups up to 5 mm. diameter, which are interrupted by the nerves on which no perithecia are formed; perithecia, 130–140 μ diameter, the upper part breaking out to the leaf-surface, with a papilla but no ostiole, the upper wall eventually breaking down; the wall is brown and formed of small cells. Stroma wanting, but occasionally some contact hyphae may be detected between neighbouring perithecia. Asci clavate, rounded or thickened at the apex, sessile, eight-spored, paraphysate, 50–70 $\times$ 12–15 μ . Spores mostly parallel in a single bundle, straight or somewhat oblique, tapering slightly towards each end, at first with a single medial septum, later each cell becomes further septate, not constricted, hyaline, 30–38 $\times$ 3–4 μ .

128. *Phyllachora anthistiriicola*, Syd., n. sp.

Stromata in folio decolorato evoluta, in utraque folii pagina, sed praecipue in superiore visibilia, irregulariter distributa, mox plus minus solitaria vel irregulariter aggregata, mox et saepissime dense aggregata et paginam folii superiorum fere crusta aterrima obtegentia, singula $\frac{1}{2}$ –1 mm. longa, confluyendo haud raro longiora, in epiphylllo convexa, in hypophylo plana et saepe tantum e stromate sterili contraposito vel clypeo epidermali constantia; clypeo epiphylllo aterrimo, 20–30 μ crasso; loculi pauci in singulo stromate, 200–350 μ lati, 175–200 μ alti, plerumque $\frac{3}{4}$ folii crassitudinis rarius totam crassitudinem occupantes; asci cylindracei, 70–80 $\times$ 12–16 μ , paraphysati, octospori; sporae 1–2-stichae, oblongae, plerumque obtusae, continuae, hyalinae, 11–16 $\times$ 7–9 μ ; conidia simul praesentia, filiformia, recta vel curvata, hyalina, 12–15 $\times$ $\frac{1}{2}$ μ .

Hab. in foliis *Anthistiriæ imberbis*, Magaliesberg, Transvaal, 23.3.12, leg. Van der Bijl [2196].

129. *Phyllachora Chrysopogonis*, Syd., n. sp.

Stromata amphigena, per totum folium pallide decoloratum plus minus aequahiter dispersa, solitaria, vix confluentia, elliptica $\frac{1}{2}$ –1 mm. longa, convexula, 2–4 locularia. clypeo unilaterali aterrimo ca. 25 μ crasso; loculi applanati, 300–420 μ lati. 130–170 μ alti, $\frac{2}{3}$ folii

crassitudinis occupantes, parietibus brunneis, 15–20 μ crassis; asci cylindracei copiose paraphysati, 65–85 $\times$ 10–14 μ , octospori; sporae plerumque monostichae, oblongae, utrinque obtusae, continuae, hyalinae, 12–15 $\times$ 7–8 μ .

Hab. in foliis *Chrysopogonis monticolae*, Armandsvlakte, Vryburg, 25.10.15, leg. Pole Evans [9302].

130. *Phyllachora Digitariae*, Syd., n. sp.

Stromata in utraque folii pagina indistincte decolorata conspicua, elliptica, $\frac{1}{3}$ –1 mm. longa, raro confluentia, leniter convexa, parum nitidula, clypeo epidermali amphigeno 20–30 μ crasso, 1–4 locularia; loculi totam folii crassitudinem occupantes, 150–200 μ lati, usque 175 μ alti, pariete 20–25 μ crasso; asci cylindracei, breviter stipitati, 50–65 $\times$ 9–14 μ , copiose paraphysati; sporae monostichae vel in superiore asci parte distichae, late ellipsoideae usque subglobosae, continuae, hyalinae, 8–10 $\times$ 7–8 μ .

Hab. in foliis *Digitariae Smutsii*, Irene, Transvaal, 1922, leg. S. Smuts [17017].

131. *Phyllachora Doidgeae*, Syd., n. sp.

Stromata amphigena, irregulariter dispersa, discreta, elliptica, $\frac{1}{2}$ –1 mm. longa, convexula, 1–4 locularia, clypeo epidermali 30–40 μ crasso aterrimo; loculi totam folii crassitudinem occupantes, 300–400 μ lati, 200–300 μ alti, parietibus 10–15 μ crassis; asci cylindracei, 80–100 $\times$ 12–16 μ , copiose paraphysati; sporae monodistichae, oblongae, plerumque late rotundatae, continuae, hyalinae, 14–18 $\times$ 8–10 μ ; loculi conidiiferi simul praesentes aequales, sed plerumque minores, 175–250 μ lati, 140–170 μ alti; conidia oblonga vel oblongo-fusoida, utrinque obtusa vel leniter attenuata, continua, dilute fuscicula, 8–12 $\times$ 2–3 μ , basidiis nullis.

Hab. in foliis *Andropogonis cerasiaeformis*, Durban, 7.7.11, leg. Doidge [1612]; Edendale, Natal, 26.12.11, leg. Doidge [1998].

132. *Phyllachora Leptocarydii*, Syd., n. sp.

Stromata plerumque in utraque folii pagina visibilia, per folii superficiem plus minus aequaliter distributa, $\frac{1}{3}$ – $\frac{2}{3}$ mm. longa, in epiphylllo convexula in hypophylo plana et fere semper e stromate contrapposito sterili tantum constantia, pauci (1–3)-locularia, clypeo epiphylllo aterrimo, 20–30 μ crasso; loculi, 100–140 μ lati, 90–120 μ alti; circiter $\frac{2}{3}$ – $\frac{3}{4}$ folii crassitudinem occupantes, pariete tenui, 10 μ crasso; asci cylindracei, 50–60 $\times$ 7–11 μ , octospori, paraphysati; sporae monodistichae, ovatae vel ellipsoideo-ovatae, continuae, hyalinae, 7–10 $\times$ 5–6 μ .

Hab. in foliis *Leptocarydii Vulpiastri*, Sydenham, Natal, 1914, leg. Franks [7814].

133. *Phyllachora Winkleri*, Syd.

Ann. Myc., X, 1912, p. 80; XIII, 1915, p. 456.

On leaves of *Paspalum scrobiculatum*, Quelimane, Mozambique, 1908, Howard [648].

Stromata convex, elliptic or oblong, often confluent, 1–3 mm. diameter, with a rough surface, lying between the vascular bundles and the upper layer of cells; the space between the bundles and the upper epidermis being filled with a perpendicular, prosenchymatous, palisade stroma, which distends the leaf to double its normal thickness. Loculi not numerous, fusing with the epidermal clypeus; in transverse section they are 340 μ in diameter and 180–200 μ high; in longitudinal leaf-sections they reach a length of 700 μ ; the wall of the locus is sharply differentiated, brown, and in most cases formed of concentric layers of very small hyphae, 2½ μ thick. Asci cylindrical, 80–120 $\times$ 11–16 μ , briefly stipitate. Spores monostichous, broadly elliptic, one-celled, hyaline, 14–17 $\times$ 12–19 μ . Paraphyses numerous, thread-like.

This species was originally described as occurring on *Paspalum scrobiculatum*, in German East Africa.

134. *Phyllachora sanguinolenta*, Theiss. et Syd., var. *microspora*.

Ann. Myc., XIII, 1915, p. 455.

On leaves of *Panicum minus*, Bloemfontein, April, 1917, Potts [11307].

Stromata thickly scattered on both leaf-surfaces, convex, somewhat shiny, elliptic, $\frac{3}{4}$ –1 mm. long. Epidermal clypeus $20\ \mu$ thick, on one or both sides of the leaf. Where the stroma is on only one side of the leaf, the loculi are elliptic, occupying two-thirds of the thickness of the leaf; they are numerous, and there is a stroma connecting the upper parts of the loculi. When two stromata occur on opposite leaf surfaces the loculi only reach to the middle of the leaf, and are more spherical, $240\ \mu$ diameter; the locular walls are thin and brown. Asci cylindrical, paraphysate, briefly stipitate, $60\text{--}70 \times 10\text{--}14\ \mu$. Spores obliquely monostichous, sometimes distichous, elliptic, hyaline, one-celled, $8 \times 5\ \mu$.

This is another tropical African species, originally collected on an undetermined grass in the Congo.

135. *Phyllachora Brachystegiae*, n. sp.

On leaves of *Brachystegia* sp., December, 1917, Salisbury, Rhodesia, Eyles [11680].

Stromata minute, epiphyllous, circular or broadly elliptic, black, convex, somewhat shiny, penetrating through the leaf and becoming visible on the lower surface. Each stroma has a single loculus, which is flattened-spherical, $350\text{--}370\ \mu$ diameter, $240\text{--}290\ \mu$ high, occupying the entire thickness of the leaf. Wall of the loculus light-brown, about $10\ \mu$ thick. Clypeus developed in both the upper and the lower epidermis, $400\text{--}500\ \mu$ in diameter. Asci paraphysate, cylindrical, eight-spored, $100\text{--}120 \times 13\text{--}16\ \mu$. Spores monostichous, occasionally subdistichous, one-celled, hyaline, ellipsoid, somewhat more rounded at one end than the other, $16.5\text{--}17 \times 6.5\text{--}7\ \mu$; conidia hyaline, filiform, curved, $16\text{--}20 \times 1\ \mu$.

Phyllachora Brachystegiae, Doidge, n. sp.

Stromata minuta, rotundata v. late elliptica, atra, convexa parum nitidula, in hypophyllo etiam perspicua; loculum tantum unicum, subglobosum, $350\text{--}370\ \mu$ latum, $240\text{--}290\ \mu$ altum, totem folii crassitudinem occupans, pariete tenue $10\ \mu$ crasso, clypeo epidermale amphigeno; asci paraphysati, cylindranei, octospori, $100\text{--}120 \times 13\text{--}16\ \mu$; sporae monostichae v. distichae, hyalinae, continuatae, ellipticae, $16.5\text{--}17 \times 6.5\text{--}7\ \mu$; conidia hyalina, filiformia, curvata, $16\text{--}20 \times 1\ \mu$.

SOUTH AFRICAN GRAMINEAE.

GRASSES OF THE TRANSVAAL AS REPRESENTED IN THE NATIONAL HERBARIUM.

By SYDNEY M. STENT.

THE Transvaal may be divided broadly into three climatic areas each with various edaphic conditions and a corresponding flora. The South-Western and Western portion takes in part of the Karroo and the sand-veld of Bechuanaland. It consists of grass and low scrub with scattered thorn trees. The climate is generally hot and dry; the average rainfall is from 15 to 25 inches. The dominant grasses of this region are xerophytic species of *Eragrostis* (*E. Lehmanniana*, *E. Atherstonii*, *E. obtusa*, *E. echinochloidea*), *Sporobolus Ludwigii*, *S. tenellus*, *Aristida* spp., *Themeda triandra*, *Digitaria eriantha*, var. *stolonifera*, *Panicum coloratum*, *Brachiaria nigropedata*, *Antheplora pubescens*, *Fingerhuthia africana*, *Enneapogon* sp. and *Schmidtia* sp. The two latter are peculiarly characteristic of this region. Northwards to the Limpopo and eastwards, passing above the Magaliesberg, to the Drakensberg is the warm and moist tropical and sub-tropical region; a great part of this is bush veld with occasional stretches of open country developing into the savanna of the north. The annual rainfall of this region averages from 15–35 ins., except along the Drakensberg range, where it rises to over 70 inches. The tropical genera of the Andropogoneae are abundantly represented, the tall, coarse tambookie grasses and the sorghums, the dek-gras or thatching grasses, as well as many of the finer species. Broad-leaved species of the Paniceae are dominant in the bushveld area—*Panicum maximum* (Guinea grass), *Brachiaria brizantha*, *Digitaria* spp., *Setaria sulcata*, *Pennisetum cenchroides*, etc. The third region is what is known as the High Veld, which extends along the southern border, taking in parts of Potchefstroom, Heidelberg, and Standerton, and north as far as Middelburg and the Magaliesberg range. The winters in the High Veld are cold and dry, and it is subject to severe frosts; the average summer rainfall is from 25 to 35 inches. This region is practically all grass-veld, flat and treeless. *Panicum natalense* is plentiful. *Panicum laevifolium*, *Setaria nigristrois*, *Themeda triandra*, *Eragrostis chalcantha*, *E. curvula*, *E. plana*, *Sporobolus indicus*, *Elionurus argenteus* (a widely distributed "sour" grass), *Koeleria cristata*, and *Cynodon dactylon* are among the dominant species.

For arrangement and numbering of genera in the following list, I have followed Dalla Torre and Harnis in their *Genera Siphonogarum*. Where a genus has been created or revived since the publication of that work, it has been placed next to that to which it shows the closest affinity and given the same number with a distinguishing letter.

* Dr. Stapf's work on the *Gramineae* of Tropical Africa (Prain's Flora of Tropical Africa, Vol. IX), which is not yet complete, has been used, as far as possible, as a basis for nomenclature. No synonymy is given except in cases where the name given differs from that in the Flora Capensis; the synonym employed in this earlier work is then added in brackets. References to original descriptions are only given when the species is not described in the Flora Capensis.

* Dr. Stapf, in his recent and not yet completed work on the grasses of Tropical Africa, has revived a number of old genera and created a number of new ones. This paper is not the place for a critical review of this work, and I have, for present purposes, accepted it as a basis on which to build up my classifications, though I do not in all cases agree with Stapf's sub-divisions.

There is no doubt for instance that the genera *Andropogon* and *Panicum* as defined by him in his earlier work in the Flora Capensis were altogether too unwieldy, and contained a number of very heterogeneous types, but in splitting these up into separate entities he has, I think, gone to the other extreme and he has based some of his small genera on characters which I find to be inconstant or mutable, and often unimportant. This question will, however, be discussed in a later paper.

Local names are given where they are known. For the agricultural notes I am indebted to Mr. H. A. Melle, except where otherwise stated.

A revision of the South African species of the genus *Eragrostis* will be undertaken later. For the purposes of this present list the species not described in the Flora Capensis are identified by letters only.

Dr. J. Burtt-Davy, for ten years prior to Union, Government Botanist and Agrostologist of the Transvaal, is responsible for the collection of the greater number of Transvaal grasses in the National Herbarium. *Eragrostis Burtt-davii* was named in his honour. Mr. E. E. Galpin, formerly of Queenstown, but now of Mosdene, Naboomspruit, an enthusiastic collector, has of recent years contributed very considerably to our knowledge of the grasses of the Waterberg. The genus *Mosdenia* was so named after his farm, where it was collected; he has also been responsible for the collection of a number of new species, some of which bear his name.

Lastly, I make my acknowledgments to General J. C. Smuts, world-famous statesman and Prime Minister of the Union of South Africa, who finds time in his hours of ease (and he has not many of them) to interest himself in the study of botany as represented in the flora of South Africa. Being a farmer as well as a politician, he is most keenly interested in the grasses, and applies himself to stimulate and assist, in South Africa, more extensive research work on this most important order.

Digitaria Smutsii and *Sporobolus Smutsii* are named in his honour.

The localities recorded in the following list are, with few exceptions, from specimens in the National Herbarium of South Africa, all of which have been examined by me. The exceptions are indicated by an asterisk (*) after the number.

I have included in the key all the South African genera, instead of confining it only to those that occur in the Transvaal and narrowing the usefulness of the key to one Province.

The key to the species will be prepared later. At the present time there is such confusion between those of some of the larger genera that, until these have been revised, it is impossible to work out a satisfactory key to the species.

KEY TO THE GENERA OF SOUTH AFRICAN GRASSES.

1. Spikelets not or dorsally compressed, falling from the pedicels entire, never more than 2-flowered, perfect floret always uppermost 2
 Spikelets compressed laterally or scarcely compressed, 1-many flowered, glumes usually persistent or subpersistent on the rachis after the fall of the flower—or if the spikelets fall entire, then not consisting of 2 heteromorphous florets of which the lower is ♂ or barren 46
2. Valves and pales much more delicate in texture than the glumes, valves awned (except in *Urelytrum*, *Hemarthria*, *Elionurus*, and *Imperata*) 3
 Valves similar in texture to the glumes or firmer, never entirely hyaline, occasionally mucronate, but seldom long awned. (In *Antheophora* the lower glume is rigidly coriaceous and longer than the valves, and in *Melinis* the barren valve is finely, sometimes long awned) 24
3. Spikelets in pairs at least in the upper part of the raceme, one sessile or shortly pedicelled and one on a longer pedicel 4
 Spikelets solitary, on filiform pedicels, arranged in lax racemes on the branches of a long narrow panicle XX. (¹) **Cleistachne**.
4. Spikelets of a raceme all alike in sex and form 5
 Spikelets differing in sex or form or both 9
5. Spikelets all pedicelled 6
 Spikelets one sessile and one pedicelled 8
6. Spikelets arranged in long solitary, binate or digitate racemes terminating the culms.
 VIII. **Trachypogon**.
 Spikelets racemosely arranged on the branches of a panicle 7
7. Panicle contracted, spike-like, dense, with long silvery-silky hairs; spikelets not awned I. **Imperata**.
 Panicle much branched, large, often somewhat contracted but not spike-like, greyish or tawny villous, spikelets awned II. **Miscanthidium**.
8. Racemes binate or digitate or subdigitate at the ends of the culms or the branches, villous with greyish or tawny hairs, spikelets awned III. **Eulalia**.
 Racemes solitary, terminating the culms and their branches, joints and pedicels fused, forming semi-cylindrical internodes, which are excavated on the inner face for the reception of the sessile spikelets. Spikelets not awned.
 VII. **Hemarthria**.
9. Racemes solitary (at least in the South African species), terminating the culms and their branches 10
 Racemes 2 or more 16
10. Racemes enclosed or partially enclosed in spathes 11
 Racemes not subtended by spathes 13

(¹) The numbers before the names refer to the order of the genera in the following list. Those without numbers have not been recorded within the limit of the Transvaal.

11. Lower glume of sessile spikelet with involute margins, not or obscurely keeled towards the apex..... 12
 Lower glume of sessile spikelet with sharply inflexed margins; 2-keeled, keels running into teeth or mucros. Racemes slender; erect grasses usually tinged with red or brown. Much used for thatching.....XIV. **Schizachyrium.**

12. Racemes bearing 6-8 fertile, short-awned spikelets, almost entirely enclosed in the conspicuous brightly coloured, permanently cymbiform, spathes.

XVIII. **Monocymbium.**

Racemes much contracted, bearing 1 (rarely 2) fertile long-awned spikelet, the lower sessile florets approximating so as to form an involucre round the upper one. "Rooigras".....XXI. **Themeda.**

13. Lower 3-10 pairs of spikelets homogamous, ♂, the sessile of the upper pairs ♀ or ♂, usually becoming dark brown, with a long hirsute twisted awn. "Speargrass."

IX. **Heteropogon.**

All pairs of spikelets heterogamous..... 14

14. Valve of fertile floret bilobed and awned. Lower glume of sessile spikelet bidentate with winged keels, pedicelled spikelets converging over the convex sides of the raceme. Raceme curved, spikelets up to 16 mm. long. Tropical grass.

V. **Sehima.**

Valves of fertile floret entire, not awned..... 15

15. Lower glume of pedicelled spikelet narrowed into a long squarrose awn, raceme long and rigid. A coarse sour grass.....VI. **Urelytrum.**

Lower glume of pedicelled spikelet not long awned, racemes delicate and silvery villous. Aromatic grass with narrow folded and usually flexuous leaves.

IX. **Elyonurus.**

16. Pedicelled spikelets ♀, racemes arranged in a densely ferruginous hirsute panicle. **Eriochrysis.**

Pedicelled spikelets ♂ or neuter or suppressed..... 17

17. Racemes not subtended by a spathe..... 18

Racemes subtended by a spathe..... 22

18. Racemes digitate or subdigitate..... 19

Racemes panicle..... 21

19. Fertile valve bilobed, racemes binate, digitate, or fascicled, joints and pedicels cuneate, hollow and convex on the back.....IV. **Ischaemum.**

Fertile valve stipiform, joints and pedicels filiform..... 20

20. Joints and pedicels solid, racemes always with 1-2 homogamous pairs of spikelets at the base.....XIII. **Dicanthium.**

Joints and pedicels longitudinally grooved and hyaline in the groove, sometimes with the lowest pair of spikelets homogamous.....XII. **Amphilophis.**

21. Racemes reduce to three spikelets, one sessile, and two pedicelled which terminate the branches of the culms, trio falling together from the thickened bearded tips of the peduncles.....XI. **Chrysopogon.**

Racemes lax, few jointed, arranged on the branches of large usually open panicles **X. Sorghum.**

22. Lower glume of sessile spikelet with sharply inflexed margins, 2-keeled..... 23
 Lower glume of sessile spikelet with involute and more or less rounded margins, awns
 hirsute, racemes binate on slender peduncles, subtended by a reddish spathe
 and collected into compound panicles.....XVII. **Hyparrhenia.**
23. Pairs of spikelets usually all heterogamous : racemes digitate or subdigitate, termin-
 ating the culm and its branches, peduncles usually long exerted from their
 narrow inrolled spathes.....XV. **Andropogon.**
- Lowest pair of spikelets on the lower raceme, or on both racemes, always homogamous;
 awns not hirsute ; racemes binate, partially enclosed, or not very long exerted
 from the reddish cymbiform spathe. Racemes collected into compound spatheate
 often dense and contracted, panicles. Mostly aromatic grasses.
 XVI. **Cymbopogon.**
24. Inflorescence of digitate, subdigitate, or racemously arranged spike-like racemes 25
 Inflorescence, an open, or contracted, or cylindrical, dense and spike-like panicle 37
25. Racemes very slender, digitate or subdigitate or less often whorled, or scattered,
 on a long common axis ; spikelets muticous, nerves of the lower valve straight
 and parallel, margins of the upper valve flat and hyaline, lower glume minute
 or suppressed.....XXXV. **Digitaria.**
- Racemes digitate, or sub-digitate, or racemously arranged on a long common axis ;
 spikelets often mucronate, lower glume well developed, nerves of the valve and
 glumes curved, margins of upper valve firm and inrolled..... 26
26. Racemes digitate or subdigitate, nerves of the upper glume and lower valve 5, the
 outer nerves approximate and close to the margin ; upper valve mucronate ;
 upper glume and lower valve somewhat indurated ; keels of the lower glume
 sometimes fimbriate, winged ; spikelets 4-6 mm. long..XXVIII. **Alloteropsis.**
- Racemes racemously arranged or paniced on an elongated common axis..... 27
27. Racemes very short (sometimes reduced to 1 spikelet), more or less sunk in the
 hollows on the face of a dorsally flattened, rather wide, common rachis. Creeping
 or prostrate perennial. The coarse couch grass of the Cape Province,
Stenotaphrum.
- Racemes variously arranged on a more or less triquetrous, or terete, rarely flattened,
 narrow, common rachis..... 28
28. Lower glume of the spikelet very much reduced or quite suppressed..... 29
 Lower glume well developed, from less than half to equalling the spikelet.... 30
29. Lower glume reduced to a narrow membranous, or hyaline scale round the swollen
 bead-like lowest node of the spikelet.....XXX. **Eriochloa.**
- Lower glume usually quite suppressed, rarely a minute scale, lowest node of spikelet
 not swollen and bead-like, upper glume convex, lower valve flat.
 XXIX. **Paspalum.**
30. Spikelets with no definite orientation as regards the rachis..... 35
 Spikelets definitely orientated with regards to the rachis..... 31
31. Spikelets with the back of the fruiting valve towards the rachis (abaxial)..... 32
 Spikelets with the face of the fruiting valve towards the rachis (adaxial). Spikelets
 closely appressed, typically 2-seriate, sometimes becoming 1-seriate.
 XXXII. **Brachiaria.**

32. Upper valve obtuse, abruptly mucronate, or aristulate; glumes mucous or cuspidate.....XXXI **Urochloa.**

Upper valve apiculate or obtuse; glumes often distinctly cuspidate, or, like the valves, with compressed callous tips..... 33

33. Fruiting floret gaping at the tip, owing to the slight recurving of the palea and valve..... 34

Upper valve embracing the palea all along, glumes and lower valve mucronate, or with distinct hirsute awns, that of the lower glume the longest.

XXXVIII. **Oplismenus.**

34. Glumes and valves with compressed callus tips or crests..... **Acroceras.**

Glumes and valves mucronate or shortly awned or at least cuspidate.

XXXIII. **Echinochloa.**

35. Spikelets not subtended by an involucre of one or more bristles, softly hairy.

Leucophrys.

Spikelets subtended by an involucre of bristles..... 36

36. Involucre usually reduced to 1 bristle; spikelets in long slenderly peduncled spike-like racemes, falling with the involucre. Glumes usually reduced to very small, delicate, nerveless, or 1-nerved scales.....XL **Pennisetum.**

Spikelets arranged racemously or in fascicles, on the branches and branchlets of a narrow spreading or contracted panicle, often some of the spikelets without a subtending bristle, bristles always persistent, glumes herbaceous, nerved.

XXXIX. **Setaria.**

37. Inflorescence a more or less dense cylindrical spike-like panicle..... 38

Inflorescence an open or contracted panicle..... 42

38. Spikelets single or in clusters, subtended by 1 or more bristles, or surrounded by an involucre of spinous bracts, or with the outer glumes indurated and forming a spurious involucre..... 39

Spikelets not subtended by an involucre of any sort. Upper glume with a curved or gibbous or saccate back, lower valve similar, but with a flatter back, hence the spikelet more or less oblique in profile. Spikes usually very dense and dark coloured..... **Sacciolepis.**

39. Spikelets subtended by 1 or more bristles..... 40

Spikelets subtended by involucre formed of spinous bracts, or of the lower glumes 41

40. Bristles of the involucre persistent; glumes well developed and nerved; upper valve finely pitted or rugose and very convex.....XXXIX. **Setaria.**

Bristles of the involucre deciduous with the spikelets, glumes small, delicate, scale-like, often nerveless; upper valve more or less resembling the lower, smooth, and not strongly concave.....XL **Pennisetum.**

41. Involucre composed of hard spinous bracts and deciduous with the enclosed cluster of spikelets, forming a spiny "burr".....**Cenchrus.**

Clusters of spikelets with the upper glumes turned towards the centre, the acuminate squarrose outer glumes contiguous at the base, and forming a spurious involucre round the deciduous cluster.....XXII. **Anthephora.**

42. Tips of upper glume and lower valve notched or bilobed, often with a fine awn; lower glume minute..... 43

Tips of glumes and valves entire, lower glume well developed (small and hyaline in *Sacciolepis*)..... 45

43. Upper glume and barren valve 5-nerved, nerves faint and anastomosing below the obtuse tips..... 44
 Upper glume 7-, barren valve 5-nerved, nerves prominent and not anastomosing below the acute tips, barren valve usually finely awned.....XXXVII. *Melinis*.
44. Upper glume and barren valve gibbous and long silky hairy....XXXVI. *Tricholaena*.
 Upper glume and barren valve not gibbous, glabrous, or at most shortly hairy.
 XXXVII. *Xyochlaena*.
45. Upper glume saccate (hence spikelets very gibbous), with 9-11 prominent rib-like nerves, lower glume small, hyaline.....*Sacciolepis*.
 Upper glume convex not gibbous, nerves not hardened and rib-like; lower valve equally convex, hence spikelet more or less symmetrical in profile.
 XXXIV. *Panicum*.
46. Leaf blades never transversely veined, in South African species, nor articulated on the sheath..... 47
 Blades transversely veined in South African species..... 139
47. Spikelets 1-flowered without barren valves or reduced florets above or below it (at least in South African species)..... 48
 Spikelets more than 1-flowered..... 69
48. Inflorescence an open, contracted, or more or less spike-like panicle 49
 Inflorescence of solitary, binate, or umbellate spikes 62
49. Spikelets disarticulating below the glumes and falling entire..... 50
 Spikelets disarticulating above the glumes..... 51
50. Spikelets in clusters or sometimes solitary in a dense cylindric spike-like panicle, the upper glume turned away from the rachis and with lines of short hooked spines on the prominent nerves.....XXIII. *Tragus*.
 Spikelets large, strongly laterally compressed, on the slender branches of an elongated spreading or contracted panicle, glumes scale or bristle-like. Valves strongly nerved.....XLII. *Oryza*.
51. Valves unawned..... 52
 Valves awned..... 55
52. Glumes well developed..... 53
 Glumes reduced to an obscure hyaline, entire or 2-lobed rim; valve 5-nerved subcartilaginous, keel and margins rigidly ciliate.....XLIII. *Leersia*.
53. Glumes equal or subequal, floret shorter than the glumes, spikelets light green 54
 Lower glumes usually smaller, often much smaller, than the upper, floret longer than the glumes, spikelets grey or greyish green, seldom gaping early.
 L. *Sporobolus*.
54. Glumes closely 3-nerved at the base, valve faintly 7-nerved, finely silky villous; pale subequal to the valve; ligule a fringe of hairs. A very delicate, much dwarfed perennial with a scanty lax panicle.....*Poagrostis*.
 Glumes 1-nerved, valve 3-nerved, glabrous or hairy (not silky), pale shorter than the valve, ligule membranous.....LI. *Agrostis*.
55. Awn from the back of the valve..... 56
 Awn from the tip or from just below..... 58

56. Callus small and long hairy, hairs usually exceeding the valve. Spikelets about 6 mm. long, straw coloured, shining, and crowded into a dense linear or linear oblong, spike-like panicle.....LII. **Calamagrostis**

Callus small, glabrous, or shortly hairy or minutely bearded..... 57

57. Glumes plumose, bristle tipped, panicle spike-like, compact, oblong, softly hairy
.....Lagurus.

Glumes glabrous, often scaberulous, not bristle tipped; panicle glabrous, spike-like, or open and lax.....LI. **Agrostis**.

58. Glumes awned; valve shorter than the glumes, very thin to hyaline; panicle contracted, spike-like, and plumose from the fine awns and bristles....Polypogon.

Glumes occasionally rather long mucronate, but never awned..... 59

59. Awn 3-branched or 3-partite (very rarely simple), ligule a fringe of hairs.

XLVII. **Aristida**.

Awn simple, ligules membranous..... 60

60. Awns rather coarse, kneed and strongly twisted below the knee, floret cylindric or linear oblong.....Stipa.

Awn straight..... 61

61. Floret shorter than the glumes; valve convolute and more or less flattened from the back, 5-nerved, with minutely bilobed tips; awn from the sinus, caducous.

XLVIII. **Oryzopsis**.

Floret exerted from the glumes; valve involute and rounded from the back, 3-nerved and continuous with the awn.....XLIX. **Pseudobromus**.

62. Spikes solitary..... 63

Spikes binate or digitate..... 68

63. Glumes linear or linear lanceolate tipped by an awn, or subulate, or bristle-like 64

Glumes not awned..... 65

64. Spikelets in groups of 3 at the nodes of the rachis of a dense bristly spike, the two lateral ♂ or barren, the middle one ♀; valve herbaceous, chartaceous, awned.

Hordeum.

Spikelets very small, solitary, on the rachis of a slender spike, awns of glumes up to ten times as long as the glume, and usually capillary and purplish, valve smaller than the glumes and delicately hyaline.....XXV. **Perotis**.

65. Spikelets more or less immersed in the hollows of the rachis of a simple terminal spike, lower glume reduced or altogether suppressed..... 66

Spikelets not immersed in hollows, lower glume present..... 67

66. Upper glume 5-7-nerved, very rigid and acute; valve hyaline acute, glabrous, finely 3-nerved; callus obsolete.....Lepturus.

Upper glume 1-3-nerved, rigid except at the hyaline margins, valve hyaline, truncate or minutely 3-toothed, 3-nerved, and hairy along the nerves, callus bearded.

LXXXI. **Oropetium**

67. Spikelets arranged unilaterally on a flattened rachis, rachilla disarticulating above the glumes, which are persistent or deciduous.....LVII. **Michrochloa**.

Spikelets arranged singly on minutely tomentose notches in spirals round a ribbed and scabrid axis, from which they fall entire. Glumes subequal, lower boat-shaped and falcate.....XXIV. **Mosdenia**.

68. Mature spikelets tardily disarticulating below the glumes, valve 1-nerved. Spikelets 12-14 mm. long on a triquetrous excurrent rachis. Spikes 1-3, yellowish. Maritime grasses.....**Spartina.**
- Mature spikelets readily disarticulating above the glumes, the lower glume persistent. Valve 3-nerved, spikelets about 2 mm. long on a slender-keeled not excurrent, rachis. Spikes digitate or subdigitate.....**LVIII. Cynodon.**
69. Spikelets arranged in secund spikes or racemes..... 70
Spikelets in open or contracted or spiciform panicles, or in spikes or racemes, but not secund..... 84
70. Spikes solitary, terminating the culms..... 71
Spikes in umbels or variously arranged on a more or less elongated common axis 77
71. Valves awned..... 72
Valves not awned..... 73
72. Lower glume symmetrical, thin, keeled; upper glume 2-3-nerved with a stiff short awn from the middle; valve entire and awned just below the tip. Spikelets in dense unilateral, usually curved spikes.....**LX. Ctenium.**
- Lower glume inequilateral, one side dilated below the middle into a tooth, lobe, or wing. Upper glume entire, or notched below the mucronate or apiculate tip, but not awned. Valve 2-fid and awned from the sinus or 4-fid with the outer lobes awned. Spikelets rather remote on a slender furrowed rachis.
LXII. Tripogon.
73. Valves 7-nerved, navicular, finely tomentose in the lower half, with clavate hairs; spikelets 2, ranked.....**Brizopyrum.**
Valves 1-3-nerved..... 74
74. Valves minutely bilobed, spikelets 2-flowered, dorsally and obliquely compressed, glabrous, lower floret hermaphrodite.....**LVII. Microchloa.**
Valves entire..... 75
75. Upper florets crowded in a club-shaped body not exceeding the hermaphrodite floret and enveloped by the valve of the lowest male floret; valves white and densely ciliate along the nerves.....**LIX. Harpechloa.**
Upper florets not as above..... 76
76. Spikelets in spike-like panicles that terminate the culms and their branches; glumes not awned, keels cartilaginous, muricate pectinate. Ligule a line of hairs.
Prionachne.
- Spikelets unilateral on short slender spreading or deflexed spikes that are racemously arranged on an elongated common axis. Glumes awned, smooth. Ligule membranous.....**LXIII. Dinebra.**
77. Spikes in umbels..... 78
Spikes arranged on a common axis..... 81
78. Valves awned..... 79
Valves not awned..... 80
79. Upper glume mucronate or more often with a curved awn, tips of the rachis barren and forming a sharp mucro.....**LXV. Dactyloctenium.**
Upper glume obtuse or acute, and with a straight bristle or mucro, tips of the rachis not bare.....**LXI. Chloris.**

80. Glumes strongly keeled, 3-5-nerved, keel crested, spikelets greenish, 3-6-flowered.
 LXIV. **Eleusine**.
 Glumes rounded or flat on the back, upper shortly and obtusely bilobed, spikelets brownish, 2-flowered.....LXI. **Eustachys**.
 [= *Chloris petraea*, Thunb., in list.]
81. Spikes forming a compound spike..... 82
 Spikes or racemes arranged on the branches of a panicle..... 83
82. Glumes and barren valves 1-nerved, fertile valves 5-nerved; styles very short.
 Tetrachne.
 Glumes 3-5-nerved, barren valves 6-8 nerved, fertile valves prominently 9-11-nerved; styles very long.....Entoplocamia,
83. Valves rigidly membranous, acuminate, quite glabrous; panicles long and narrow with spreading, more or less curved, branches.....LXXIII. **Pogonarthria**.
 Valves obtuse, with broad hyaline tips and finely silky side nerves; branches of the panicle few and erect.....LXXIV. **Diplachne**.
84. Ligule a fringe of hairs..... 85
 Ligule membranous, scarious, or hyaline..... 105
85. Spikelets 2-flowered..... 86
 Spikelets more than 2-flowered..... 93
86. Fertile valve awnless, sometimes with a bristle-like mucro..... 87
 Fertile valves awned..... 89
87. Panicle open with fine branches and shining straw-coloured or purple-tinged spikelets, or if contracted into a spike-like panicle, then the lowest spikelets not consisting of a few empty glumes, and the valves not navicular with a submarginal line of hairs on each side.....Achneria.
 Panicle densely spiciform..... 88
88. Spiciform panicle dense, compact, cylindric, never interrupted, greyish-green, often tinged with purple, usually from 2½-5 cm. long; spikelets strongly laterally compressed, lowest spikelets consisting of a few empty glumes; glumes long and softly or rigidly ciliate along the keels.....LXXI. **Fingerhuthia**.
 Spiciform panicle smaller with fewer less strongly compressed spikelets, often much interrupted, no empty glumes at base of spike, glumes usually hispid with coarse tubercle based hairs.....Lasiochloa.
89. Lower floret awnless, ♂ or barren..... 90
 Both florets perfect..... 92
90. Spikelets solitary on distinct pedicels..... 91
 Spikelets in clusters of threes, rather large—up to over 1 inch long; valve of upper florets 2-lobed, with a coarse-kneed and twisted awn from the sinus
 LV. **Tristachya**.
91. Spikelets small, never more than 5 mm., valve of upper floret 2-setose, minutely 2-toothed or entire, awn sometimes reduced or absent....XXVI. **Arundinella**.
 Spikelets from 4 mm. to 3-25 cm., upper valve always distinctly 2-toothed or 2-lobed, awn always kneed.....LVI. **Tricopteryx**.
92. Ovary obovoid with a very dense, deciduous tomentum, consisting of branched hairs, at the top. Spikelets 14-24 mm. long.....Pentameris.
 Ovary oblong, glabrous, spikelets 2-12 mm. long.....Pentastichis.
93. Fertile valves lobed..... 94
 Fertile valves entire or minutely lobed, notched or toothed..... 98

94. Fertile valves 2-lobed..... 95
 Fertile valves 4-9-lobed..... 97
95. Spikelets deciduous with the bearded upper part of the pedicel; rachilla tough above the glumes, but readily disarticulating between the valves; awn from the sinus, kneed. Perennials, with flat soft leaves and narrow contracted panicle.
Chaetobromus.
 Spikelets disarticulating above the glumes and between the valves..... 96
96. Glumes not keeled, prominently nerved on the back, greenish with white hyaline margins; awn from the sinus of the 2-lobed valve, small and straight. Small annuals..... **Schismus.**
 Glumes keeled, often distinctly but not prominently nerved, usually straw-coloured and shining; awn usually kneed and twisted; hairs of the valve often gathered into tufts or lines. Perennial, seldom annual..... **Danthonia.**
97. Fertile floret 1; valve cleft into 9 subulate awn-like lobes. LXVIII. **Enneapogon.**
 Fertile florets, 3-5; valves cleft into four hyaline lobes, alternating with five straight awns from the sinuses..... LXIX. **Schmidtia.**
98. Fertile valves not awned..... 99
 Fertile valves awned..... 104
99. Valves 3-nerved..... 100
 Valves more than 3-nerved..... 102
100. Florets usually numerous and far exserted from the glumes; glumes and valves entire, awnless, side nerves of the latter distant from the middle nerve and often short; spikelets usually panicle..... LXXV. **Eragrostis.**
 Florets not, or shortly, exserted from the glumes; spikelets in a dense compact cylindric spike-like panicle..... 101
101. Spikelets jointed on and deciduous from the pedicels; glumes long and softly or rigidly ciliate; spikelets at the base of the panicle reduced to empty glumes.
 LXXI. **Fingerhuthia.**
 Spikelets disarticulating above the glumes and between the valves, purplish, and villous from soft grey hairs; no empty glumes at base of spike
 LXXVII. **Stiburus,**
102. Glumes 1-nerved..... LXXI. **Fingerhuthia.**
 Glumes 4-7-nerved..... 103
103. Spikelets 2-4-flowered, florets not exserted from the glumes; panicle spiciform, dense or sometimes with spiciform branches; spikelets mostly hispid from short coarse tubercle-based hairs..... **Lasiochloa.**
 Spikelets 4-9-flowered, florets distinctly, often far exserted from the glumes.
Brizopyrum.
104. Panicle ovoid, compact, very small, embraced at the base by the uppermost sheath, deciduous from the terminal node, culms 3-6 in. long..... **Urochlaena.**
 Panicle of subligitate spiciform branches; florets about 6, the lowest one or two ♂, the upper ones gradually reduced and embracing each other, forming a tuft of awns..... LXVI. **Lophacme.**
105. Spikelets always with 2 florets..... 106
 All or many of the spikelets with more than 2 florets..... 109
106. Spikelets of two kinds, the fertile surrounded by the sterile..... **Lamarkia.**
 Spikelets all alike..... 107

107. Spikelets sessile, imbricate in a dense cylindric spike.....**Secale.**
Spikelets in an open or contracted panicle..... 108
108. Spikelets disarticulating from the tips of the pedicels; glumes pectinate ciliate on the keels, the lower 1-, the upper 3-, nerved; panicle usually contracted, occasionally spike-like.....**LIII. Holcus.**
Spikelets disarticulating above the glumes and between the valves; glumes not pectinate ciliate on the keels, both 1-nerved; panicle delicate, open, with capillary branches.....**Aira.**
109. Valves 3-nerved..... 110
Valves more than 3-nerved..... 115
110. Lowest floret ♂ or barren, the rest ♀; spikelets enveloped by long hairs that spring from the long callus; panicle large, lax, and silky.....**LXXII. Phragmites.**
Lowest floret ♀, upper ♂ barren or reduced..... 111
111. Valves minutely toothed, or notched or occasionally entire, but not distinctly lobed 112
Valves lobed..... 114
112. Upper glume 3-nerved, lower 1- or 3-nerved.....**LXXVI. Koeleria.**
Upper and lower glumes 1-nerved..... 113
113. Valves truncate with 4 minute hyaline teeth, mid-nerve excurrent into a fine bristle, callus slender and acute.....**Leptocarydon.**
Valve 2-toothed or minutely notched or entire, callus very small or wanting.
LXXIV. Diplachne.
114. Valves shortly 2-lobed, side nerves not excurrent, callus small hairy; panicle spreading or contracted, with simple branches on which the subsessile or shortly pedicelled spikelets are appressed and rather far apart.....**LXVII. Crossotropis.**
Valves 3-lobed, the middle lobe more or less 2-fid and awned from the sinus, the side nerves also excurrent into fine awns; callus slender, acute, and bearded.
LXX. Triraphis.
115. Spikelets enveloped by long hairs that spring from the back of the valve; panicle large compound.....**LXXIIA. Arundo.**
Spikelets not enveloped by long hairs..... 116
116. Lowest florets of the spikelet ♂ or barren, or reduced..... 117
Florets all perfect or uppermost reduced (in *Cynosurus* this applies only to the terminal spikelets of the fascicle)..... 120
117. Lower valves much larger than the upper fertile valve..... 118
Lower empty valves reduced to minute scales..... 119
118. Lower male or barren valves awned from the back, membranous and equal. A sweet scented grass.....**XLVI. Anthoxanthum.**
Lower barren valves awnless or awned from the tip, cartilaginous..**XLIV. Ehrharta.**
119. Glumes reduced to minute scales or to an obscure, entire, or bilobed rim. A weak flaccid grass.....**XLI. Potamophila.**
Glumes large, enclosing the valves; leaves flat; panicle stiff and more or less spici-form.....**XLV. Phalaris.**
120. Lower 1-3 florets ♀, upper florets barren, embracing each other and forming a club-shaped or oblong body.....**Melica.**
Upper valves not forming a club-shaped body..... 121
121. Valves awned from the back..... 122
Valves awnless or awned from or near the tip or from the sinus..... 124

122. Ovary glabrous.....**Trisetum**.
 Top of ovary hairy..... 123
123. Spikelets large ; glumes rounded on the back, 7-11- nerved; panicle larger and lax with pendulous spikelets.....**Avena**.
 Spikelets medium sized; glumes more or less keeled, 1-3- (rarely up to 7-) nerved; narrow, stiff, erect panicle.....**LIV. Avenastrum**.
124. Valves awnless..... 125
 Valves awned from the tip or from the sinus..... 132
125. Valves closely imbricate, broadly boat-shaped, saccate, and scarious ; spikelets on long capillary pedicels.....**Briza**.
 Valves not broadly boat-shaped, or saccate and scarious..... 126
126. Spikelets sessile and solitary at the hollows of a simple spike..... 127
 Spikelets not sessile on a simple spike..... 130
127. Spikelets with their flattened side appressed to the rachis.....**Agropyrum**.
 Spikelets with their flattened side radial to the rachis.....**LXXX. Lolium**.
128. Styles lateral from the hairy 2-3-lobed appendage of the ovary...**LXXIX. Bromus**.
 Styles terminal or subterminal..... 129
129. Glumes and valves distinctly keeled, valves often with a tuft of long curled wool at the base.....**LXXVIII. Poa**.
 Glumes and valves rounded on the back or slightly keeled towards the tips, or glumes only keeled..... 130
130. Glumes and valves rounded on the back ; valves hyaline, variegated towards the tips, firm below, broad, obtuse.....**Atropis**.
 Glumes distinctly keeled, valves rounded or keeled towards the tips..... 131
131. Short annuals, with stiff panicles ; spikelets shortly pedicelled and more or less appressed to the short stiffly spreading branches ; valves oblong, obtuse ; ligules hyaline.....**Scleropoa**.
 Perennials with compound, usually open, flexuous, panicles (panicles reduced almost to a raceme in *F. vulpioides*) ; valves lanceolate, acute or subacute ; ligules scarious.....**Festuca**.
132. Valves with shining hyaline tips and margins ; awn very short ; panicle spike-like, cylindric.....**LXXVI. Koelaria**.
 Valves entirely herbaceous or chartaceous, or if with hyaline tips and margins, then not in a cylindric spiciform panicle..... 133
133. Ligules oblong-acute, up to 6 mm. long ; spikelets collected in compact, usually secund clusters, on a contracted or spiciform panicle.....**Dactylis**.
 Ligules rounded, truncate, or obtuse ; usually very short and often reduced to a membranous rim..... 134
134. Glumes of all, or at least of the lateral, spikelets very unequal, the lower glume minute or suppressed..... 135
 Glumes not very unequal..... 136
135. Spikelets on short club-shaped pedicels, the lower glume of all the spikelets minute or obsolete.....**Vulpia**.
 Spikelets sessile on a simple spike, lower glume of terminal floret equal or subequal to the upper.....**LXXX. Lolium**.
136. Ovary with an often large villous appendage, styles lateral from the appendage 138
 Ovary without a villous appendage..... 137

137. Spikelets of two kinds, the fertile surrounded by the sterile.....**Cynosurus.**
Spikelets all alike.....**Festuca.**
138. Spikelets subsessile in a simple raceme or false spike, with the flattened side to the axis.....**Brachypodium.**
Spikelets pedicelled, in open or contracted panicle.....**LXXIX. Bromus.**
139. Spikelets monoecious; ♂ spikelets smaller than the ♀ on the lower branches of the panicle.....**Olyra.**
Spikelets all alike ♀ 140
140. Stamens 3.....**Arundinaria.**
Stamens 6.....**Bambusa.**

I. IMPERATA, Cyr (109).

Spikelets all alike, generally 2-nate (one short-, the other long-pedicelled) on the continuous branches (racemes) of a spike-like or narrowly thyrsiform panicle, disarticulating from the pedicels. Florets 2; the lower reduced to an empty valve (very rarely ♂ ?). Glumes sub-equal, membranous, 3-9-nerved, rarely nerveless, enveloped by very long silky hairs from the obscure callus and the lower portion of both glumes. Valve of the lower floret generally much smaller than the glumes, hyaline; of upper floret still smaller, hyaline, awnless, rarely 0. Valvule broad, hyaline, nerveless. Lodicules 0. Stamens 1-2. Styles connate below; stigmas linear, exserted from the top of the spikelet. Grain oblong; embryo half the length of the grain or more.

Perennial; basal leaves crowded and, like those of the innovation shoots, long; panicle silvery-silky.

I. *arundinacea*, Cyr., var. *Thunbergii*, Hack.

Pretoria: *A. O. D. Mogg* in *National Herb.* 39, *Burtt-Davy* 828; Tzaneen, *Rogers* 12509; Warmbaths, Waterberg, *Stent*; Walven Kraal, Waterberg, *Pole Evans* 614.

A coarse grass with creeping underground stolons, common in vleis and on the banks of streams, widely distributed throughout South Africa, also in Tropical Africa; of no agricultural value.

II. MISCANTHIDIUM, Stapf (112).

Spikelets all alike, 2-nate, those of each pair unequally pedicelled on the articulate and tough or very tardily disarticulating rhachis of paniced racemes, falling from their pedicels. Florets 2; lower reduced to an empty valve, upper always ♀. Glumes equal, chartaceous to coriaceous, lower dorsally flattened, more or less distinctly 2-keeled, with inflexed margins and a variable number of intracarinal nerves, upper 1-3-nerved, keeled. Valves hyaline, of the lower floret muticous or mucronate, of the upper awned; awn

perfect. Valvules small, hyaline, nerveless. Lodicules 2, cuneate. Stamens 3. Stigmas usually laterally exserted. Grain oblong to linear-oblong; embryo half the length of the grain.

Perennial, erect; blades long, narrow, midrib very stout or the whole blade terete (tropical African species); ligules membranous; panicle much-branched, often large, silky, and showy, spikelets usually villous, and with an involucre of hairs at the base.

M. junceum, Stapf (Stapf in Hook. Ic., Pl. XXXI, tab. 3084 = *Erianthus junceus* Stapf in Dyer, Flor. Cap., VII, 324).

Potgietersrust, Waterberg, *Burt-Davy* 2306; near Nylstroom, Waterberg,

Burt-Davy 2000; Pyramid Estate, near Potgietersrust, Waterberg, *Galpin* 8873;

Almans Drift, Bushveld, Pretoria District, *Pole Evans* 597.

A tall reed-like grass growing in vleis and on river banks.

III. EULALIA, Kunth. (113A).

Spikelets all alike, or nearly so, 2-nate, one sessile, the other pedicelled, on the articulate fragile rachis of 2-nate, digitate or fascicled spike-like racemes, the pedicelled falling from their pedicels, the sessile deciduous together with the contiguous joint of the rachis and the pedicel. Florets typically 2; lower reduced to an empty valve or represented by a minute scale or quite suppressed; upper always ♀. Glumes equal, rigidly membranous to coriaceous; lower dorsally flattened or shallowly concave (not grooved), more or less 2-keeled with inflexed margins; upper 1-3-nerved, keeled. Valves hyaline; lower mucous, sometimes much reduced or quite suppressed; upper very short, 2-lobed, awned. Valvules small or 0. Lodicules 2, small, cuneate. Stamens 3. Stigmas linear, laterally exserted. Grain oblong; embryo almost half the length of the grain or longer; hilum basal, punctiform.

Perennial. Culms simple, erect, or ascending. Leaf-blades convolute in bud, then flat, usually narrow, gradually passing into the sheath. Racemes often coloured (brown or purplish).

E. villosa, Nees. (Stapf in Prain, Flor. Trop. Afr., IX, 99 = *Pollinia villosa*, Spreng., Stapf in Dyer, Flor. Cap., VII, 325).

Waterval Boven, *Burt-Davy* 1437; Sterk Hill, Lydenburg, *Burt-Davy* 450.

E. geniculata, Stapf. (in Prain, Flor. Trop. Afr., IX, 101).

Mosdene, Naboomspruit, in moist places. *Galpin* 442.

A tropical grass.

IV. ISCHAEMUM, Linn. (119).

Spikelets of each pair alike, or differing only in sex, or more or less heteromorphous, one sessile or sub-sessile, the other pedicelled on the articulate fragile rachis of 2-nate, digitate or fascicled, spike-like racemes, the pedicelled falling from their pedicels, the sessile deciduous together with the adjacent joint of the rachis and the pedicel. Florets 2; lower generally ♂; upper ♀, rarely ♂ or neuter in the pedicelled spikelets. Sessile spikelet: glumes equal or sub-equal, lower dorsally flattened or somewhat convex and usually coriaceous below, chartaceous and markedly nerved upwards, rarely shallowly concave and more or less chartaceous throughout (§ *Coelischæum*), more or less 2-keeled with inflexed margins; upper boat-shaped keeled at least above, sometimes awned. Valves rigidly membranous to hyaline, of lower floret mucous, of upper usually 2-fid and awned from the sinus, rarely mucronate or mucous. Valvules more or less equalling their valves, hyaline. Lodicules 2, cuneate. Stamens 3, sometimes smaller or rudimentary in the fertile flower. Stigmas linear-oblong, laterally exserted. Grain oblong or lanceolate, dorsally compressed; embryo reaching to the middle of the grain. Pedicelled spikelet in structure like the sessile, or more or less reduced; ♂ or neuter and awnless.




Plate I.

Ischaemum Mellei, Stent.

A, Pedicelled spikelet; B, Lower glume; C, Upper glume; D, Upper valve; E, Lower glume of sessile spikelet, back view; F, Lower glume of sessile spikelet, front view; G, Upper glume of sessile spikelet; H, Lower valve of sessile spikelet; I, Pale; J, Upper valve of sessile spikelet; K, Pale; L, Pistil and stamens

Generally perennial; blades convolute when young, at length flat; ligules generally membranous; racemes compressed, joints flattened or sub-concave on the inner side and often stout, sessile spikelets dorsally compressed, often rather broad, the pedicelled sometimes apparently laterally compressed with a median keel owing to the more or less complete suppression of one side.

I. arcuatum Stapf. (Prain, Flor. Trop. Afr., IX, 33 = *Ischaemum fasciculatum*, var. *arcuatum* Hook. ex Stapf. in Dyer, Flor. Cap., VII, 327).

Warmbaths, *Burt-Davy* 2339.

Also from Natal and Tropical Africa.

I. glaucostachyum, Stapf.

Strubens Farm, Pretoria, *Burt-Davy* 4482; Skinners Court, Pretoria, *Stent* H. 5173; Springbok Flats, Waterberg, *Burt-Davy* 1138; Rooikop, Bushveld, Pretoria District, *Pole Evans* 542, Nietverdiend; Zeerust, *Pole Evans* 329; Klipdrift, Marico District, *Pole Evans* 310.

A very glaucous grass with creeping underground stolons, good fodder for cattle, makes palatable hay, one of the best bushveld species. Commonly known as "turf-grass."

I. Mellei, Stent, n. sp.

Gramen perenne. *Culmi* erecti vel geniculati, ascendentes, simplices, vel ramis prope basim et aliquando e nodis superioribus, ad 120 cm. alti, glabri, tereti vel sub compressi. *Vaginae* foliorum laxae, striatae, tomentosae vel hirsutae, saepe longiores quam internodi, nodis albedo-barbatis; *ligulae* membranaceae, truncatae, ciliatae, ad 2 mm. longae; *laminae* lineares ad 20 cm. longae, 12 mm. latae, planae, dense hirsutae, cum circa 5 nervis lateralibus latere costa utroque. *Racemi* 3, raro 2-4, 6-9 cm. longi, medii breviter pedunculati; articuli triquetri, 3-4 mm. longi, recti vel paulo curvati, purpurescentes, ciliati; pedicelli articuli aequales et sub paralleli, et apud junctionem non excisi. *Spicula sessilis* 8 mm. longa; *glumae* inaequales, maculis purpureis, superior paulo longior; inferior late ovato-lanceolata, basi contracta, epice paulo bifido, 5-9 nervis, dorso rotundo, longo hirsuto, apicem versus bi-carinata, lato-alataque; superior cymbiforma, acuminata, mucronata, 5 nervi, margines superiores atque carina angusto-alata ciliati. *Anthoecium inferum*: valva glumam sub-equalis, lanceolata, acuta, uno-nervata, membranacea, marginibus superioribus ciliatis, palea valvam sub-aequalis, carinis asperulis. *Anthoecium superum*: valva 5 nerva, ad medium bilobata; lobae ciliatae, arista circa 7 mm. longa; *palea* angusto-oblonga, fere valvam aequalis, carinis scabris, antherae 2-5 mm. longae. *Spicula pedicellata* 6 mm. longa; gluma inferior cymbiforma, lanceolata, acuminata, 11 nervis, dorso longe-hirsuta, carina apicem versus late alata; cetera similis spiculae sessilis sed valva superior haud profunde lobata.

Pastills Farm, Pretoria. *Melle* in *Nat. Herb.* H. 21423.

Very near to *I. purpurascens*, Stapf. (ex. descr.), from which it differs in being larger in all its parts and very hairy, the joints and pedicels not being excised, and in the broadly winged keels of the lower glumes.

A very early vlei grass, very palatable to stock, but not frost resistant.

V. SEHIMA, Forsk. (119A).

Spikelets 2-nate, those of each pair differing in sex and (in the African species) also much in shape, one sessile, the other pedicelled on the articulate fragile rhachis of solitary spike-like racemes, the pedicelled tardily separating from their pedicels, the sessile deciduous together with the adjacent joint of the rhachis and the pedicel. Florets 2; lower ♂, upper ♀ in the sessile, ♂ or neuter in the pedicelled spikelets. Sessile spikelets: Glumes equal or sub-equal; lower deeply grooved, rarely flat, 2-dentate or 2-mucronate, more or less chartaceous, upwards acutely 2-keeled with inflexed margins, keels winged; upper glume boat-shaped, keeled upwards with a bristle-like awn. Valves hyaline, of lower floret entire, muticous, of upper 2-fid and awned from the sinus. Valvules more or less

equalling their valves, hyaline. Lodicules 2, cuneate. Stamens 3. Stigmas linear-oblong, laterally exserted. Grain oblong, obtusely trigonous; embryo reaching to the middle of the grain. Pedicelled spikelets flat, with (in the African species) a strongly nerved or ribbed lower glume, and 2 florets resembling the lower floret of the sessile spikelets, the lower or both more or less reduced and barren.

Perennial or annual; blades convolute when young, at length flat, narrow; ligules a line of stiff hairs; racemes usually gently curved, dorsiventral and laterally compressed, with the pedicelled spikelets converging over the convex side, joints and pedicels sublinear and parallel; sessile and pedicel spikelets strikingly heteromorphous in the African species.

S. galpinii, Stent, n. sp.

Gramen perenne. *Culmi* erecti, ad 1.5 m. alti, teretes, laeves, glabri, simplices, 2-4 nodi, internodi superiores, longissimi, exserti. *Folia* glauca; vaginae tenuite, sulcatae, laeves, glabrae vel nonnunquam nodi inferiores tomentosae; superiores aliquantum arctae, inferiores confertae fere flabellatae; *ligulae* ovatae, setis rigidis 2 mm. longis constitutae; *laminae* anguste lineares, circa 4 mm. latae, 15-30 cm. longae, longè in apicem setaceum attenuatae, planae, glabrae, scaberrulae, marginibus albis scabris cartilaginibus; costa gracilis, supra albida, 4-5 nervis lateralibus costae utrimque. *Racemus* solitarius; ad 19 cm. longus, strictus vel aliquantum curvus; articuli pedicellique paralleli, 7 mm. longi, subcompressi vel tenuite concavi, glabri nitensque, dense albo ciliatos margines. *Spicula* sessilis 15 mm. longa, subfalcata; *glumae* aequales; *inferior* praete apicem membranaceum chartacea, subfalcata, cum 9 nervibus inaequalis et cum transversis saepe brachiatis in parte superiore venulis, admodum carinata, carina exterior ala angusta ciliata et margine latiore inflexo, interior et exterior scabrida, dorsus flavus viridis nervibus carinisque; *superior* navicularis, chartacea, nitens, laevis, marginibus longe ciliatis, 5 nervis cum venulis transversis, apice carinata, carina ala ciliata, in setam longam excurrente. *Anthoecium inferum* ♂: *valva* 11 mm. longa, minute biloba, hyalina, 2 nervis, apicem versus scabridis, marginibus inflexis; *palea* similis valvae sed rigidior et omnino scabrida; *Antherae* 3, 6 mm. longae; *lodicae* cuneatae, carnosae, 1 mm. longae. *Anthoecium superum* ♀: *valva* 3 mm. longa, biloba, 3 nervis, parte inferiore hyalina, margines ciliatis; *arista* 40 mm. longa, columna bruno-fulva, marginibus involutibus minutissime ciliatis; *palea* valvam equalis, acuminata, minute biloba; *antherae* et lodicae iis in anthoecio infero similis; *styli* compressi, distincti; *caryopsis* trigona, urceolata. *Spiculae* pedicellatae 15 mm. longae: gluma inferior subfalcata, flava, purpureo-tincta, 7 nervis, intercarinas cum venulis transversis, nervi carinaeque viridi; *gluma* superior inferiorem subaequalis, sub-cymbiforma, membranacea, scabrida, 3 nervis, marginibus hyalinis, ciliatis; *anthoecia* 2, ♂; *valva* inferior, gracilis, membranacea, ciliata, minute bidentata, 3 nervis, nerva media in punctum callosum deltoideam excurrente; *palea* linearis, 2 nervis; *anthoecium* superum anthoecii inferi in spicula sessile simile.

Geluk, Springbok Flats, Waterberg, *Galpin* M. 557.

This comes nearest to the Indian species, *S. notata*, but has much larger spikelets. The flat lower glume of the sessile spikelet without a depressed channel on the median line separates it from other known African species, none of which are recorded so far south.

Mr. Galpin says of this grass: "In addition to growing on the farms Geluk and Zaai-plaats, there are also large patches of it growing on the adjoining farms Zandfontein Oog and Welbekend. . . . It is an excellent thatching grass, the culms being stout and very free from leaves, but it is not eaten by stock."

VI. URELYTRUM, Hack. (125).

Spikelets 2-nate, those of each pair different in shape and sex, one sessile, the other pedicelled, on the very fragile rhachis of solitary, paired, or racemed subcylindric spike-like racemes, the sessile deciduous with the adjacent joint of the rhachis and pedicel; joints obliquely truncate with appendaged tips. Sessile spikelets; florets 2, lower ♂




Plate II.

Sekima Galpinii, Stent.

A, Lower glume of sessile spikelet; B, Upper glume of sessile spikelet; C, Lower valve of sessile spikelet; D, Pale; E, Upper valve of sessile spikelet; F, Pale; G, Pistil and stamens; H, Pedicelled spikelet; I, Lower glume, front view; J, Upper glume; K, Lower valve.

upper 8. Glumes equal or subequal; lower coriaceous, dorsally flattened, 2-keeled, mucicous; upper subchartaceous or membranous, boat-shaped, keeled. Valves hyaline, subequal, slightly shorter than the glumes, mucicous, of lower floret 2-, of upper 3- to sub-5-nerved. Valvules almost equalling their valves, hyaline, 2-nerved. Lodicules 2, cuneate. Stamens 3. Stigmas linear, laterally exserted. Grain narrow, oblong, terete at the middle; scutellum to the middle; hilum obovate, basal. Pedicelled spikelets with 2 ♂ florets and with the lower glume narrowed into a subulate often long awn, otherwise as in the sessile spikelets, or more or less reduced, sometimes to a pair of rudimentary glumes, the lower of which is represented by a long awn, flattened and widened at the base.

Perennial, rarely annual; culms erect, few-noded, simple; blades narrow, linear, flat or convolute; racemes long, rigid.

U. squarrosus, Hack.

Paardevalle No. 12, near Zeerust, *Burt-Davy* 7196; Observatory Ridge, Johannesburg, *Burt-Davy* 3098; Houghton Estate, Johannesburg, *Bryant* C. 16.
A coarse sour grass.

VII. HEMARTHRIA, R.Br. (127A).

Spikelets 2-nate on the tough or tardily disarticulating rachis of spike-like spathe-supported racemes which terminate the culms and their often fascicled branches, alike in sex and shape, or at least similar; joints and pedicels fused into roughly semicylindric internodes, hollowed out on the inner face for the reception of the sessile spikelet; disarticulation at a right angle to the rachis or slightly oblique, tips of joints truncate, not hollowed out or appendaged. Sessile spikelet: florets 2, lower reduced to a barren valve, upper 8, awnless. Glumes equal or subequal, lower flat on the back, 2-keeled, very narrowly inflexed along the margins, coriaceous or subcoriaceous, closing up the cavity formed by the adjacent joint and pedicel, upper membranous adhering to the inner face of the cavity. Valves hyaline, of lower floret 2-nerved, of upper usually nerveless. Valvule (of upper floret) hyaline, small, nerveless. Lodicules 2, cuneate. Stamens 3. Stigmas laterally exserted. Grain oblong, dorsally slightly compressed; embryo about two-thirds the length of the grain; hilum conspicuous, punctiform, subbasal. Pedicelled spikelet with more elongated acuminate glumes, particularly the terminal, the upper glume mucronate or aristate.

Decumbent or ascending perennial grasses with many-noded branched compound culms; blades linear, conduplicate in bud, then flat; ligules very short, membranous; racemes compressed, often curved, tips more or less subulate owing to the slender terminal spikelet; spikelets pseudo-opposite owing to the fusion of joints and pedicels, each pair made up of a sessile (secondary) spikelet and the pedicelled companion of the sessile spikelet of the next lower node.

H. fasciculata, Stapf (Prain, *Flor. Trop. Afr.*, IX, 55 = *Rottboellia compressa*, var. *fasciculata*, Hack. ex Stapf in Dyer, *Flor. Cap.*, VIII, 329).

Irene, near Pretoria, *Burt-Davy* 3843, *Pole Evans* 366; Christiana, Bloemhof, *Burt-Davy* 11706; Waterval Boven, *Burt-Davy* 1409; Leeuwpoot, Carolina, *Burt-Davy* 112; Almans Drift, Bushveld, Pretoria District, *Pole Evans* 596.

A common species of vleis and streams usually semiprostrate, with creeping base. Found throughout Africa, moderately frost-resistant. Well liked by cattle.

VIII. TRACHYPOGON, Nees. (131).

Spikelets 2-nate, those of each pair differing in sex and slightly in shape, both pedicelled on the tough or upwards very tardily disarticulating many-jointed rachis of solitary, 2-nate, or digitate racemes, terminating the culms; pedicels unequal (the lower very short), fused towards the base with each other and the adjacent internode of the rachis, this and the pedicels terete or subterete, the longer of the latter disarticulating very obliquely;

internodes marked off by very oblique and usually shortly bearded articulation lines; subsessile spikelets persistent; long-pedicelled spikelets falling separately. Florets 2, lower reduced to an empty valve, upper ♂ or neuter in the subsessile, ♀ in the upper long-pedicelled spikelet. Fertile (upper) spikelets more or less terete, awned; callus acute to very pungent, densely bearded. Glumes equal or subequal; the lower firmly chartaceous to coriaceous, subconvolute, finely 2-keeled upwards, the inflexed margins very narrow or reduced to a mere rim, nerves 5-9 in addition to the keel-nerves, mostly obscure except in transmitted light, more or less connected by transverse veins in the upper part; upper thinner, 3-nerved, grooved on either side of the rounded keel. Valve of lower floret hyaline, 2-nerved, ciliate or ciliolate upwards, of upper floret stipitiform, hyaline and 3-nerved at the base, cartilaginous upwards and passing into a flexuous or geniculate awn. Valvule very small, hyaline or 0. Lodicules 2, small, glabrous. Stamens 3. Stigmas laterally, exserted; styles terminal. Grain terete; embryo about half the length of the grain. Male or neuter (lower) spikelets dorsally flattened (particularly the lower of a raceme) or more or less terete, awnless; callus 0. Glume and valve of the lower floret similar to those of the fertile spikelets. Valve of the upper floret narrow, hyaline, entire or 2-toothed, with or without a minute mucro, rarely produced into a short bristle. Stamens 3. Pistil, if present, rudimentary.

Perennial (very rarely annual?), densely caespitose grasses, with more or less slender wiry culms conspicuously bearded below the nodes; the beards appressed, covering the constricted nodes; leaf-blades narrow, frequently convolute.

T. plumosus, Nees. (ex Stapf in Prain, *Flor. Trop. Afr.*, IX, 403=*T. polymorphus*, var. *capensis*, Hack. ex Stapf in Dyer, *Flor. Cap.*, VII, 331).

Henley-on-Klip, *Stent* in *Nat. Herb.*, H. 21144; Groenkloof, Pretoria, *Phillips*, H. 21683; Johannesburg, on steep hillsides, *Bryant* D. 22; near Shilovane, Zoutpansberg, *Junod* 4077; Ermelo, *Burt-Davy* 5471; Leeuwpoot, Carol na, *Burt-Davy* 7343; Sterk Hill, Lydenburg, *Burt-Davy* 476.

Fairly widely distributed through the tropical and subtropical regions of the world.

A coarse grass only palatable when young; tested as a material for paper-making at the Imperial Institute, it was found to yield good pulp, but the ultimate fibres were rather short; good thatching grass.

IX. **ELYONURUS**, Humb. and Bonpl. (132).

Spikelets similar, usually awnless, but differing in sex, 2-nate, one sessile, the other pedicelled, on the articulate fragile rhachis of solitary spike-like racemes, the sessile deciduous with the adjacent joint of the rhachis and the pedicel. Florets 2; lower reduced to an empty valve; upper ♀ in the sessile, ♂, rarely barren, in the pedicelled spikelet. Glumes equal; lower subcoriaceous to herbaceous, often 2-toothed or 2-fid, rarely awned, dorsally flattened, 2-keeled, usually with fine filiform transparent balsam ducts close to the ciliate or penicillate keels; upper membranous, lanceolate, acute, rarely awned. Valves hyaline, awnless. Valvule obsolete or 0. Lodicules 2, cuneate. Stamens 3. Stigmas laterally exserted. Grain oblong, dorsally compressed; embryo about half the length of the grain.

Generally perennial, caespitose, aromatic; blades flat or folded; ligules membranous, very short; racemes erect, joints strongly compressed, usually villous, tips oblique, not appendaged.

E. argenteus, Nees.

Eloffs Plantation, Pretoria, *MacDonald* in *Nat. Herb.* 118; Derby Station, Marico, *Burt-Davy* 7168; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 410; Meintjes Kop, Pretoria, *Landsell* in *Nat. Herb.* 65; Henley-on-Klip, *Stent*, *Nat. Herb.* H. 21138; Potchefstroom, *Burt-Davy* 14569; Ermelo, *Burt-Davy* 952; Bloemhof, *Burt-Davy* 12953; Malans Farm, Hartbeestnek, *Burt-Davy* 758.

Widely spread throughout South Africa as well as in the tropics. A xerophytic grass of little value, usually characteristic of sour veld, known locally as "Wildebeeste gras," and "Zuur pol." Occasionally useful for winter grazing.

X. SORGHUM, Pers. (134A).

Spikelets 2-nate, those of each pair differing in shape and sex, one sessile, the other pedicelled or represented by a pedicel only, on the articulate fragile or (in cultivated forms) tough rhachis of panicle few- (sometimes 1- or, the other extreme, 6-8-) jointed racemes, the sessile spikelet falling with the contiguous joint and the accompanying pedicelled spikelet or at least its pedicel. Florets 2, lower reduced to an empty valve, upper ♂ in the sessile, ♂ or neuter in the pedicelled spikelets, if present at all. Sessile spikelets dorsally compressed. Glumes equal, coriaceous, at least when mature, rarely permanently chartaceous, mucous, lower with a broad flattened or convex back with the margins narrowly inflexed near the tips and otherwise involute, upper boat-shaped, keeled upwards with narrow hyaline, usually upwards ciliate margins. Valves hyaline, ciliate, of lower floret 2-nerved or nerveless, of upper oblong to ovate, 2-dentate or 2-lobed, with the lobes free or more or less adnate to a perfect or variously reduced awn or a mucro rising from the sinus, rarely entire and mucronate or mucous, 1- or 3-nerved. Lodicules 2, ciliate or glabrous. Valvule hyaline, often minute or 0. Stamens 3. Stigmas laterally exserted; styles terminal or sub-terminal. Grain in the spontaneous species mostly obovoid, dorsally compressed, in cultivated forms frequently enlarged, globose or sub-globose; embryo as long as or slightly longer than half the grain. Pedicelled spikelets, if present, much narrower than the sessile, lanceolate to subulate, ♂ or neuter, sometimes reduced to the glumes or one glume only, or quite suppressed. Glumes permanently herbaceous, like the hyaline 2-1-nerved ciliate; valves awnless.

Annual or perennial, often robust, grasses. Leaf-blades convolute in bud, usually flat, herbaceous, often large. Panicles erect or nodding, with verticillate or scattered branches, often large, in the spontaneous species mostly loose, in the cultivated forms frequently variously contracted to compact.

S. halepense, Nees. (Flor. Afr. Austr., 83=*Andropogon halepensis*, var. *effusus*, Stapf in Dyer, Flor. Cap., VII, 346).

Springbok Flats, Waterberg, *Burtt-Davy* 1128; Mountain View, Pretoria, *Wagstaff* in *Nat. Herb.* H. 162; Skinners Court, Pretoria, *Native Aaron*, *Nat. Herb.* H. 109; Johannesburg, *Nat. Herb.* H. 108; Groot Marico, *Burtt-Davy* 121; Barberton, *Burtt-Davy* 248.

This is the stoloniferous *Sorghum* known in America as Johnson grass. Stapf has followed Piper in doubting the existence of this grass as indigenous to tropical Africa and has omitted it from his Flora of that region. It certainly occurs spontaneously in subtropical parts of the Transvaal, mostly along streams. It is possible that in some localities from which it has been recorded as a troublesome weed, the seed has been introduced as an impurity probably with Sudan grass seed, but in other localities there appears little doubt that it occurs naturally indigenous to the soil. As in the case of most of the other species of *Sorghum*, it contains a certain amount of prussic acid.

S. versicolor, J. N. Anders (ex Stapf in Prain, Flor. Trop. Afr., IX, 138).

Mosdene, Naboomspruit, *Galpin* M. 419; Wonderboompoort, Pretoria, *Mogg* in *Nat. Herb.* 182; Potgietersrust, Waterberg, *H. M. Dewar* in *Nat. Herb.* H. 20947; Onderstepoort, Pretoria, *Native Aaron* in *Nat. Herb.* H. 183; Pyramids, near Pretoria, *Burtt-Davy* 4480.

Plentiful in the Rustenburg and Waterberg districts, sometimes behaving as an annual or at least flowering the first year. "Only found on black turf; much liked by stock and cut for fodder by some farmers."—*Galpin*.

S. ethiopicum, Rupr. (ex Stapf in Prain, Flor. Trop. Afr., IX, 119).

Zeerust, in *Nat. Herb.* H. 21416.

S. micratherum, Stapf (l.c. 142).

Mosdene, Naboomspruit, Waterberg District, *Galpin* M. 413; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4402.

XI. *CHRYSOPOGON*, Trin. (134b).

Spikelets in threes at the ends of the branchlets of terminal panicles, one sessile, the other 2 pedicelled, the trio falling entire from the thickened, nearly always bearded, oblique tips of the peduncles; exceptionally 2-nate in 2-jointed racemes, one sessile, the other pedicelled, each sessile spikelet falling with the contiguous joint and its pedicelled companion, pedicels (and joints, where any) linear-filiform, never longitudinally grooved not appendaged. Florets 2; lower reduced to an empty valve, upper ♂ in the sessile, ♂ or neuter in the pedicelled spikelet. Sessile spikelets usually laterally compressed, awned. Glumes subequal; lower coriaceous or chartaceous, involute with a rounded back or complicate and more or less keeled upwards, upper boat-shaped, more or less keeled. Valves hyaline, of lower floret 2-nerved, of upper linear, entire or 2-dentate with a usually perfect awn from the sinus. Valvule 0 or small, hyaline, nerveless. Lodicules 2, small, glabrous. Stamens 3. Stigmas exerted laterally low down. Grain linear, laterally compressed; embryo half the length of the grain; scutellum linear-oblong. Pedicelled spikelet dorsally compressed, awnless or aristulate.

Mostly perennial (all the Old World species). Leaf-blades narrow. Panicles usually lax, of whorls of simple or basally divided filiform branches, rarely the branches 2-nate or solitary.

C. montanus, Trin., var. *tremulus*, Stapf (= *Andropogon monticola*, var. *trinii*, Stapf Dyer, Flor. Cap., VII, 349).

Warmbaths, Waterberg District, *Burtt-Davy* 2338.

XII. *AMPHILOPHIS*, Nash (134c).

Spikelets 2-nate, one sessile, the other pedicelled, similar in shape or the pedicelled reduced and smaller, the latter always different in sex, except sometimes the lowermost pair, which may be homogamous (♂ or neuter), on the fragile rhachis of many-jointed shortly peduncled subdigitate or paniced racemes; joints and pedicels filiform, longitudinally grooved and hyaline in the groove, disarticulating horizontally sessile and pedicelled (always ?) spikelets deciduous, the former with the adjacent joint and pedicel. Florets 2 in the sessile spikelets; lower reduced to an empty valve, upper ♀, 2 or 1 in the pedicelled spikelet, the lower ♂ or neuter, the upper neuter or usually quite suppressed. Sessile spikelet dorsally compressed, awned; callus small, shortly bearded. Glumes equal, thinly chartaceous to membranous; lower 2-keeled, with narrow sharply inflexed margins; upper boat-shaped, 3-nerved, acutely keeled. Valve of lower floret hyaline, nerveless, of upper floret reduced to a hyaline linear stipe, firmer upwards, passing into a slender awn. Valvules 0 or very minute. Lodicules 2, minute, glabrous. Stamens 3. Stigmas exerted laterally, usually low down, longer than the styles. Grain oblong, obtuse, dorsally slightly compressed; embryo about half the length of the grain. Pedicelled spikelet awnless, valves (if present) hyaline, nerveless.

Perennial grasses with slender simple or branched culms, bearded or beardless at the nodes. Panicles mostly subdigitate with a short primary axis, rarely the racemes on branches of the second order; racemes always shortly peduncled. Spikelets small.

A. glabra, Stapf (Prain, Flor. Trop. Afr., IX, 172 = *Andropogon intermedius*, var. *punctatus*, Stapf in Dyer, Flor. Cap., VII, 345).

Hartebeestpoort Dam, Pretoria, *Pole Evans* 701; Mosdene, Naboomspruit, *Galpin* M. 437; Godwan River, *Pole Evans* 696; Rooikop, Bushveld, *Pole Evans* 263; Pienaars River, *Pole Evans* 166; Wonderfontein, Marico, *Burtt-Davy* 7220; Bushveld, Pretoria, *Reck* in *Nat. Herb.* 99; Waterberg, *Burtt-Davy* 2294; Shilovane, Zoutpansberg, *Junod* in *Nat. Herb.* 101.

Stapf describes the pedicelled spikelet of this species as being "not pitted," but in nearly all these specimens the pit on the lower glume of the pedicelled spikelet is distinct. *Burtt-Davy* 7220 exhibits a panicle with all of the spikelets, except a few towards the top

of the inflorescence, unpitted; Reck's specimen from Pretoria has some of the pedicelled spikelets unpitted, while others have one or occasionally two pits. Galpin's specimen presents a very robust form with leaves hairy beneath and up to 15 mm. broad and panicle 17 cm. long.

A. insculpta, Stapf (in Prain, *Flor. Trop. Afr.*, IX, 175=*Andropogon pertusus*, var. *capensis*, Stapf in Dyer, *Flor. Cap.*, VII, 345).

Barberton, *Burt-Davy* 237; Onderstepoort, Pretoria, *Native Aaron* in *Nat. Herb.* 93; Sterkstroom, *Burt-Davy* 132; Waterval Boven, *Burt-Davy* 1446; Pyramid Estate, near Potgietersrust, *Galpin* 8880; Mosdene, Naboomspruit, *Galpin* M. 411; Rooikop, Bushveld, Pretoria, *Pole Evans* 545.

XIII. **DICHANTHIUM**, Willemet (134b).

Spikelets 2-nate, one sessile, the other pedicelled, similar in shape, different in sex, except the lowermost 1 or 2 pairs of each raceme which are (with occasional exceptions in *D. annulatum*) homogamous (σ or neuter), in many-jointed shortly peduncled subdigitate, rarely subpanicled or racemously arranged, racemes; joints and pedicels filiform, solid, disarticulating subhorizontally, except the lowest barren pairs; fertile sessile and pedicelled spikelets deciduous, the former with the adjacent joint and pedicel. Florets 2 in the fertile sessile spikelets (lower reduced to an empty valve, upper σ), 1 in the barren sessile and all the pedicelled spikelets, σ or neuter, or suppressed. Sessile spikelet dorsally compressed, awned (except the basal homogamous ones); callus small shortly bearded. Glumes equal, thinly chartaceous; lower usually very obtuse, 2-keeled, with narrow sharply inflexed margins; upper boat-shaped, 3-nerved, acutely keeled. Valve of lower floret hyaline, nerveless, of upper floret reduced to a hyaline upwards firmer linear stipe, passing into a slender awn (very rarely finely 2-toothed with the awn from the sinus in a doubtful or aberrant Indian species). Valvule 0 or minute. Lodicules 2, minute, glabrous. Stamens 3. Stigmas exerted laterally at or above the middle or near the tips. Grain oblong, obtuse, dorsally compressed; embryo rather more than half the length of the grain. Pedicelled spikelet awnless; valve, if present, hyaline, nerveless.

Perennial, very rarely annual, grasses with simple or branched usually many-noded culms, bearded or beardless at the nodes. Panicles usually subdigitate with a short or very short primary axis, rarely the latter elongated; racemes always shortly peduncled. Spikelets small, rarely the male somewhat larger.

D. annulatum, Stapf (Prain, *Flor. Trop. Afr.*, IX, 178=*Andropogon annulatus* Forsk., Stapf in Dyer, *Flor. Cap.*, VII, 350).

Mosdene, Naboomspruit, *Galpin* M. 411A.

D. papillosum, Stapf (in Prain, *Flor. Trop. Afr.*, IX, 179).

Near Selous River, Rustenburg, *Burt-Davy* 153; Meintjes Kop, Pretoria, *Burt-Davy* 3552; Komatipoort, Barberton, *Burt-Davy* 369; Skinners Court, Pretoria, *Native Aaron* in *Nat. Herb.* 102.

XIV. **SCHIZACHYRIUM**, Nees. (134E).

Spikelets 2-nate, of each pair differing in sex and mostly also more or less in shape and size, one sessile, the other pedicelled, on the articulate fragile rachis of many-jointed solitary racemes terminating the culms and their branches, supported by spathes and frequently collected into a false panicle, the sessile spikelets falling with the contiguous joint, and the accompanying pedicelled spikelet; joints and pedicels thickened upwards, often rather stout, with a scarious cupuliform and more or less dentate terminal appendage. Florets 2; lower reduced to an empty valve, upper σ in the sessile, σ , neuter or 0 in the pedicelled spikelets or both suppressed in these. Sessile spikelets dorsally compressed or sometimes in their lower half almost terete, awned; callus short, with a short beard at the very base. Glumes equal or subequal, lower chartaceous (often thickly) to subcoriaceous, more or less convex (to sub-semiterete) or flat on the back with, at least from

the middle upwards, sharply inflexed and mostly narrow margins, 2-keeled, the keels running out into teeth or mucros; upper thinner to membranous, narrowly boat-shaped to dorsally flattened, keeled (at least upwards), 1-3-nerved, the delicate margins ciliate. Valves ciliate, hyaline, of lower floret membranous downwards and often purplish 2-nerved, of upper usually 2-fid, sometimes very deeply, or 2-dentate, rarely entire, awned, awn from the sinus or continuing the entire valve. Valvule 0 or a microscopic hyaline scale. Lodicules 2, minute, glabrous. Stamens 3, rarely 2. Stigmas laterally exserted low down; styles terminal. Grain narrowly linear in outline or tapering upwards, subterete; embryo short. Pedicelled spikelet similar to the sessile, but usually relatively broader and flatter or more or less reduced and then sometimes quite small. Glumes more or less membranous, the lower aristulate or muticous. Valves (if present) hyaline, ciliate, muticous.

Annual or perennial grasses, rarely suffrutescent, never very tall. Culms slender. Leaf-blades folded in bud, usually narrow. False panicles varying from very loose and scanty to densely fascicled with the spathes mostly very narrow scarious, membranous or lower down herbaceous.

S. semiberbe, Nees. (ex Stapf in Prain, Flor. Trop. Afr., IX, 195 = *Andropogon hirtiflorus*, var. *semiberbis*, Stapf in Dyer, Flor. Cap., VII, 337).

Mosdene, Naboomspruit, *Galpin* M. 414; Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21114; Pretoria, *Stent* in *Nat. Herb.* H. 110; Waterval Boven, *Burtt-Davy* 1447; Crocodile River, *Burtt-Davy* 210; Johannesburg, on hill-slopes, *Bryant* C 38; Springbok Flats, Waterberg, *Burtt-Davy* 1121; Warmbaths, *Baylis* in *Nat. Herb.* H. 125.

"There is no doubt that this grass would be quite suitable for the manufacture of paper-pulp on a commercial scale."—*Imp. Inst. Rep.* 8/5/18.

XV. ANDROPOGON, Linn. (134).

Spikelets 2-nate, those of each pair differing in sex and mostly also more or less in shape and size, or those of the lowest pair of the lowest raceme homogamous (♂ or imperfect), one sessile, the other pedicelled on the articulate fragile rachis of many-jointed paired (very rarely solitary) or corymbose (digitate or subdigitate) racemes terminating the culms, and, if these be branched, also their branches, or at least, the upper; raceme-pairs or corymbs supported by a spathe or spatheole and frequently collected into a false panicle; the sessile spikelets falling with the contiguous joint and the accompanying pedicel; joints and pedicels filiform or thickened upwards (rarely stout in members of *Piestium*) and then the tips frequently more or less cupular or auricled. Florets 2; lower reduced to an empty valve, upper ♀ in the sessile spikelets of the heterogamous pairs, ♂, neuter or suppressed in those of the homogamous pairs and in the pedicelled spikelets. Sessile spikelets dorsally or laterally compressed, nearly always awned; callus short, mostly quite obtuse, shortly bearded. Glumes equal or subequal, subcoriaceous to membranous, the lower flat or concave or channelled on the back, with, at least from the middle upwards, sharply inflexed margins, 2-keeled, the upper more or less boat-shaped, keeled upwards 3-1-nerved, sometimes aristulate. Valves ciliate or ciliate, rarely glabrous, of lower floret hyaline, 2-nerved, of upper 2-fid or 2-dentate, hyaline or firmer and sometimes substipitiform below the insertion of the awn. Valvule a hyaline nerveless scale, or 0. Lodicules 2, minute, glabrous. Stamens 3. Stigmas laterally exserted; styles terminal. Grain narrowly lanceolate to oblong in outline, subterete to plano-convex; embryo about half the length of the grain. Pedicelled spikelets often very different from the sessile in shape and less so in size, always more or less compressed dorsally, never concave or channelled on the back, sometimes reduced and then often small or quite suppressed. Glumes herbaceous-chartaceous to membranous, the lower muticous or very rarely aristulate. Valves, if present, hyaline, ciliate, muticous.

Mostly perennial grasses of varying habit.

A. eucomis, Nees.

Malmanie Oog, Lichtenburg, *Burt-Davy* 80; Vereeniging, *Brandmuller* 44; Panplaats, Middelburg, *Burt-Davy* 13260; Skinners Court, Pretoria, *Burt-Davy* 1023.

A. appendiculatus, Nees.

Schweizer Reneke, *Burt-Davy* 1697; Experiment Farm, Ermelo, *Burt-Davy* 3914; Skinners Court, Pretoria, *Stent* in *Nat. Herb.* 148; Volksrust, Wakkersstroom, in *Nat. Herb.* 150; Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21129; Heilbron, Vereeniging, *Brandmuller* 3; Tzaneen, Zoutpansberg, *Samson* in *Nat. Herb.* 141; Beginsel, Standerton, *Burt-Davy* 926; Johannesburg, *Burt-Davy* 832.

"A useful hay-grass valued for feeding."—*J. Burt-Davy*. Strongly stoloniferous.

A. schirensis, Hochst., var. *augustifolia*, Stapf.

Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21140; Aapies River, Pretoria, *H. Bolus* 10851; Downs, Zoutpansberg, *Junod* 4092; Billey's Vlei, Ermelo, *Burt-Davy* 9264; Middelburg, *Burt-Davy* 13295.

A. Schinzii, Hack.

Kosters River, Rustenburg, *Burt-Davy* 151; Pillansberg, Rustenburg, *Pole Evans* 633; Pyramid Estate, near Potgietersrust, *Galpin* 8885.

A. amplexens, Nees.

Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21139; Vereeniging, *Brandmuller* 42; Groenkloof, Pretoria, *Phillips* H. 21684; open veld, Downs, Zoutpansberg, *Junod* 4096; Johannesburg, *Bryant* D. 8; Experiment Farm, Ermelo, *Burt-Davy* 7720; Uitgevalen, Heidelberg, *Burt-Davy* 13667; Potchefstroom, *Burt-Davy* 2700.

"Not liked by stock, grows in tufts on well-drained sandy soil devoid of humus."—*Brandmuller*.

A. filifolius, Steud.

Mariesstad, *Pott* 5196; Belfast, Middelburg, *Burt-Davy* 1336; near Bossis, Carolina, *Burt-Davy* 2959; Lake Chrissie, *Hamilton* in *Nat. Herb.* 89; Spionkop, Ermelo, *Burt-Davy* 9239.

A. huillensis, Rendl.

Panplaats, Pan Station, Middelburg, *Burt-Davy* 13264; Johannesburg, *Hutton* 254; Pyramid Estate, near Potgietersrust, *Galpin* 8884.

XVI. **CYMBOPOGON**, Spreng. (134F).

Spikelets 2-nate, those of each pair differing in sex and more or less in shape—except those of the lowest pair of the lower or of both racemes, which are homogamous (♂ or neuter)—one sessile, the other pedicelled on the articulate fragile rhachis of many-jointed paired racemes, terminating the culms and their branches; raceme-pairs supported by a spatheole, collected into often decompound or supra-decompound spatheate panicles; the fertile spikelets falling with the contiguous joint and the accompanying pedicel; joints and pedicels filiform or linear with frequently more or less cupular or auricled tips, those of the lowest pair (raceme-base) often conspicuously swollen, oblong or barrel-shaped, and hard. Florets 2; lower reduced to an empty valve, upper ♂ in the sessile spikelets of the heterogamous pairs, ♂, rarely neuter, in those of the homogamous pairs and in the pedicelled spikelets. Fertile spikelets dorsally, rarely laterally, compressed, awned (normally); callus very short, obtuse, shortly bearded. Glumes equal or subequal, more or less chartaceous, the lower almost flat or slightly depressed or narrowly grooved on the back with at least from the middle upwards sharply inflexed margins, 2-keeled, the upper more or less boat-shaped, keeled upwards, usually 1-nerved. Valves ciliate or ciliolate (sometimes obscurely), of the lower floret entire, hyaline, 2-nerved, of the upper 2-fid or 2-lobed,

hyaline, rarely firmer and almost stipe-like below the insertion of the awn; column of awn, if any, smooth. Valvule 0. Lodicules 2, minute, glabrous. Stamens 3. Stigmas laterally exserted; styles terminal. Grain oblong in outline, subterete to plano-convex in cross-section; embryo about half the length of the grain. Pedicelled spikelets usually slightly different in shape and size from the sessile, but never depressed or grooved on the back. Glumes muticous, the lower chartaceous to subchartaceous, the upper thinner. Of the valves, only that of the lower floret present hyaline, 2-nerved; upper floret usually represented by a naked ♂ flower, rarely suppressed.

Perennial, densely tufted and usually aromatic grasses, with often very coarse leaves, and frequently much compound and contracted spatheate panicles.

C. marginatus, Stapf. (in Prain, *Flor. Trop. Afr.*, IX, 280 = *Andropogon nardus*, var. *marginatus*, Stapf in Dyer, *Flor. Cap.*, VII, 351).

Groenkloof, Pretoria, *Phillips*; Beginsel, Standerton, *Burt-Davy* 1807; Houtbosch, Pietersburg District, *Bolus* 10932; Crocodile River, near Pretoria, *Burt-Davy* 9281; Wonderboom, near Pretoria, *Burt-Davy* 8912; Pilgrims Rest, Lydenburg, *Burt-Davy* 439.

C. validus, Stapf. (= *Andropogon nardus*, var. *validus*, Stapf in Dyer, *Flor. Cap.*, VII, 352). Hamanskraal, near Pretoria, *Burt-Davy* 1085.

One of the tambookie grasses.

C. plurinodes, Stapf. (= *Andropogon pluri nodis*, Stapf in Dyer, *Flor. Cap.*, VII, 353).

Ermelo, *Burt-Davy* 3897, 5462, 966; Henley-on-Klip, *Stent* in *Nat. Herb.* 21110; Koster River, Rustenburg, *Burt-Davy* 170, 137; Vierfontein, Schweizer Reneke, *Burt-Davy* 1631; Uitgevalen, Heidelberg, *Burt-Davy* 9173; near Smitskraal, Bloemhof, *Burt-Davy* 10099; Doornkloof, Irene, near Pretoria, *Pole Evans* 384.

C. excavatus, Stapf (in Prain, *Flor. Trop. Afr.*, IX, 285 = *Andropogon Schoenanthus*, var. *versicolor*, Hack, ex Stapf in Dyer, *Flor. Cap.*, VII, 354).

Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21142; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 417; Heilbron, Vereeniging, *Brandmüller* 27; Johannesburg, *Bryant* D. 25; Skinners Court, Pretoria, *Native Aaron* in *Nat. Herb.* H. 213; Downs, Zoutpansberg, *Junod* 4084; Welverdiend Station, Potchefstroom, *Burt-Davy* 14583; Sterk Hill, Lydenburg, *Burt-Davy* 470; Waterval Boven, *Burt-Davy* 1438; Crocodile River, Rustenburg, *Burt-Davy* 9288; Irene, near Pretoria, *Pole Evans* in *Nat. Herb.* 216, 353, 204.

One of the turpentine grasses containing an essential oil that gives the grass a hot, bitter taste. It is seldom eaten by stock.

XVII. HYPARRHENIA, Anderss. (134g).

Spikelets 2-nate, those of each pair differing in sex and more or less in shape, or those of the lowest pair or pairs homogamous (♂ or neuter), one sessile, the other pedicelled, on the articulate fragile rhachis of few- to many-jointed paired racemes, terminating the culms and their branches; raceme-pairs supported by a spatheole, collected into scanty or compound to decompound spatheate panicles; the fertile spikelets falling with the contiguous joint and the accompanying pedicel; joints and pedicel finely filiform with more or less obliquely truncate sometimes obscurely auricled tips, in racemes with homogamous spikelets, the joint or joints below the lowest fertile spikelet entering into the formation of the raceme-base, shorter to very short, ciliate and tardily disarticulating. Florets 2; lower reduced to an empty valve, upper ♀ in the sessile spikelets of the heterogamous pairs, ♂ or neuter (rarely quite suppressed) in those of the homogamous pairs and in the pedicelled spikelets. Fertile spikelets more or less flattened dorsally, but with rounded sides, awned, very rarely awnless; callus short and obtuse or more often acute to pungent, shortly bearded. Glumes equal, more or less chartaceous, mostly thin, the lower narrowly truncate or 2-dentate, sometimes with a very shallow

longitudinal depression, with very narrow involute margins up to two-thirds or more, then equally narrowly inflexed from the very fine short or obscure keels, the upper narrowly and shallowly boat-shaped, rounded on the back below, finely keeled towards the tips only, 3-nerved. Valves ciliate or ciliate or that of the upper floret glabrous, of lower floret hyaline, 2-nerved (often very faintly), rarely quite nerveless, of upper stipitiform with a very fine delicate base, above it gradually hardened with narrow hyaline margins, which usually run out into minute teeth, passing between the teeth into the mostly distinctly kneed more or less hirtellous or subplumose awn. Valvule 0, rarely a small hyaline scale. Lodicules 2, minute, glabrous. Stamens 3. Stigmas laterally exerted; styles terminal. Grain oblong in outline, subterete to plano-convex in cross-section; embryo about half the length of the grain. Pedicelled spikelets acute, often mucronate or aristulate from the lower glume, usually somewhat longer than the fertile and more distinctly 2-keeled (often from below the middle). Valves of both florets developed, even in neuter spikelets, rarely the upper or both suppressed.

Perennial or annual, mostly coarse, grasses, with often large loose, rarely much contracted, panicles, never aromatic.

H. hirta, Stapf (in Prain, Flor. Trop. Afr., IX, 315 = *Andropogon hirtus*, Linn.; Stapf in Dyer, Flor. Cap., VII, 355).

Henley-on-Klip, *Stent* in *Nat. Herb. H.* 21127; Pretoria, *Stent* in *Nat. Herb. H.* 5161; *Burt-Davy* 3395, 3802; Heilbron, *Brandmuller* 26; near Wolhuters, Sterkstroom, *Burt-Davy* 133; Malmanie Oog, *Burt-Davy* 77, 70; Uitgevalen, Heidelberg District, *Burt-Davy* 12668; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb. H.* 4383; Henley-on-Klip, *Stent* in *Nat. Herb. H.* 21127; Kaalfontein, near Pretoria, *Pole Evans* in *Nat. Herb. H.* 17561; Irene, near Pretoria, *Burt-Davy* 33 *Pole Evans* 226, 354, 403.

"Useful as a thatch grass when no better grass is at hand."—*Brandmuller*.

H. auctus, Stapf (= *Andropogon auctus*, Stapf, in Dyer, Flor. Cap., VII, 357).

The Willows, Pretoria, *Burt-Davy* 1694; Johannesburg, *Bryant* C. 40; Vlakfontein, near Onderstepoort, Pretoria, *Burt-Davy* 4029; Sterkstroom, Rustenburg, *Burt-Davy* 138.

H. rufa, Stapf (in Prain, Flor. Trop. Afr., IX, 304 = *Andropogon rufus*, Kunth., ex Stapf in Dyer, Flor. Cap., VII, 358).

Tzaneen, Zoutpansberg, *Rogers* 12510; Barberton, *Burt-Davy* 278, 246; Zoutpansberg, *Wordsdeel* in *Nat. Herb. H.* 5918; Sterk Hill, Lydenburg, *Burt-Davy* 469.

H. dregeana, Stapf (= *Andropogon dregeanus*, Nees, ex Stapf in Dyer, Flor. Cap., VII, 359).

Stream at Houtenbek, near Lydenburg, *Burt-Davy* 517; Sterk Hill, Lydenburg, *Burt-Davy* 469.

"This grass yielded pulp of good quality; an excellent brown paper could be made from it by treatment with small amounts of caustic soda, while the pulp obtained with larger proportions of caustic soda could be readily bleached and used for the manufacture of white paper of good quality."—*Imp. Inst. Rep.*, 8/5/18.

H. dichroa, Stapf (in Prain, Flor. Trop. Afr., IX, 302 = *Andropogon dichroos*, Steud., ex Stapf in Dyer, Flor. Cap., VII, 360).

Tzaneen, Zoutpansberg, *Burt-Davy* 2631.

"Used as a thatching grass in the Orange Free State."—*Hilder*.

H. Buchanani, Stapf (= *Andropogon Buchanani*, Stapf in Dyer, Flor. Cap., VII, 362).

Premier Mine, Pretoria, *Menzies* 8; Skinners Court, Pretoria, *Native Aaron* in *Nat. Herb. H.* 7906; Sterk Hill, Lydenburg, *Burt-Davy* 469; Irene, near Pretoria, *Burt-Davy* 36.

"Its stems are slender, 4-5½ ft. long with pith in the internodes and somewhat refractory nodes 10-15 in. apart; the ultimate fibres are mostly 1.5-2.5 mm. in length,



Plate III.

Hyparrhenia glauca, Stent.

A, Lower glume of pedicelled spikelet; B, Upper glume of pedicelled spikelet; C, Sessile spikelet (showing lower glume); D, Upper glume of sessile spikelet; E, Lower valve of sessile spikelet; F, Upper valve of sessile spikelet; G, Pistil and stamens of sessile spikelet; H, Ligule.

and somewhat drastic treatment is required to bleach the pulp to a pale cream colour, but the paper produced is opaque, of good strength and quality, and does not shrink appreciably in the drying. It gives a good, long fibre pulp suitable for strong brown paper, or, after bleaching, for fairly good cream-coloured or white paper, and is regarded as of about the same value as Algerian esparte."—*Imp. Inst. Bull.* XIX, 2, 1921.

H. cymbaria, Stapf (Prain, Flor. Trop. Afr., IX, 332 = *Andropogon cymbarius*, Linn., ex Stapf, Flor. Cap., VII, 360).

New Agatha, Zoutpansberg, *McCallum* in *Nat. Herb.* 189; Irene, near Pretoria, *Hutchins* in *Nat. Herb.* H. 193; Malta, Zoutpansberg, *Junod* 4431; Zeerust, *Burt-Davy* 119; White Umbelusi, *Burt-Davy* 10609; Magoobaskloof, Houtboschberg, *Burt-Davy* 2604; Pyramid Estate, near Potgietersrust, *Galpin* 8887.

H. tamba, Anderss. (Stapf in Prain, Flor. Trop. Afr., IX, 336).

Waterval Onder, Carolina, *Burt-Davy* 393; Waterval Boven, Carolina, *Burt-Davy* 1419, 392.

H. variabilis, Stapf (Prain, Flor. Trop. Afr., IX, 334).

Godwan River, Carolina, *Pole Evans* 692.

H. filipendula, Stapf (in Prain, Flor. Trop. Afr., IX, 322 = *Andropogon filipendulus*, Hochst., ex Stapf in Dyer, Flor. Cap., VII, 362).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 415; Wolverkraal, Kameel River, *Pole Evans* 610; Godwan River, *Pole Evans* 693; Shilovane, Zoutpansberg, *Junod* 2213, 96.

H. Ruprechtii, Fourn. (Stapf in Prain, Flor. Trop. Afr., IX, 326 = *Andropogon Ruprechtii*, Hack., ex Stapf in Dyer, Flor. Cap., VII, 365).

Warmbaths, *Burt-Davy* 1153; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 418; between Nelspruit and White River, Barberton, *Burt-Davy* 1513; Komatipoort, Barberton, *Burt-Davy* 382; Barberton, *Burt-Davy* 288; Springbok Flats, Waterberg, *Burt-Davy* 1120.

"Zulu name 'unTshaba. It forms dense clumps reaching 10 ft. in height with hard nodes 11–14 in. apart. It gives a good yield of pulp, but the ultimate fibres are somewhat short, mostly ranging from .8 to 2 mm. in length. There is a considerable native demand for most of these tambookie grasses for thatch."—*Imp. Inst. Bull.*, Vol. XIX, 2, 1921.

H. glauca, Stent, n. sp.

Gramen perenne, ad 240 cm. altum, innovationibus intravaginalibus; culmi valedissimi, erecti, 4–8 mm. diam. tereti, glabri, saepe rubescentes, plus minus pruinosi, 7–9 nodis, supra nodis tumidi, simplices vel ramosi.

Vaginae foliorum firmæ, teretes vel apicem versus tenuiter, carinatae, glabrae. *Ligulae* scariosae, rotundatae, ad 4 mm. longae, flavae vel rufae, glabrae. *Laminae* lineares, apicem setiformam paullatim, attenuatae, ad 80 cm. longae, 12 mm. latae, glabrae, glaucae, subtus laevae, supra marginibusque scabrae; costa parvis valida, ad basim 2–4 mm. lata, supra albida, subtus prominente, nervis utrinque lateralibus 4–5. *Panicula* longa, laxa, gracilis, 5–8 nodis; internodis infimis ad 29 cm. longis, sequentibus brevioribus, folia subtudenta perfecta, sed apicem versus minora. *Rami* primarii fasciculati, plerumque 3–4 nati, inferiores 30–60 cm. longi ab basim 10–30 cm. indivisi, plerumque 3–4-nodi; ramuli filiformi, 1–4 pedunculis spatheates simplices ferenti. *Spatheolae* angustae, 4–6 cm. longae, setaceo-acuminatae, apicem versus rubescentes, cetera glauco-virides, deinde omnino rubescentes, glabrae, vel pilis longis dispersis, minute scaberulae. *Pedunculi* communes filiformes, ad 45 mm. plerumque circa 15 mm. longi, in spatheolis inclusi, apicem versus longe albedo-hirsutae. *Racemorum basi* breviores, dense pubescentes, et pilis albidis longis et tuberculis ortis barbata, spicula infimae basi in appendici scarioso denticulato 1 mm. longo producti. *Racemi* spatheola a latere exserti, plus-minus (aliquando late) divaricati, tandem reflecti et plus minus contigui, circa 15 mm. longi, aristis 8 quoque racemorum pare; articuli pedicellique gracilis, ciliati, spiculae duae racemis infimis basi mucronatae, 12 mm. longae. Spicula pedicellata ♂ 12 mm. longa; *gluma inferior* angusta,

virida, carinis scabridis, minute bidentata, mucrona ad 2 mm. longa, 11–13 nervis, prominentibus sparse appresseque pilis longis hirsuta; *gluma superior* gracile membranacea, 3–5 nerva, lineare lanceolata, acute acuminata, pilis longis appressis; *valva* hyalina, glumis paulo brevior, ciliata 1-nerva; *lodiculæ* carnosae, cuneatae; 5 mm. longae; *antherae* ad 4 mm. longae. *Spicula sessilis* ♂, pallido-virida, 5 mm. longa, callum brevior obtusum sparse barbatum annuerans; *gluma inferior* minute truncata, hirsuta, dorso plano, in parte superior carinis duobus, 7-nervis; *gluma superior* tenuiter chartacea, marginibus hyalinis 3-nervis, carina lata rotundata; *valva inferior* paulo glumis brevior, angusta, hyalina, enerva, apice abtuse; *valva superior* glumis multa brevior, hyalina, biloba, lobis angustis, obtusis; *arista* gracila, circa 10 mm. longa; *caryopsis* oblonga; *stylis* distincti.

Groenkloof Valley, Pretoria, *Stent* in *Nat. Herb.* H. 21425.

The upper valve of the hermaphrodite floret often appears to be stipitiform, the delicate hyaline margins and lobes being closely rolled round the base of the awn.

A remarkably glaucous species; the light blueish-grey colour of the foliage renders it most conspicuous from a distance.

XVIII. MONOCYMBIUM, Stapf (134H).

Spikelets 2-nate, those of each pair differing in sex and structure otherwise alike in shape, one sessile, the other pedicelled on the articulate fragile rhachis of many-jointed solitary racemes, terminating the culms and their branches; each raceme supported by a spatheole and forming with it a monostachyum; monostachya collected into loose spatheate panicles; the fertile spikelets falling with the adjacent joint and the accompanying pedicel; joints and pedicels finely filiform with truncate tips. Florets 2; lower reduced to an empty valve, upper ♀ in the sessile spikelets, ♂ in the pedicelled. Sessile spikelets dorsally flattened with rounded sides, awned; callus short and obtuse, indistinctly marked off from the lower glume, obscurely bearded. Glumes sub-equal, thinly cartilaginous, the lower obtuse, with narrow involute margins, not or very obscurely keeled near the tips, the upper boat-shaped, rounded on the back, laterally compressed and keeled in the upper third, usually aristulate, 3-nerved. Valve of the lower floret hyaline, ciliate, of the upper eciliate, deeply 2-fid, with a hyaline base, margins and lobes, otherwise cartilaginous, eciliate, delicately awned from between the lobes. Valvule 0. Lodicules 2, minute, cuneiform, glabrous. Stamens 3. Stigmas laterally exserted; styles terminal. Grain oblong, dorsally slightly compressed; embryo about half the length of the grain. Pedicelled spikelets in outline and size, very similar to the sessile, but mucous. Valves of both florets developed, hyaline, or the upper more frequently suppressed.

A perennial grass, with the coloured cymbiform spatheoles of a *Hyparrhenia* of the *Cymbaria* group, but easily recognized by its solitary racemes.

M. cerisiiforme, Stapf (in Prain, *Flor. Trop. Afr.*, IX, 386 = *Andropogon cerisiiformis*, Nees, in Dyer, *Flor. Cap.*, VII, 337).

Groenkloof, Pretoria, *Phillips* H. 21685; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 416; Johannesburg, *Bryant* D. 6; Sterk Hill, Lydenburg, *Burt-Davy* 457; Ermelo, *Burt-Davy* 5501; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* 131; Heidelberg, *Burt-Davy* 7769; Panplaats, Pan Station, Middelburg, *Burt-Davy* 13263; Cypress Point, Zoutpansberg, *Junod* 4085; Naboomfontein, *Schlechter* 4300; Belfast, Middelburg, *Burt-Davy* 1335; Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21113; Pyramid Estate, near Potgietersrust, *Galpin* 8888.

XIX. HETEROPOGON, Pers. (124I).

Spikelets 2-nate, those of the lower (1 to many) pairs alike in sex and shape, ♂ or neuter, of the upper pairs differing in sex and strikingly so in shape, one of each pair sessile, the other pedicelled on the many-jointed rhachis of solitary racemes, terminating the culms and their upper branches; rhachis tough or upwards tardily disarticulating and glabrous between the homogamous pairs, readily disarticulating above them; homogamous pairs

ong-persistent, the spikelets of the heterogamous pairs falling separately, the pedicelled with the pedicel, the sessile with the adjacent joint and the adjacent pedicel or its base. Florets 2; lower reduced to an empty valve, upper ♀ or ♂ in the sessile spikelets of the heterogamous pairs, ♂ or neuter in all the other spikelets. Fertile spikelet terete or subterete, awned; callus long, pungent, densely bearded upwards. Glumes equal; the lower coriaceous, rarely chartaceous, more or less tightly involute, quite keelless, nerves obscure, often connected by a few transverse nerves in the upper part; the upper with a deep longitudinal groove on each side, coriaceous, rarely chartaceous, between them, thinner towards the margins, membranous at the tips, 3-nerved. Valve of lower floret hyaline, nerveless, of upper floret stipitiform from a hyaline very slender base, cartilaginous upwards and passing into a usually stout geniculate awn. Valvule small or 0. Lodicules large or more or less reduced, to very minute. Stamens 3, often rudimentary, or 0. Stigmas exerted terminally or laterally. Grain more or less linear in outline, subterete, slightly dorsally compressed; embryo somewhat exceeding the middle of the grain. Male (or neuter) spikelets dorsally flattened, usually slightly asymmetric, and often somewhat twisted, muticous, imbricate. Lower glume herbaceous, many-nerved, winged upwards from one or both keels; upper membranous, lanceolate-oblong, acute, 3-nerved. Valves hyaline, 1-nerved, well developed or more or less reduced. Stamens 3 or 0.

Perennial or annual grasses, with simple or more often upwards branched culms; branches few to many, mostly flowering and gathered into a spatheate panicle; racemes conspicuously dorsiventral, the bases of the ♂ (or neuter) spikelets subimbricate on the back of the raceme, their upper parts bending forward around the sides, forming a hollow in which the fertile spikelets are enclosed, with their awns exerted anticously and often intertwisted.

H. contortus, Roem. & Schult. (Stapf in Prain, *Flor. Trop. Afr.*, IX, 410 = *Andropogon contortus*, L. in *Flor. Cap.*, VII, 350).

Kimberley, *Wilman*; Mosdene, Naboomspruit, Waterberg, in loam formation, *Galpin* M. 412; Koster River, Rustenburg, *Burtt-Davy* 151; Leeuwpoot No. 195, Carolina, *Burtt-Davy* 7332; Spitzkop, Ermelo, *Pott* 5210; Warmbaths, Waterberg, *Burtt-Davy* 2225; near Nylstroom, Waterberg, *Burtt-Davy* 2104; Schweizer Reneke, *Burtt-Davy* 1665; Welverdiend Station, Potchefstroom, *Burtt-Davy* 14578; Uitgevalen, Heidelberg, *Burtt-Davy* 9183; Experiment Farm, Ermelo, *Burtt-Davy* 915.

"Spear grass"—"Steek gras." A useful fodder grass when young, but the sharp-bearded calli of the mature spikelet render the grass, when in seed, very troublesome and often dangerous to stock, the "seeds" often pierce right into the flesh of small stock, causing great irritation and even in some cases death; in cattle, these "seeds" have been known not only to injure the beasts, but to materially damage the hide for commercial purposes.

XX. CLEISTACHNE, Benth. (134).

Spikelets solitary, all alike, ♀, pedicelled on the tough rachis of racemously arranged or panicked racemes, falling entire from the thickened tips of the pedicels. Florets 2; lower reduced to an empty valve, upper ♀. Glumes equal, very similar, with involute margins, more or less coriaceous, delicately 7-9-nerved, muticous. Valves hyaline, lower 2-nerved, upper 2-dentate or subentire, 3-nerved, with a twisted flexuous awn from the sinus or tip. Valvule very minute, ciliate. Lodicules 2, broad-cuneate, sparingly ciliate. Stamens 3. Stigmas laterally exerted, plumose. Grain oblong to obovoid-oblong, very obtuse to truncate; embryo half the length of the grain.

Annual, rather coarse grasses. Leaf-blades long, flat, with stout midribs. Panicles narrow, more or less contracted, greyish or fulvously hairy.

C. sorghoides, Benth. (Stapf in *Flor. Trop. Afr.*, IX, 154).

Godwan River, Carolina, *Pole Evans* 694.

A tropical grass and a new record for the Transvaal.

XXI. **THEMEDA**, Forsk. (136).

Spikelets 2-nate (the terminal 3-nate), those of the lower two pairs alike in shape and sex, ♂ or neuter, sessile to subsessile and closely approximate so as to form an involucre, of the following pair or pairs differing in sex and conspicuously so in shape, one of each pair sessile, the other pedicelled, on the rhachis of solitary racemes, terminating the culms and their upper branches; rhachis terete, tough or tardily disarticulating and glabrous between the homogamous pairs, readily disarticulating below the fertile spikelets; homogamous pairs or at least their sessile members persistent, the spikelets of the heterogamous pairs falling separately, the pedicelled with the pedicel, the sessile with the adjacent joint (if more than 1) and the base or bases of the adjacent pedicels. Florets 2; lower reduced to an empty valve, upper ♂ or neuter in the homogamous spikelets and the pedicelled members of the heterogamous pairs, ♀ in the sessile members of the latter. Fertile spikelets terete, awned or awnless; callus usually acute to pungent, densely bearded, beard more or less concealing the adjacent joint or pedicel-bases. Glumes equal, coriaceous, except at the submembranous tips; lower tightly involute, quite keelless, nerves very fine, obscure except in transmitted light, the upper with a deep longitudinal groove on each side, very firm between them, thin long, the margins 3-nerved. Valve of lower floret hyaline, nerveless, of upper floret stipitiform from a hyaline 1-nerved base, cartilaginous upwards and passing into a stout geniculate awn or hyaline and 1-nerved throughout, with the nerve becoming stouter upwards and passing into a more or less reduced awn or quite awnless. Valvule hyaline, nerveless, small or 0. Lodicules 2, somewhat large, glabrous. Stamens 3. Stigmas exerted subterminally or from near the middle of the floret. Grain linear-obovate, subterete, grooved on the face; embryo about half as long as the grain. Involucral spikelets dorsally much compressed, awnless. Lower glume more or less herbaceous, 2-keeled; upper membranous, very rarely suppressed. Both florets present or one or both suppressed, their valves hyaline, the upper with or without a valvule. Pedicelled spikelets of the heterogamous pairs similar to those of the homogamous pairs, but narrower.

Annual or perennial; usually coarse grasses, with scanty or compound to decompound spatheate panicles, and scattered or more or less crowded and often glomerate short 1- or 2- (very rarely 3-) awned or awnless monostachya.

T. triandra, Forsk. (Stapf in Flor. Trop. Afr., IX, 3, 416 = *Anthistiria imberbis*, Retz., ex Stapf in Dyer, Flor. Cap., VII, 366).

Var. **Burchelli**, Stapf.

Schweizer Reneke, Bloemhof, *Burt-Davy* 1642.

Var. **hispida**, Stapf.

Near Nylstroom, Waterberg, *Burt-Davy* 2025A; Arcadia, Pretoria, *Burt-Davy* 12324; Ermelo, *Miss Tennant* in *Nat. Herb.* 259; Carolina, *Burt-Davy* 7425; Crocodile River, Rustenburg, *Burt-Davy* 224; Uitgevalen, Heidelberg, *Burt-Davy* 9174; Komatipoort, Barberton, *Burt-Davy* 362; Potchefstroom Farm, *Burt-Davy* 1052; Skinners Court, Pretoria, *Burt-Davy* 3809; Springbok Flats, Waterberg, *Burt-Davy* 1119.

Var. **glauca**, Stapf.

Meintjes Kop, Pretoria, *Miss Landsdell* in *Nat. Herb.* 253; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 420.

Var. **punctata**, Stapf.

Near Nylstroom, Waterberg, *Burt-Davy* 2012; Mariestad, *Pott* 5212; Belfast, Lydenburg, *Burt-Davy* 1333; Sterk Hill, Lydenburg, *Burt-Davy* 455.

This is the well-known rooi gras—an excellent pasture and hay grass if cut before it gets too ripe. It is one of the grasses examined at the Imperial Institute for paper-making and was found to yield about the same quality and quantity of pulp as the tambookie and dek grasses previously reported on, except that the nodes being harder would require more boiling to render them soft enough for the purpose.

XXII. **ANTHEPHORA**, Schreb. (140).

Spikelets lanceolate to ovate-oblong, sub-sessile in deciduous squarrose clusters 4-6, on the very short branches of a cylindric apparently simple spike; lower floret reduced to an empty valve, upper ♀ or ♂, with or without the rudiments of an ovary. Glumes very dissimilar, lower always turned away from the centre of the cluster, coriaceous, oblong to lanceolate, distinctly nerved on the inner side, upper hyaline, setaceously subulate from a small ovate base. Valves more or less equal, usually shorter than the lower glume; lower oblong, hyaline, 5-7-nerved; upper similar, but firmer and 3-nerved. Pale equalling the valve and of similar substance, 2-nerved. Lodicules 0. Stamens 3; filaments fleshy at the base when young. Ovary lanceolate-oblong. Styles free or nearly so, very slender. Stigmas plumose, slender, exerted from or near the apex. Grain oblong-ellipsoid, enclosed in the spikelet; hilum punctiform; embryo large, equalling about half the length of the grain.

Perennial or annual; culms tufted and erect, or ascending from a prostrate base; ligules large scarious; spikelets in subspicate clusters with the irrespective lower glumes turned outwards so as to form a false involucre, usually with gaps between their bases.

A. pubescens, Nees.

Hamanskraal, Pretoria District, *Pole Evans* 392; Doornkloof, Irene, near Pretoria, *Pole Evans* 222.

Also known to occur plentifully near Naboomspruit in the Waterberg District and at Potchefstroom. A tufted grass with very glaucous leaves, useful pasture for stock, one of the earliest grasses in Bechuanaland.

XXIII. **TRAGUS**, Haller (143).

Spikelets sessile, in deciduous clusters of 2-4 on the filiform continuous axis of a cylindric spike-like panicle; rachilla tough, not continued beyond the floret. Floret 1, ♀, somewhat shorter than the upper glume. Glumes very dissimilar, lower facing the rachis, minute, hyaline, or suppressed, upper 5-ribbed or 5-nerved, membranous between the hispid or spinehooked ribs or nerves, exceeding the valve. Valve lanceolate or lanceolate-oblong, membranous, 3-nerved. Pale as long as the valve, 2-nerved. Lodicules 2, broad, cuneate, fleshy. Stamens 3. Styles distinct, very slender. Stigmas narrow, plumose, terminally exerted. Grain enclosed by the valve and pale, oblong to ellipsoid, slightly compressed from the back; embryo about one-third the length of the grain; hilum punctiform, basal.

Annual or perennial; culms erect, ascending or decumbent; blades linear, rather rigid with cartilaginous spinulously ciliate margins; ligules reduced to a delicate ciliate rim; panicles cylindric, slender; all the spikelets of a cluster, fertile, or often one more or less reduced.

T. racemosus, All.

Kafraria, Christiana, Bloemhof, *Burt-Davy* 14109; near Bloemhof, *Burt-Davy* 1504; Warmbaths, Waterberg, *Burt-Davy* 863; near Nylstroom, Waterberg, *Burt-Davy* 2084; Potgietersrust, Waterberg, *Burt-Davy* 2244; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 409; Barberton, *Burt-Davy* 241; Pretoria, *Miss Tennant* in *Nat. Herb.* H. 6313; Vereeniging, *Brandmüller* 30; Zeerust, Marico District, *Burt-Davy* 115.

"Carrot seed grass"—a common weed of most warm countries.

T. Koelerioides, Aschers.

Vierfontein, Schweizer Reneke, *Burt-Davy* 1630; near Bamboespruit, Bloemhof, *Burt-Davy* 1521.

Creeping carrot seed grass—a short creeping grass, useful only as a crop for sheep.

XXIV. **MOSDENIA**, Stent (143A).

Spikelets sessile, solitary on the notches of a ribbed and scabrid common axis disarticulating from below the glumes and falling entire. Floret 1. Glumes subequal to equal, 3-4 mm. long, lower rather longer and broader than the upper, cymbiform, falcate rigidly membranous with a prominent scabrid keel; upper like the lower, but narrower and straighter. Valve about 3 mm. long, delicately hyaline, 1-nerved or sub 3-nerved. Pale nearly equalling the valve. Stamens 3. Ovary ovate. Styles short, distinct. Lodicules truncate, about as large as the ovary.

A prostrate ascending stoloniferous perennial, the stolons being covered with short, dry, densely imbricating scales. Leaf-sheaths tight, terete, the lower short. Ligule short, membranous. Inflorescence of solitary spikes terminating the culms.

M. Waterbergensis, Stent (*Bothalia* I, 3, 170).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 447.

A monotypic genus.

XXV. **PEROTIS**, Ait. (148).

Spikelets very small, narrow, sessile or sub-sessile on the continuous axis of a spike or a lax spike-like raceme, jointed on and falling entire from the axis or the rudimentary pedicels; rhachilla not continued beyond the floret. Floret 1, ♂, much shorter than the glumes. Glumes equal, linear or linear-lanceolate, rigidly membranous, 1-nerved, passing into capillary awns. Valve lanceolate, acute, delicately hyaline, 1-nerved. Pale very minute, hyaline nerveless. Lodicules 2, broad, cuneate. Stamens 3. Styles distinct, short. Stigmas plumose, laterally exserted. Grain cylindric, slender, exserted from the unchanged floret and enclosed with it in the glumes; embryo about one-third the length of the grain; hilum punctiform, basal.

Annual or subperennial; culms leafy; blades usually broad, rigid, and ciliate; ligules hyaline; spikes or pseudo-spikes slender, crinite from the long capillary awns.

P. latifolia, Ait.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 441; Warmbaths, Waterberg, *Burt-Davy* 861; Barberton, *Burt-Davy* 8014; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4436; Riviera, Pretoria, *Mogg* in *Nat. Herb.* 10085; Waterval Boven, *Burt-Davy* 1410; Koster River, Pietersburg, *Burt-Davy* 157; Pyramid Estate, near Potgietersrust, *Galpin* 8883.

XXVI. **ARUNDINELLA**, Raddi. (153).

Spikelets small, $2\frac{1}{2}$ lin. to less than 1 lin. long, acute or acuminate, panicked, continuous with or imperfectly jointed on the pedicels; rhachilla disarticulating between the valves, glabrous, not produced beyond the upper floret. Florets 2, heteromorphous; lower ♂, rarely barren or ♀, upper ♀. Glumes persistent, acuminate, membranous, strongly 3-5-nerved, upper usually longer. Lower valve thin, equalling the lower glume or slightly longer, subacute or minutely truncate, 3-7-nerved; upper valve terete, rarely dorsally subcompressed, finally cartilaginous, entire or minutely 2-fid, 3-7-nerved, minutely scaberulous, lobes sometimes produced into fine bristles; awn terminal or from between the lobes, short, kneed or straight, or 0; callus very short, obtuse. Pales linear, 2-keeled; flaps more or less auricled. Lodicules 2, cuneate, rather fleshy or 0 in the lower floret. Stamens 3 or 0 in the lower floret. Ovary glabrous, oblong or more or less arrested, or 0 in the lower floret. Styles distinct. Stigmas plumose, laterally exserted. Grain oblong to ellipsoid, terete or dorsally subcompressed, tightly embraced by the valve and pale, free; hilum punctiform; embryo large; albumen hard.

Perennial, rarely annual; leaves various; panicles usually rather stiff; branches stiff, divided from the base.

A. Ecklonii, Nees.

Vlakfontein, near Amersfort, Wakkerstroom, *Burt-Davy* 4055; Skinners Court, Pretoria, *Stent in Nat. Herb.* H. 5076; Ermelo, *Burt-Davy* 5407; White Umbelusi, Zoutpansberg, *Burt-Davy* 10607; Waterval Boven, Lydenburg, *Burt-Davy* 1400; Vereeniging, *Brandmuller* 2.

"A coarse, hard grass of vleis and swamps; of apparently little feeding value"—*Burt-Davy*.

XXVII. MELINIS, Beauv. (154).

Spikelets linear-oblong, small, laterally compressed, panicle, deciduous from the capillary pedicels; lower floret reduced to an empty valve; upper ♂. Glumes very dissimilar, lower reduced to a minute scale, upper membranous, shortly 2-lobed, mucronulate from the sinus, prominently 7-nerved, glabrous, rarely hairy. Valves very dissimilar; lower rather like the upper glume, but more deeply lobed and usually with a fine awn from the sinus, 5-nerved; upper smaller, very thin and rigidly membranous, minutely 2-lobed, shining, obscurely 3-1-nerved with an equal, obscurely 2-nerved pale of similar substance. Lodicules 2, very delicate. Stamens 3. Styles free, slender. Stigmas plumose, laterally exerted. Grain unknown.

Perennial or annual (?); culms ascending from a prostrate, very slender, many-noded, branched base; ligules reduced to a fringe of short hairs; panicle contracted narrow, much divided, with capillary branchlets and pedicels; spikelets usually purple, at least, at the tip.

M. tenuinervis, Stapf (= *M. minutiflora*, var. *pilosa*, Stapf in Dyer, *Flor. Cap.*, VII, 447). Barberton, *Burt-Davy* 270; Tzaneen, *Mogg in Nat. Herb.* H. 10454.

Nearly related to the molasses grass of tropical Africa—*Melinis minutiflora*.

XXVIII. ALLOTEROPSIS, Presl. (161A).

Spikelets ovate or elliptic to lanceolate-oblong, acute or acuminate, mostly awned, slightly or conspicuously compressed from the back, falling entire from the pedicels, 2-nate or fascicled, subsecond and abaxial on the triquetrous rachis of more or less spiciform digitate or subdigitate racemes; lower floret usually ♂, upper floret ♀. Glumes unequal, lower smaller, membranous to hyaline, 3-1-nerved, very acute, often mucronulate, upper equal or subequal to the spikelet, membranous to chartaceous, 5-nerved with the outer nerves submarginal, densely ciliate along them. Lower floret: valve resembling the upper glume, but eciliate, the tip minutely truncate and the lateral nerves approximate to each other and in or near the marginal flexure; valvule short, deeply 2-fid, with conspicuously auricled flaps. Upper floret: valve chartaceous, glabrous, delicately ciliate upwards, 5-nerved, produced into a straight awn or mucronate; valvule equal to the valve, 2-keeled, with broadly auricled flaps. Lodicules 2, broadly cuneate. Stamens 3. Styles distinct. Stigmas laterally exerted. Grain enclosed by the valve and valvule, elliptic-oblong much compressed; scutellum about half the length of the grain; hilum basal, punctiform.

Perennial or annual; leaf-blades flat or more or less convolute; ligules membranous, ciliate or ciliate, short or reduced to a mere rim; racemes sessile or peduncled, often more or less compound towards the base, digitate or subdigitate on a more or less elongated common axis.

A. semialata, Hitch, var. *Ecklonii*, Stapf (*Flor. Trop. Afr.* IX, 3, 483 = *Axonopus semialatus*, Hook, var. *Ecklonii*, Stapf in Dyer, *Flor. Cap.*, VII, 418).

Vereeniging, *Brandmuller* 47; Ermelo, *Tennant in Nat. Herb.* 289; Ermelo, north of Blesbokspruit, *Burt-Davy* 946; Carolina, *Burt-Davy* 7367, 7446; Belfast, *Burt-Davy* 1323; Johannesburg, *Bryant* D. 3; Germiston, Witwatersrand, *Fenn in Nat. Herb.* 316; Beginsel, Standerton, *Burt-Davy* 1020, 3082; Meintjes Kop, Pretoria, *Fouche*; Derby Station, Marico, *Burt-Davy* 7170; Mariestad, *Pott* 5214; Downs, Zoutpansberg, *Junod* 4081.

"Occasionally grazed when growing among other grasses, but rather coarse."—*Burt-Davy*.

XXIX. **PASPALUM**, Linn. (161).

Spikelets orbicular to oblong, mostly decidedly plano-convex, falling entire from the short rudimentary pedicels, solitary or paired, secund and abaxial on the dilated or narrow rhachis of spike-like racemes; lower floret barren, reduced to its valve; upper floret ♂. Glumes: the lower typically suppressed, very rarely represented by a small scale; the upper more or less equal to the spikelet, rarely shorter or absent, convex, membranous, 3- or more-nerved. Lower floret: valve very similar to the upper glume, but usually flat or at least less convex, rarely subconcave, the middle nerve sometimes suppressed. Upper floret: valve chartaceous to crustaceous with firm margins, obtuse, emucronate, faintly nerved; valvule subequal to the valve and similar in substance, tightly embraced by the narrowly involute margins of the valve. Lodicules 2, minute, broadly cuneate. Stamens 3. Styles distinct. Stigmas laterally exerted near the tip of the floret. Grain tightly enclosed by the slightly hardened valve and valvule, more or less biconvex; scutellum orbicular-elliptic, shorter than half the grain; hilum subbasal, punctiform.

Mostly perennial grasses of varying habit.

P. distichum, Linn. (= *P. digitaria*, Poir., Stapf in Dyer, *Flor. Cap.*, VII, 370).

Wonderboom, near Pretoria, *Nat. Herb.* H. 19604.

P. scrobiculatum, Linn., var. **Commersonii**, Stapf (in Prain, *Flor. Trop. Afr.*, 573 = *P. scrobiculatum*, L., Stapf in Dyer, *Flor. Cap.*, VII, 370).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 389; Middelburg, *Nat. Herb.*

302; Downs, Zoutpansberg, *Junod* 4099; Tzaneen, Zoutpansberg, *Sampson* in

Nat. Herb. 304; Lydenburg, *Nat. Herb.* 306.

"Ditch grass," the Kodo Millet of India—a useful pasture grass in some localities. "Known to be poisonous and injurious to animals and man in India. The seed, especially the testa and pericarp, contain a narcotic poison. . . . Cattle should not be allowed to feed on it when it is ripening."—*Pammel*.

We have no records of poisoning from this grass in South Africa. Susceptible to ergot.

P. dilatatum, Poir.

Occurs as an alien, probably an escape from cultivation, in moist localities. It is largely cultivated as a pasture and hay crop, but is subject to attacks of ergot (*Claviceps Paspali*). Very good pasture grass for sour veld.

XXX. **ERIOCHLOA**, H. B. and K. (164).

Spikelets ovate-lanceolate to oblong, acute or acuminate, sometimes finely and shortly aristulate, slightly compressed from the back, falling entire from the pedicels, 2-nate or solitary, secund and abaxial on the triquetrous rhachis of more or less spiciform simple or compound racemously arranged racemes; lower floret ♂ or barren, upper floret ♂. Glumes very unequal; the lower reduced to a minute cupular rim clasping the callously thickened bead-like lowest internode of the spikelet, upper corresponding in outline and size to the spikelet, sometimes shortly aristulate, membranous, faintly 5-nerved, the submarginal nerves sometimes obscure. Lower floret: valve very like the upper glume, if not slightly shorter; valvule subequal to the valve, 2-keeled or reduced or, like the lodicules and flower, quite suppressed. Upper floret: valve chartaceous to almost crustaceous, glabrous or very minutely puberulous at the apex, faintly 5-nerved, with a fine barbellate mucro of varying length; valvule equal to the valve, 2-keeled, with moderately narrow flaps. Lodicules 2, broadly cuneate. Stamens 3. Styles distinct. Stigmas laterally exerted near the tips of the spikelets. Grain enclosed in its floret and the tightly appressed lower floret and upper glume, dorsally compressed; scutellum rather exceeding the middle of the grain; hilum subbasal, punctiform.

Perennial or annual; leaf-blades more or less flat, ligules reduced to a ciliate rim. Racemes peduncled or the upper sessile, racemously arranged in a narrow panicle.



Plate IV.

Eriochloa Fouchetii, Stent.

A, Rachis of raceme with pedicels ; B, Spikelet ; C, Upper glume ; D, Lower valve ;
E, Upper valve ; F, Pale ; G, Pistil and stamens.

E. Fouchei, Stent, n. sp.

Gramen annua. Culmi erecti, vel geniculati, 4-7 nodi, simplices vel nodis inferioribus ramosi, compressi, glabri, internodii infimi 4-5 cm., cetera breviores, breviter exserti, vel supremi basin paniculae inclusi. *Foliorum vaginæ* compressae, carinatae, tenuiter striatae. *Ligulae* ad coronam dense ciliatam 1.5 mm. longa redacta. *Laminae* lineares, in apicem tenuem attenuatae, ad 8 mm. latae, 22 cm. longae, glabrae vel basin pubescente, planae vel complicatae; costa gracila, distincta, supra tenuiter canaliculata, subtus carinata. *Panicula* angusta, ad 21 cm. longa, racemo pluris (ad 20), appressis, vel obliquo patulo, solitarius, vel 2-3 natis. *Rachis* communis tenuiter sulcatis, minute pubescentis. *Racemi* ad 5 cm. longi, internodios excedens, breviter pedunculati, infimi saepe basin compositi: rachis circa .7 mm. latis, minute pubescentis, supra carina flexua, apex in spiculam terminans. *Pedicelli* plerumque 2-nati, longiore ad 2 mm. longi, carinam utroque alternis dispositus, ideo spiculae in ordines 4 videtae (in paucis racemis pedicelli solitarii ideo spiculae certe bi-seriatae), hirsuti, pilo longo-rigido albo, apices disciformes infra. *Spiculae* ad 4.5 mm. longae, ovato-lanceolatae, aristato-acuminatae, pallido-viridae, internodiis basin globuliformis, viridis, vel purpureus, .5 mm. diam.; gluma inferior membrana, cupula redacta: gluma superior ovato-acuminata, mucronis scabridis .4 mm. longis, 5-sub 7-nervis, quadrans superior glabrus, cetera sparse hirsuta. *Valva inferior* gluma superior similis, sed parce brevior, vacua, ne quidem palea minuta. *Valva superior* angusto-ovata, glabra sed apice minute hirsuto, tenuiter rugosa, 2.5 mm. longa, 5-nerva, mucrona scabrida, .4 mm. longa; palea angusta, plana, dorso tenuiter rugoso; antherae 3, purpureascentes. Styli distincti.

Gezina, near Pretoria, *Fouche* 7. The plants growing in the open veld have been much browsed by stock and they exhibit a prostrate habit with ascending branched culms, while those within a fenced area where the land has been ploughed are taller; simple and erect, or only slightly geniculate from the lowest node.

This plant is very near to *E. acrotricha*, Hack. It differs from Hackel's description of that species in having longer spikelets and a longer panicle with more numerous racemes, spikelets in most of the racemes distinctly 4-seriate, and the mid-nerve of the leaf grooved above and prominent below. I have not seen an authentically named specimen of *E. acrotricha*, and it may be, as Dr. Stapf suggests, a polymorphic type that will include this species as well as *E. ramosa*, but this point can only be settled by the study of a wide range of material.

This is the first record that we have of the genus occurring in the Union of South Africa. It appears to be a good fodder grass and stock apparently relish it.

XXXI. UROCHLOA, Beauv. (166A).

Spikelets broad-ovate to elliptic- or lanceolate-oblong, awnless, usually more or less flattened or slightly depressed abaxially, convex on the back, falling entire from the pedicels solitary or 2-nate or in fascicles of 3-4, secund and abaxial on the more or less triquetrous rarely strap-shaped rachis of moderately slender or somewhat stout spiciform racemes; lower floret ♂ or barren, with a well-developed valvule; upper floret ♂. Glumes similar and subequal or more often dissimilar and very unequal in length, the lower being the shorter, upper resembling and more or less equalling the valve of the lower floret, 5-11- (mostly 7-nerved), with the nerves evenly distributed. Lower floret: valve 5-7-, rarely more-nerved, the inner lateral nerves somewhat distant from the mid-nerve; valvule subequal to the valve, with well-developed inflexed flaps and sharp, sometimes marginate, keels. Upper floret: valve elliptic to rotundate-elliptic in outline, very obtuse with a usually scabrid or barbellate mucro, crustaceous with firm margins, narrowly involute, faintly 5-7-nerved; valvule almost as long as the valve, 2-keeled, its sides tightly embraced by the valve all along, of the same texture as the latter. Lodicules 2, small, broadly cuneate. Stamens 3. Styles distinct. Stigmas plumose, laterally exerted upwards. Grain tightly enclosed by the more or less hardened valve and valvule, broadly to rotundate-elliptic, dorsally compressed, more or less flattened on both faces with broadly rounded

sides; hilum subbasal, punctiform; embryo half to over three-quarters the length of the grain.

Perennial or annual; blades linear to lanceolate, flat; ligules reduced to a ciliate rim; racemes sessile or subsessile on a common axis, simple or nearly so; rhachis more or less triquetrous with a low almost straight or zigzag facial angle or keel, rarely strap-shaped; pedicels solitary or in pairs, alternately to the right and the left of the facial angle, usually reduced to short disc-tipped stumps; or if 2-nate, the primary slightly larger; spikelets close, contiguous or slightly discontinuous, 2- or irregularly pluri-seriate, glabrous or hairy.

U. trichopus, Stapf (in Prain, Flor. Trop. Afr., IX, 589 = *Panicum trichopus*, Hochst., Stapf in Dyer, Flor. Cap., VII, 391).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 398; Pienaars River Station, Pretoria District, *Burt-Davy* 7042; Hector Spruit, *Kirk*.

"Only grows on old lands, and along roads, etc., but has the merit of being the first grass to become fresh and green in the early spring before the rains."—*Kirk*.

U. pullulans, var. **mosambicensis**, Stapf (in Prain, Flor. Trop. Afr., IX, 580).

Mosdene, Naboomspruit, *Galpin*; Rooikop, Bushveld, Pretoria District, *Pole Evans* 155; Hamans Kraal, Pretoria District, *Pole Evans* 394; Pienaars River, Pretoria District, *Pole Evans* 165.

"Very drought resistant; . . . stock are very fond of it in the growing state; . . . it has a spreading habit, except where it comes up very thickly, and there it grows erect."—*Galpin*.

U. helopus, Stapf (in Prain, Flor. Trop. Afr., IX, 595 = *Panicum helopus*, var. *glabrescens*, K. Schum., Stapf in Dyer, Flor. Cap., VII, 392).

Johannesburg, *Bryant* B. 6; Pretoria, *Burt-Davy* 789; Smitskraal, Bloemhof, *Burt-Davy* 12919; Potgietersrust, Waterberg, *Burt-Davy* 2224 and *Galpin* 8866; Hamanskraal, near Pretoria, *Burt-Davy* 1094; Shilovane, Zoutpansberg, *Junod* in *Govt. Herb.* 4682.

An annual weed, known as "sinande."

U. bulbodes, Stapf (in Prain, Flor. Trop. Afr., IX, 593).

Pyramid Estate, Potgietersrust, in moist shady places, *Galpin* 8868; Potgietersrust, *Burt-Davy* 2307.

XXXII. BRACHIARIA, Gris. (166b.)

Spikelets oblong to ovate- or elliptic-oblong, awnless, usually more or less flattened or slightly depressed, convex on the back, falling entire from the pedicels, 1-2-nate, more rarely in fascicles of 3 or more, secund and abaxial on the triquetrous or flat rhachis of moderately slender or stout mostly spiciform racemes; lower floret ♂ or barren, with a usually well-developed valvule, very rarely the latter suppressed. Glumes dissimilar and mostly very unequal in length, the lower being the shorter, upper resembling, and more or less equalling, the valve of the lower floret, 5-7- (very rarely 9-) nerved, with the nerves evenly distributed. Lower floret: valve 5-, very rarely 7-nerved, the lateral nerves placed towards the margins and distant from the middle nerve; valvule usually only slightly shorter than the valve, with well-developed inflexed flaps or the latter vanishing above the middle. Upper floret: valve oblong to elliptic in outline, emucronate, though sometimes contracted into a small projecting point, more or less crustaceous or subcoriaceous with firm margins, narrowly involute, faintly 5-nerved; valvule almost as long as the valve, 2-keeled, its sides tightly embraced by the valve all along, of the same texture as the latter. Lodicules 2, small, broadly cuneate. Stamens 3. Styles distinct. Stigmas plumose, laterally exerted from the upper part of the spikelet. Grain tightly enclosed by the more or less hardened valve and valvule, broad, oblong, or elliptic, dorsally compressed, more or less flattened on both faces with broadly rounded sides; hilum subbasal, punctiform; embryo half to over three-quarters the length of the grain.

Perennial or annual; blades linear to lanceolate, usually flat; ligules reduced to a narrow ciliate or ciliolate rim; racemes usually sub-sessile and solitary on a common axis.

sometimes bare at the base owing to the arrest of spikelets, rarely truly peduncled and panicle (B. *pooveoides*), simple or compound near the base, rarely to or beyond the middle; rhachis filiform, triquetrous or more or less flattened and herbaceous with a wavy or zigzag midrib, which projects as a mostly acute keel on the face; pedicels solitary or in pairs, alternately to the right and the left of the facial angle or the midrib, if solitary, all short or very short, if paired, the primary slightly to very much longer; spikelets closely appressed, always 2-seriate in the plan, but frequently becoming 1-seriate by the dovetailing of the alternate spikelets of the closely approximate ranks, more or less contiguous with their sides or imbricate forming dense spike-like racemes, or distant by almost their own length or more, glabrous or hairy.

B. brizantha, Stapf (in Prain, Flor. Trop. Afr., IX, 531 = *Panicum brizanthum*, Hochst., Stapf in Dyer, Flor. Cap., VII, 386).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 399; Warmbaths, Waterberg District, *Burt-Davy* 1151; near Komatipoort, Black M'belusi to Komati, Barberton, *Burt-Davy* 10658 and 285; Skinners Court, near Pretoria, *Burt-Davy* 3555; Zeerust, Marico, *Hughes* in *Nat. Herb.* H. 321; also in tropical Africa.

A very promising pasture grass; birds are very fond of the seed, and on this account it is difficult to collect sufficient for sowing. "Large-seeded millet grass," frost-resistant.

B. mutica, Forsk. (?) (Stapf ex Prain, Flor. Trop. Afr., IX, 527).

Groenkloof, Pretoria, *Stent* in *Nat. Herb.* H. 21501.

This grass, which originally came from Warmbaths and which has been designated "Warmbaths grass," has never produced flowers. It, however, makes a dense and luxurious mass of vegetation, the details of which agree in every respect with the description of *Brachiaria mutica*, Forsk.; it also resembles exactly plants of Para grass of which we received roots some two years ago from Australia under that name.

The Penhalonga grass of Rhodesia is also apparently the same grass, and has never flowered either here or in Rhodesia.

It is a late grass, and susceptible to cold; can be recommended for soil erosion on account of its long runners.

B. nigropedata, Stapf (in Prain, Flor. Trop. Afr., IX, 535 = *Panicum nigropedatum*, Munro, Stapf in Dyer, Flor. Cap., VII, 388).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 397; Hamanskraal, Pretoria District, *Burt-Davy* 1096; Springbok Flats, Waterberg, *Reck*, *Gort. Herb.* 3531; Kaffraria, Bloemhof, *Burt-Davy* 12974; Vierfontein to Schweizer Reneke, *Burt-Davy* 1625.

Krul gras—a sweet grass much relished by all stock.

B. serrata, Stapf (in Prain, Flor. Trop. Afr., IX, 537 = *Panicum serratum*, Spreng., Stapf in Dyer, Flor. Cap., VII, 388).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 395; near Nylstroom, Waterberg, *Burt-Davy* 2030; Downs, Zoutpansberg, *Junod* 4090; Skinners Court, near Pretoria, *Burt-Davy* 3811; Bushveld, Pretoria, *Reck*, *Gort. Herb.* 3531; Johannesburg, *Bryant* D. 33, C. 13; Beginsel, Standerton, *Burt-Davy* 1019; Heidelberg, *Burt-Davy* 3145; Vereeniging, *Burt-Davy* 7026; Schweizer Reneke, *Burt-Davy* 1666; Leeuwpoot, Carolina, *Burt-Davy* 7426; Potchefstroom, *Burt-Davy* 1061.

Of little value.

B. Isachne, Stapf (in Prain, Flor. Trop. Afr., IX, 552 = *Panicum Isachne*, Rott., Stapf in Dyer, Flor. Cap., VII, 390).

Springbok Flats, Waterberg, *Burt-Davy* 1140; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4392; Skinners Court, near Pretoria, *Burt-Davy* 588; Beginsel Farm, Standerton, *Burt-Davy* 1793; Waterval Boven, *Burt-Davy* 1444; Vlakfontein, near Amersfoort, Wakkerstroom, *Burt-Davy* 4055; Kolonies Plaats, Potchefstroom, *Burt-Davy* 9094.

A troublesome annual weed.

B. Marlothii, Stent. comb. nov. (= *Panicum Marlothii*, Hack., Stapf in Dyer, Flor. Cap., VII, 390).

Schweizer Reneke, Bloemhof, *Burt-Davy* 1672; Kaffraria, near Christiana, Bloemhof, *Burt-Davy* 11414.

B. arrecta, Stent, comb. nov. (= *Panicum arrectum*, Nees, Stapf in Dyer, Flor. Cap., VII, 393).

Spionkop, Lake Chrissie, Ermelo, *Bott and Bell* in *Nat. Herb.* H. 3401.

XXXIII. **ECHINOCHLOA**, Beauv. (166c).

Spikelets ovate to elliptic- or lanceolate-oblong, usually cuspidate or awned, very convex on the back, flat or slightly depressed in front, falling entire from the pedicels, 2-nate or clustered, secund and abaxial on the triquetrous rachis of racemosely arranged false spikes; lower floret ♂ or barren, upper floret ♀. Glumes unequal, membranous, the lower much shorter, more or less ovate from a clasping base, 3-5-nerved, often mucronate, the upper corresponding in length and outline to the spikelet (as seen from the back), very concave, 5-7-nerved, acute, cuspidate or cuspidate, rarely produced into a short awn. Lower floret equalling the upper glume (excluding cusps or awns); valve very similar to the upper glume, but flat or depressed on the back and often with a more pronounced cusp or an awn; valvule equal to the body of the valve, or in barren florets more or less reduced, hyaline, finely 2-keeled. Upper floret: valve ovate to elliptic-oblong, apiculate or obtuse, very convex on the back, subcoriaceous or crustaceous, polished, faintly 5-nerved, margins firm, involute up to near the tip, then flat, not embracing the tip of the valvule; valvule subequal to the valve and similar in substance, with rounded keels and flaps which thin out towards the flat slightly recurved tips. Lodicules 2, cuneate, fleshy. Stamens 3. Styles distinct. Stigmas plumose, exerted from near the tips. Grain broad-elliptic, dorsally flat, ventrally convex; hilum punctiform, subbasal; embryo elliptic-oblong or ovate, exceeding the middle of the grain.

Perennial, and often tall, or annual; blades linear from a slightly constricted or equally wide rarely much attenuated base; ligules 0 or represented by a transverse fringe of hairs; panicles of crowded or loosely arranged secund spiciform branches mostly bearing spikelets from the base or near it; spikelets more or less hispid on the nerves of the glumes, and the lower valve, rarely almost smooth.

E. Holubii, Stapf (in Prain, Flor. Trop. Afr., IX, 606 = *Panicum Holubii*, Stapf in Dyer, Flor. Cap., VII, 394).

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 396; Vereeniging, *Brandmuller* 29; Vaal River, Heidelberg, *Burt-Davy* 12766; vleis at Hamanskraal, near Pretoria, *Burt-Davy*, 1645.

A grass of moist situations, much relished by cattle.

E. stagnina, P. Beauv. (Stapf in Prain, Flor. Trop. Afr., IX, 617 = *Panicum stagninum*, Retz., Stapf in Dyer, Flor. Cap., VII, 394).

Potchefstroom, *Smith* in *Nat. Herb.* H. 13975; Johannesburg, *Kling* in *Nat. Herb.* H. 8793; Pyramid Estate, near Potgietersrust, Waterberg, *Galpin* 8909.

E. colona, Link. (Stapf in Prain, Flor. Trop. Afr., IX, 607).

Potchefstroom, *Bell* in *Nat. Herb.* 7635; Hartbeestdam, near Pretoria, *Pole Evans* 699.

An annual, probably introduced, not very common, valued in India as a fodder-grass and for the grain, which in some districts is used by the natives for food.

XXXIV. **PANICUM**, Linn. (166).

Spikelets lanceolate to oblong, elliptic or orbicular in outline, symmetrical in profile, rarely somewhat oblique, falling entire or almost so, from the often elongated pedicels of a compound or decompound panicle, without a definite orientation towards the axis;

lower floret ♂ or barren, and then usually with a distinct valvule; upper floret ♀. Glumes more or less herbaceous-membranous; the lower usually shorter than the upper (often by very much), rarely equaling it, usually with 1 or more nerves or if very small nerveless; upper as long as the spikelet, rounded on the back, 5-9-nerved. Lower floret: valve very similar to the upper glume and equally rounded and curved on the back (rarely less when the spikelets become oblique), 5-9-, rarely 3- or 11-nerved; valvule thinly membranous to subhyaline, sub-equal to the valve or more or less reduced, rarely suppressed. Upper floret: valve subcoriaceous to coriaceous with firm margins, obtuse to subacute, emucronate (see 1, *P. meyerianum*), faintly nerved; valvule subequal to the valve and of similar substance, tightly embraced by the more or less involute margins of the valve. Lodicules 2, broadly cuneate. Stamens 3. Styles distinct. Stigmas laterally exerted near the tip of the floret. Grain tightly enclosed by the hardened valve and valvule, dorsally compressed bi-convex to almost plano-convex; scutellum elliptic to ovate-elliptic, about half as long as the grain; hilum subbasal, punctiform.

Perennial or annual; rarely suffrutescent, of varying habit and size; leaf-blades mostly linear to linear-lanceolate, but in some groups ovate or filiform to subulate; panicles usually much divided and at least temporarily open; spikelets usually loosely scattered, glabrous or hairy.

P. deustum, Thunb.

The Pyramids, Pretoria, *Maynard* 21; Pilansberg, Rustenburg, *Gibbs* in *Nat. Herb.* H. 3442.

P. maximum, Jacq.

Skinnars Court, Pretoria, *Burt-Davy* 3807; Pretoria, *Burt-Davy* 2190; Bushveld, Pretoria, *Reck* in *Nat. Herb.* H. 178; De Kroon, Brits, Rustenburg, *Stent* in *Nat. Herb.* H. 6121; Crocodile River, *Burt-Davy* 9284; Spitzkop Farm, Ermelo, *Burt-Davy* 13631; Warmbaths, Waterberg, *Burt-Davy* 1147; Pyramid, near Potgietersrust, *Galpin* 8871; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 403; Springbok Flats, Waterberg, *Burt-Davy* 1136; Waterval Boven, Lydenburg District, *Burt-Davy* 1417; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4450.

The "buffels gras" of the bushveld, more commonly known in other parts of the world as Guinea grass; an excellent pasture and hay grass.

The degree of hairiness varies considerably. *Burt-Davy* 1417 has very hairy sheaths and culms, and rather large, glabrous, purple spikelets. *Reck*, in *Nat. Herb.* H. 178 and *Galpin* 403 exhibit a glaucous form with smaller, greenish, pubescent spikelets, which may prove to be specifically distinct when more material can be collected and examined.

P. laevifolium, Hack.

Johannesburg, Observatory Hill, *Bryant* B. 41; Vereeniging, *Burt-Davy* 7518; Warmbaths, Waterberg, *Burt-Davy* 1152; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 401; Nylstroom, Waterberg, *Driver* in *Nat. Herb.* H. 8947 and *Hilliard* in *Nat. Herb.* H. 4124; Pietersburg, Zoutpansberg, *Van Rooyen* in *Nat. Herb.* H. 9034; Barberton, *Lea* in *Nat. Herb.* H. 5875; Middelburg, *Ballard* in *Nat. Herb.* H. 6307; Schweizer Reneke, *Burt-Davy* 1677 and 1623; Ermelo Experiment Farm, *Burt-Davy* 3390; Teakworth, Standerton District, *Stent* in *Nat. Herb.* H. 19941; Platrand, Standerton, *Burt-Davy* 2356; Potchefstroom, in vlei lands, *Burt-Davy* 1071, 2375; Belfast, Lydenburg, *Burt-Davy* 1380; Christiana, Bloemhof District, *Burt-Davy* 12790.

"Roll gras" or "Blaauwzaad gras" of the Orange Free State.

An annual sweet grass, valuable as a hay and pasture grass, very fast growing. It is sometimes attacked by the parasite *Striga lutea*. *Stent*, H. 19941, shows a specimen of the *Panicum* bearing *Striga lutea* on its roots—both host and parasite are in a flourishing condition; there were, however, patches on the lands on which I collected this specimen where the *Panicum* was dwarfed and stunted, and not more than a few inches high on account of the attacking parasite.

P. longijubatum, Stapf (in Prain, Flor. Trop. Afr., IX, 718 = *Panicum proliferum*, var. *longijubatum* and var. *paludosum*, Stapf, in Dyer, Flor. Cap., VII, 406).

Pyramid Estate, Potgietersrust, *Galpin* 8872.

P. coloratum, Linn.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 405; near Station, Warmbaths, Waterberg District, *Burt-Davy* 860; Kolonies Plaats, Potchefstroom, *Burt-Davy* 9083; Machavie Station, between Potchefstroom and Klerksdorp, *Burt-Davy* 1480; Kaffraria, Christiana, Bloemhof, *Burt-Davy* 12869, 14125, 14134; Bloemhof, *Burt-Davy* 1509; Smitskraal, Bloemhof, *Burt-Davy* 12924.

P. minus, Stapf.

Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21125; Christiana, Bloemhof District, *Burt-Davy* 11711.

Var. **planifolium**, Stapf.

Kaffraria, Christiana, Bloemhof District, *Burt-Davy* 14137; Potchefstroom, in black loam near furrow, *Burt-Davy* 1211.

"A useful grass; very closely resembling forms of *P. coloratum*."—*Burt-Davy*.

P. dregeanum, Nees.

Pyramid Estate, near to Potgietersrust, *Galpin* 8870.

P. natalensis, Hochst.

Groenkloof, Pretoria, *Phillips* H. 21686; Johannesburg, *Bryant* D. 13; Observatory Ridge, Johannesburg, *Burt-Davy* 3096; Pretoria, *Macdonald* in *Nat. Herb.* H. 5464; Henley-on-Klip, *Stent* in *Nat. Herb.* H. 2112; Billy's Vlei, Ermelo, *Burt-Davy* 9312; one mile north of Robinsons, Carolina, *Burt-Davy* 2973; Heidelberg, *Burt-Davy* 3143.

P. Ecklonii, Nees.

Belfast, *Stent* in *Nat. Herb.* H. 10627.

P. novemnerve, Stapf (in Prain, Flor. Trop. Afr., IX, 702).

In sandy places near Mara, north Zoutpansberg, *Schlecht.* 4614.

Dr. Stapf, in the Flor. Trop. Afr., IX, 705, has created a separate species for *Schlechter* 4614, and described it under the name of *Panicum arcuatum*. I cannot, however, find sufficient differences between *Schlechter*'s specimen and those Dr. Stapf has classified under *P. novemnerve* to warrant such distinction, and his description of the latter grass agrees well with *Schlechter*'s specimen in which the upper glume is more often than not 9-nerved. The arrangement of the spikelets is also not a constant character.

P. miliaceum, L.

Groenkloof, Pretoria, *Turner* in *Nat. Herb.* H. 10013; Pyramid, Pretoria, *Maynard* 21; Rustenburg, *Weeber* in *Nat. Herb.* 6621.

Indian buffels grass. Cultivated; there is no records of this grass occurring spontaneously in the Transvaal, except perhaps as an escape from cultivation.

P. obscurans, Stapf (*ex* Woodrow in Journ. Bomb. Nat. Hist. Soc., XIII, 434, 1901).

Pretoria, *Franks* in *Nat. Herb.* H. 10616; Wonderboompoort, Pretoria, *Mogg* in *Nat. Herb.* H. 9937; Crocodile River, *Burt-Davy* 9290.

The culms of this grass break off at the top node when the seeds are ripe, and the widely spread, straw-coloured panicle is blown over the veld. The seeds contain a great quantity of oil, and were responsible for seriously impeding the movements of trains, principally in the Pretoria, Witwatersrand, and Standerton Districts, where the tops of the grass were constantly being blown in great masses on to the railway line.

Another report of strange behaviour on the part of this *Panicum* comes from Krugersdorp, which suddenly received a shower of this grass which fell on the streets of the town like manna from the heavens above. The probable explanation of this phenomenon was

that masses of the grass had been collected in the veld by a passing whirlwind, commonly known in this country as a "dust devil," and this had carried the grass along, whirling it high into the clouds, and finally releasing it over Krugersdorp.

XXXV. *DIGITARIA*, Hall (166D).

Spikelets lanceolate to elliptic-oblong, flat or more or less concave in front, convex on the back, falling entire from the pedicels, usually 2-3- (more rarely 4-6-) nate, second and abaxial on the triquetrous or flat rhachis of slender to very slender spiciform racemes; lower floret barren, reduced to the valve and a very minute rudimentary valvule, accompanied by equally minute lodicules; upper floret usually equal to the lower, ♂. Glumes very dissimilar, the lower a minute membranous scale or a very delicate readily disintegrating hyaline membrane, or sometimes quite suppressed, upper membranous, sometimes very thin, as long as the fertile floret to whose back it is appressed, or shorter to very short, 5- or more often 3-nerved, very rarely 1-nerved or almost nerveless, or 0. Barren floret: valve sometimes resembling the upper glume, but generally larger and determining the outline and size of the spikelet, 7-5-nerved; very rarely 3-nerved or reduced to a small scale, nerves parallel, prominent if not obscured by hairs. Fertile floret: valve chartaceous to subchartaceous with thin hyaline margins, faintly 3-nerved; valvule equal or subequal to the valve, embraced by it all along, finely 2-nerved, of the same texture. Lodicules 2, minute, broadly cuneate. Stamens 3. Styles distinct. Stigmas laterally exerted near the top of the floret. Grain tightly enclosed by the slightly hardened valve and pale, oblong, dorsally more or less compressed, plano-convex in cross-section, with rounded sides; hilum subbasal, punctiform; embryo usually less than half the length of the grain.

Perennial or annual, very rarely suffrutescent; blades linear to linear-lanceolate, usually flat and often more or less flaccid; ligules membranous; racemes sessile, digitate or more or less distant on a common axis, sometimes bare at the base owing to the arrest of spikelets, rarely truly peduncled and panicle, simple or sometimes compound near the base; rhachis triquetrous, lateral angles with narrow green margins or herbaceous wings or flattened with a terete midrib; pedicels unequally long; spikelets closely appressed and more or less imbricate or lax to very lax, usually silky though often apparently glabrous owing to the very tight application of the extremely fine hairs, rarely really glabrous.

D. monodactyla, Stapf.

Pretoria, *Macdonald* in *Nat. Herb.* H. 5408; Belfast, Middelburg, *Burt-Davy* 1345; Botsabelo, *Schlechter* 3781.

Var. **explicata**, Stapf (*ex* Stapf in Prain, *Flor. Trop. Afr.*, IX, 442).

Billy's Vlei, Ermelo, *Burt-Davy* 9254.

Macdonald and Burt-Davy's specimens, which I have classified as belonging to the type as limited by Stapf in the *Flor. Trop. Afr.* on account of the narrow convolute leaves, both have some of the basal leaves flat and with well-developed motor cells, and occasionally three lateral ribs on either side of the midrib.

D. argyrograpta, Stapf.

Potchefstroom Farm, Potchefstroom, *Burt-Davy* 1068; Kaffraria, Christiana, Bloemhof, *Burt-Davy* 14148; Springbok Flats, Waterberg, *Burt-Davy* 1141.

D. eriantha, Steud.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 393; Downs, Zoutpansberg, on open veld, *Junod* 4086; Uitgevallen, Heidelberg, *Burt-Davy* 13666; Vierfontein, Schweizer Reneke, Bloemhof, *Burt-Davy* 1639; Potchefstroom, *Burt-Davy* 11459.



Plate V.

Digitaria Smutsii, Stent.

A, Base of leaf with ligule; B, Pair of spikelets; C, Glumes; D, Lower valve; E, Scale like pale; F, Upper valve; G, Pale; H, Pistil and stamens.

Var. *stolonifera*, Stapf.

Potchefstroom, *Burt-Davy* 2376; Hamanskraal, near Pretoria, *Burt-Davy* 1091.

A useful pasture grass. "Finger gras"; one of our best sweet grasses.

D. Smutsii, Stent, n. sp.

Gramen perenne, dense caespitosum, rhizoma obliquum, gemmis innovationalibus. *Culmi* erecti, geniculati ad 1.5 m. alti, glabri; nodi plurimi ramos florentes ferentes. *Foliorum vaginæ* glabrae, vel inferiores pilis tuberculis insidentibus conspersa, infimæ basis tomentosae. *Ligulae* membranaceae, 2-3.5 mm. longae. *Laminae* lineares, apice setaceo-acuminatae, ad 12 mm. latae, 50 cm. longae, culmorum folia plerumque breviora, illa innovationum longiores angustiora, laeves, vel marginibus parce asperis, planae, subtus carinatae, nervis lateralibus primariis utrimque 5-7. *Racemi* 4-10, plerumque 8-10, sessiles, solitarii vel infimi verticillati ex axe communis, plerumque late patentibus, ad 15 cm. longi, plumbeo-viridi vel purpurentes, inferiores saepe e basi ramis brevibus; rachis tenuis .5 mm. lata, anguste marginata, supra carinata; pedicelli binati, aliquando ternati, vel solitarii, longissimos, ad circa 7 mm. longos. *Spiculae* appressae, subimbricatae, anguste ovato-lanceolatae, 3.5 mm. longae, 7 mm. latae; *gluma inferior* brevissima, membranacea, enerva; *gluma superior* angusta, circa 1-2 mm. longa, 3 nervis, pilis bombycinis inter nervoset in marginibus; *palea* squama hyalina .5 mm. longa, rare longiore.

Doornkloof, Irene, near Pretoria, *Pole Evans* 217; Division of Botany, Pretoria (cult.), *Stent in Nat. Herb.* H. 21418; Irene, *Burt-Davy* 3839; Uitgevallen, Heidelberg, *Burt-Davy* 13666; Hamanskraal, near Pretoria, *Burt-Davy* 1089.

Near to *D. eriantha*, Steud., but a taller, coarser grass, with much-branched culms; broader leaves, with 5-7 distinct lateral nerves and a prominent keel; racemes nearly always widely spreading on an elongated axis and with numerous short additional racemes at the base of the lower ones. The spikelets are often ternate, sometimes 4-5 on short branches, and the pedicels are usually long.

This grass has been frequently collected and distributed as *D. eriantha*. I had seen only the dried material till General Smuts drew my attention to it growing on his farm at Irene, and the difference in habit and general appearance between it and *D. eriantha* was obvious. It has since been cultivated at the Dry-land Station at Groenkloof, where it has done well and is proving a very promising grass.

D. setivalva, Stent, n. sp.

Gramen perenne, stolonifer. *Culmi* erecti, vel geniculati, 46-107 cm. alti, 3-5-nodis, simplices vel plerumque nodis ramatis; nodus superior longissimus vagina longa exsertus. *Foliorum vaginæ* tenuae, laxae, viridae, gracile sulcatae, hirsutae, pilis longis albis e tuberculis patentibus (*Galpin* M. 390 and 394) vel omnino glabrae (*Galpin* M. 392) vel tantum pauci nodi hirsuti, ceteri glabri (*Burt-Davy* 864); *ligulae* tenuae membranaceae, ovatae, integrae, vel tandem fimbriatae, ad 4 mm. longae. *Laminae* lineares, laevolatae, base tenuite constrictae et rotundatae, in apicem setaceum attenuatae, glabrae vel exigue hirsutae, scaberulae; margines cartilaginei, stricti vel aliquantum crispi; costa triua, nervis lateralibus inconspicuis. *Racemi* 5-11, digitati vel sub-digitati, erecti vel tenuites patenti, stricti, ad 20 cm. longi, rachis strictus vel tenuite flexuosus, marginibus argute alatis, scaber, 6 mm. latus. *Spiculae* appressae, 3 mm. longae, haud imbricatae, acute acuminatae, pallide viridae vel flavescentes, hirsutae; *gluma inferior* parvissima obtusa, enerva; *gluma superior* angusta, acute acuminata, circa 2 mm. longa, 3-nervis, pilis appressis costa utrinque et cum marginibus longe molliteque ciliatis; *valva inferior* spiculam aequans, 7-nervis, pilis tenuis obtusis inter nervos e marginibusque; ordines interiores pilorum cum setis validis albidis vel flavibus nitentibus, acutis, mixti, setae pilique tamdem patenti; *lodiculæ* et *palea* minutae; *valva superior* 3-nervis, valvam inferiorem fere aequans, sub-apiculata.

Mosdene, Naboomspruit, in black turf, *Galpin* M. 392; on loam formation, *Galpin* M. 390, 394, 391; Warmbaths, *Burt-Davy* 864; Springbok Flats, *Burt-Davy* 1122.

This resembles *D. marginata*, var. *fimbriata*, Stapf (*ex descr.*), but is a more robust grass and a perennial; the thick-walled, sharp-pointed bristles on the lower valves are constant in all the material examined. In *Burt-Davy* 1122, *Galpin* M. 390, 394, and 391, they are bright golden, while in *Burt-Davy* 864 and *Galpin* M. 392 they are whitish with a tendency to become yellow in some of the spikelets.

Stapf refers *Burt-Davy* 1122 to *D. eriantha* (Prain, *Flor. Trop. Afr.*, IX, 430), and notes that it is the only specimen he has seen with the stiff yellow bristles.

The same grass occurs in Southern Rhodesia (*Mundy* 2830), and Mr. Walters writes of it: "It is a native Rhodesian grass and when grown alone covers the ground well, being both perennial and stoloniferous. It has shown great promise as a pasture constituent."

D. ternata, Stapf.

Belfast, Lydenburg, *Burt-Davy* 1375; Spionkop, Ermelo, *Burt-Davy* 9238; Volksrust, Wakkerstroom, *Burt-Davy* 7732.

D. debilis, Willd.

Shilovane, Zoutpansberg, *Junod* in *Nat. Herb.* H. 367.

D. horizontalis, Willd.

Potgietersrust, Waterberg, *Burt-Davy* 2233; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* 368.

D. marginata, Link. (Stapf in Prain, *Flor. Trop. Afr.*, IX, 439).

Pretoria, *Burt-Davy* 886; Rietvlei Farm, Belfast, Lydenburg, *Burt-Davy* 1262.

D. longiflora, Pers. (Stapf in Prain, *Flor. Trop. Afr.*, IX, 469).

Downs, on veld, Zoutpansberg, *Junod* 4100; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 378.

D. uniglumis, Stapf (Stapf in Prain, *Flor. Trop. Afr.*, IX, 474).

Downs, Zoutpansberg, *Junod* 4093; Meintjes Kop, Pretoria, *Burt-Davy* 3549.

I have not seen an authentically named specimen of this species, but the specimens quoted above agree with Stapf's description, except that the spikelets are slightly larger (2.25 mm.) and the roots and bases of the culms are densely covered with silky tomentose cataphylls, and the sheaths do not appear to break up into fibres. *Schlechter* 6801 from Sand River, distributed as *D. diagonalis*, is the same grass from a very different locality. It is possible that both the above grasses should be placed under Stapf's *D. diagonalis*, but the localities suggest rather an extension of the tropical species. More material is necessary before I can decide definitely on this point.

D. tricholaenoides, Stapf.

Leeuwpoot, Carolina, *Burt-Davy* 7445; Ermelo, *Burt-Davy* 5464; Spitzkop, *R. Pott* 5206; Amersfoort, *Burt-Davy* 9049.

Good fodder grass, relished by sheep.

D. Brazzae, Stapf (in Prain, *Flor. Trop. Afr.*, IX, 447).

Houghton Estate, Johannesburg, *Bryant* C. 37.

XXXVI. **TRICHOLAENA**, Schrad. (168).

Spikelets oblong, laterally compressed, more or less gaping, panicked, deciduous from capillary pedicels; lower floret usually ♂ or barren, upper ♀. Glumes very dissimilar or at least unequal, lower reduced to a minute scale or obsolete, very rarely half the length of the spikelet, somewhat remote from the upper; upper glume membranous, (marginate, mucicous or finely mucronate or aristate from the sinus, 5-nerved, usually hairy. Valves very dissimilar, lower like the upper glume with a hyaline 2-nerved subequal pale, upper much smaller, thinly chartaceous, glabrous, shining, obtuse or submarginate, obscurely 5-nerved, with an equal 2-nerved pale of similar substance. Lodicules 2, very small. Stamens 3. Style free, slender. Stigma densely plumose, laterally exerted. Grain



Plate VI.

Digitaria seticalva, Stent.

A, Lower glume; B, Upper glume; C, Lower valve; D, Fertile floret.

oblong-ellipsoid, closely embraced by the valve and pale; hilum basal, punctiform; embryo about half the length of the grain.

Perennial, rarely annual; blades linear to setaceous; ligule a fringe of hairs; panicles open or contracted, often much divided, with capillary branchlets and pedicels. the latter with thickened tips; spikelets often completely enveloped by soft shining hairs, rarely pubescent or quite glabrous.

T. setifolia, Stapf.

Johannesburg, Observatory Hill, *Bryant* B. 36; Houghton Estate, *Bryant* C. 23; Pretoria, Wonderboom, *Burtt-Davy* 8059; The Willows, *Burtt-Davy* 4488; Eloff's Plantation, *Macdonald* in *Nat. Herb.* H. 5440; near Irene, *Burtt-Davy* 30; Barberton Mts., *Burtt-Davy* 337; Belfast, Lydenburg, *Burtt-Davy* 1378; Waterval Boven, *Burtt-Davy* 1423; Schoeman's Kloof, Lydenburg, *Burtt-Davy* 7251; Spitzkop, Sabie, Lydenburg, *Burtt-Davy* in *Nat. Herb.* H. 1573; Billy's Vlei, Carolina, *Mitchell* in *Nat. Herb.* H. 5666; Christiana, Bloemhof District, *Burtt-Davy* 13227.

T. rosea. Nees.

Bezuidenhout Valley, Johannesburg, *Bryant* B. 1; Skinners Court, Pretoria, *Burtt-Davy* 4233; Schweizer Reneke, Bloemhof, on the kopje, *Burtt-Davy* 1668; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 406; Tzaneen, Zoutpansberg, *Mogg* in *Nat. Herb.* H. 9452; Lydenburg, *Burtt-Davy* 405; Komatipoort, *Burtt-Davy* 375; Mooibank Settlement, Potchefstroom, *Jones* in *Nat. Herb.* H. 170.

Natal Red Top—a good fodder grass.

XXXVII. **XYOCHLAENA**, Stapf (168A).

Spikelets oblong or ovate-oblong, deciduous from capillary pedicels, more or less gaping, not gibbous, glabrous or pubescent; lower floret usually ♂ or barren; upper floret ♀. Glumes very unequal, the lower often reduced to a minute scale; upper 5-nerved, emarginate, mucicous, or minutely mucronate, glabrous or pubescent, rarely shortly villous. Lower valve like the upper glume, with a hyaline 2-nerved subequal pale; upper valve smaller, membranous or chartaceous, glabrous, shining, obtuse or subemarginate, obscurely 5-nerved with an equal 2-nerved pale. Lodicles 2, very small. Stamens 2. Styles free, slender.

Perennial; blades linear to setaceous; ligule a fringe of hairs; panicles open or contracted, often much divided, with capillary branchlets and pedicels. Differs from *Tricholaena* in the non-gibbous upper glume and barren valves, which also lack the enveloping long silky hairs.

X. monachne, Stapf (Hook. Icon. Plt. V, 1, t. 3098 = *Tricholaena glabra*, Stapf, in Dyer, Flor. Cap., VII, 446).

Bushveld, Pretoria, *Reck* in *Nat. Herb.* H. 176.

Xyochlaena, sp. nov. (?)

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 404.

XXXVIII. **OPLISMENUS**, P. Beauv. (169).

Spikelets oblong to ovate-oblong or lanceolate, awned from the glumes, falling entire from the pedicels, in small clusters on a common axis or 2-nate or solitary, second and abaxial on the angular rhachis of racemosely arranged densely or more rarely interruptedly spiciform racemes; lower floret ♂ or barren, with a perfect or reduced or no valvule; upper floret ♀. Glumes similar in shape and texture, subequal, herbaceous to membranous, 3-7 nerved, both, at least the lower, awned, more or less keeled upwards. Lower floret exceeding the glumes; valve resembling them, 5-9-nerved, mucicous or mucronate, or very shortly awned; valvule subequal to the valve, and 2-nerved when perfect. Upper

floret: valve oblong, broadly involute with firm margins, muticous, chartaceous to subcoriaceous; valvule equalling the valve and embraced by the valve all along. Lodicules 2; broadly cuneate, often very delicate. Stamens 3. Styles distinct, long. Stigmas terminally or subterminally exerted, plumose. Grain tightly enclosed by the somewhat hardened valve and valvule, oblong; hilum oblong, one-third to half the length of the grain.

Annual or more usually perennial; often weak grasses with slender many-noded ascending culms, the internodes of which have mostly a decurrent villous line adaxial to the subtending leaf; blades flat, thin, often slightly asymmetrical, often minutely cross-veined; lower and intermediate racemes mostly distant, the upper decreasing upwards and more or less approximate; the awn of the lower glume the longest.

O. africanus, Beauv., var. *simp'lex*, Stapf.

Houtbosch, Transvaal, *Rehmann* 5735*.

O. hirtellus, P. Beauv. (Stapf in Prain, Trop. Afr., IX, 631 = *O. africanus*, Beauv., Stapf in Flor. Cap., VII, 416).

Shilovane, Zoutpansberg, *Junod* in *Nat. Herb. H.* 4675.

XXXIX. **SETARIA**, Beauv. (171).

Spikelets ovate to oblong, falling entire from the pedicel, subtended by 1 to many persistent bristles (modified branchlets), which often form a one-sided involucre (see section *Ptychophyllum*), subsessile in contracted spike-like or more or less open panicles. Lower floret ♂, or reduced to the valve, and a more or less arrested pale; upper floret ♀. Glumes membranous, lower generally much smaller, usually 3-5, rarely 1- or 7-nerved, upper usually 5-, sometimes 7-nerved. Lower valve more or less exceeding and resembling the upper glumes; upper valve chartaceous to coriaceous, 5-nerved. Pales subequal to their valves or that of the lower floret more or less arrested, flat, 2-nerved, hyaline in the lower, of the same substance as the valve in the upper floret. Lodicules 2, broadly cuneate. Stamens 3. Styles distinct. Stigmas laterally exerted. Grain tightly enclosed by the hardened valve and pale, oblong or ellipsoid; hilum basal, punctiform or orbicular; embryo about half as long as the grain.

Perennials or annuals of various habit; ligules usually reduced to a ciliate rim, rarely a distinct membrane; panicle mostly cylindric, spike-like, dense, with the solitary or clustered spikelets on very short branches, which are more or less produced into bristles beyond the spikelets or divided into a one-sided bristly involucre at their base, or more or less open with elongate branches and more distant spikelets, often with or without subtending bristles in the same inflorescence; bristles always persistent.

S. sulcata, Raddi.

Skidders Court, Pretoria (cult. ?), *Burt-Davy* 611; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb. H.* 4460; Shilovane, Zoutpansberg, *Junod* in *Nat. Herb. H.* 4667; Barberton, *Burt-Davy* 267.

Natal buffel grass; valuable pasture grass; horses are especially fond of it. Seeds are harmful to small birds.

S. lindenberiana, Stapf.

Johannesburg, on hillsides at Rosebank, *Bryant* F. 6; Meintjes Kop, Pretoria, *Burt-Davy* 5608; Crocodile River, near Pretoria, *Burt-Davy* 207; slopes of the Magaliesberg, *Fouche* H. 21687.

"Kopje gras"; an excellent pasture grass.

S. nigrirostris, Dur. and Schinz.

Ermelo District, Ermelo, *Tennant* in *Nat. Herb. H.* 6927; Athol, near Amsterdam, *A. C. Macdonald* in *Nat. Herb. H.* 94; Vlaktefontein, Wakkerstroom District, *Burt-Davy* 4155; Beginsel Farm, Standerton, *Burt-Davy* 934; Leeuwpoot, Carolina, *Burt-Davy* 7380; Johannesburg, *Bryant* C. 8; Skidders Court, Pretoria, *Burt-Davy* 791; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* 4382.

An excellent pasture grass for high veld.

S. Gerrardii, Stapf.

Skinnners Court, Pretoria, *Burtt-Davy* in *Nat. Herb.* H. 4367; Struben's Farm, Pretoria, *Burtt-Davy* 4481; Onderstepoort, Pretoria, *Native Aaron* in *Nat. Herb.* H. 7938; Mosdene, Naboomspruit, Waterberg, *Galpin* 408; Potchefstroom, *Burtt-Davy* 2176; Pan Station, Middelburg, *Burtt-Davy* 13336; Uitgevalen, Heidelberg, *Burtt-Davy* 9158; Standerton, *Burtt-Davy* 3316.

S. Holstii, Herrm. (in Rosen, *Beitrag. Biol., Pflanzen* X, 45).

Mosdene, Naboomspruit, *Galpin* M. 407; Malalane, Barberton, *Atmore* in *Nat. Herb.* H. 4727.

S. sp.

Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4393.

Belongs to an undescribed tropical species, the description and name of which will be published by Dr. Stapf.

S. perennis, Hack.

Kopjes at Sunnyside, Pretoria, *W. Macdonald* in *Nat. Herb.* H. 5549; Crocodile and Magalies Rivers, *Burtt-Davy* 202; Koster River, Rustenburg, *Burtt-Davy* 141.

S. flabellata, Stapf.

Eloff's Plantation, Pretoria, *W. Macdonald* in *Nat. Herb.* H. 5433; near Nylstroom, Waterberg, *Burtt-Davy* 2032, 2025; Carolina, *Burtt-Davy* 7375, 7403; Uitgevalen, Heidelberg, *Burtt-Davy* 9171; Ermelo, *Burtt-Davy* 5457; Panplaats, Pan Station, Middelburg, *Burtt-Davy* 13309; Potchefstroom, Machavie Station, *Burtt-Davy* 1479; Welverdiend Station, *Burtt-Davy* 14580; Cawoods Hope, Bloemhof, *Burtt-Davy* 12973.

S. aurea, A. Brown.

Skinnners Court, Pretoria, *Stent* in *Nat. Herb.* H. 4361, 3388, 5830; Johannesburg, *Bryant* C. 87; Sterk Hill, Lydenburg, *Burtt-Davy* 451; Louis Trichardt, Zoutpansberg, *Fripp* in *Nat. Herb.* H. 10358; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4403 and 4398; Ermelo, *Burtt-Davy* 1659.

South African golden millet grass; good hay grass.

S. imberbis, R. & S.

Arcadia, Pretoria, *Burtt-Davy* 11745; Johannesburg, *Bryant* D. 86, B. 13, C. 11; Shilovane, Zoutpansberg, *Junod* in *Nat. Herb.* H. 4674; Kolonies Plaats, Potchefstroom, *Burtt-Davy* 9084; Rietvlei, Belfast, *Burtt-Davy* 1263; Experiment Farm, Ermelo, *Burtt-Davy* 7716.

An annual weed in cultivated lands.

S. italica, Beauv.

Houghton Estate, Johannesburg, *Bryant* C. 35; Standerton, *Burtt-Davy* in *Nat. Herb.* H. 5064.

A cultivated species occurring sometimes as an escape.

S. verticillata, Beauv.

Pretoria, *Burtt-Davy* 1040; Potchefstroom, *Burtt-Davy* 9090, *Eldred* in *Nat. Herb.* H. 4220; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 436; Rietvlei, Belfast, *Burtt-Davy* 1257.

"Klitsgras"; a troublesome weed in cultivated lands.

XL. **PENNISETUM**, Pers. (175).

Spikelets oblong or lanceolate, solitary or in clusters of 2-4, subtended by and deciduous with sessile or peduncled involucre of naked or plumose bristles (rarely reduced to a solitary bristle in section *Beckeropsis*), and arranged round the axis of spike-like, usually cylindric panicles; lower floret ♂ or barren, with or without a pale; upper ♀. Glumes usually small and hyaline, lower sometimes suppressed, upper rarely half the length of the spikelets

or more, and then several- to 7-nerved. Valves equal or subequal, membranous to chartaceous, 5-7-nerved, or the lower more or less reduced, thinner, fewer-nerved. Pales subequal to the valve and of similar texture, 2-nerved, or more or less reduced in the lower floret. Lodicules small, usually in front and outside the pale, or 0. Stamens 3. Styles distinct, slender or connate. Grain enclosed by the slightly changed valve and pale (see also *P. typhoideum*), broadly oblong, slightly dorsally compressed to subglobose; hilum basal, punctiform; embryo large, half-three-quarters the length of the grain.

Perennial or annual; culms simple or often profusely branched; blades flat or convolute; ligules usually reduced to a ciliate rim or a fringe of hairs, rarely membranous; panicle spike-like, usually dense, branches very numerous all around the axis, very short, simple with a solitary spikelet, or scantily divided, with the spikelets in clusters of 2-5; the solitary spikelets or the clusters subtended by and deciduous with an involucre (very rarely a solitary bristle) of often very numerous and usually unequal scabrid or plumose, simple, rarely branched bristles.

P. spicatum, R. & S. (= *P. typhoideum*, in Dyer, Flor. Cap., VII, 432).

Nylstroom, Waterberg, *Burtt-Davy* 1735, 1734.

Cultivated under the names of pearl millet and babala grass.

P. cenchroides, Rich.

Waterberg, near Nylstroom, *Burtt-Davy* 2006; Warmbaths, *Burtt-Davy* 2334;

Mosdene, Naboomspruit, *Galpin* M. 435.

One of several grasses known as buffel grass; useful pasture plant.

P. macrourum, Trin.

Belfast, *Burtt-Davy* 1463.

Occurs also in the Waterberg District, but not very common in the Transvaal.

P. sphacelatum, Dur. & Schinz., var. **tenuifolium**, Stapf.

Wonderboom, near Pretoria, *Burtt-Davy* 8060.

P. Thunbergii, Kunth.

Pinedene, near Pretoria, *Burtt-Davy* 2325; Rustenburg, *Burtt-Davy* 140;

Belfast, *Franks* in *Nat. Herb.* H. 9774.

Var. **Galpinii**, Stapf.

Pretoria, *Burtt-Davy* 830; Vereeniging, *Brandmuller* 6.

XLI. POTAMOPHILA, R. Br. (191).

Spikelets laterally slightly compressed, on short or rather long pedicels, paniced; rhachilla jointed above the rudimentary basal glumes. Florets 3; lower two reduced to minute empty valves, uppermost ♀ or unisexual, with the organs of the other sex reduced. Glumes reduced to very minute rounded or truncate scales, or to an obscure entire or bilobed hyaline rim. Empty valves very small, hyaline, nerveless, subulate or elliptic and rounded or lobed; fertile valve membranous, 5-nerved, awnless, nerves raised, sometimes slightly winged. Pale 3-nerved; otherwise similar to the fertile valve. Lodicules 2, finely nerved. Stamens 6. Styles distinct. Stigmas feathery. Grain obovate, compressed, crowned by the thickened bases of the styles, enclosed by the unaltered glume and pale, free.

Rather tall aquatic grasses; blades flat; ligules membranous; panicle effuse or contracted.

P. prehensiles, Benth.

In bush, Cyprus Point, Zoutpansberg, *Junod* 4103.

XLII. ORYZA, Linn. (193).

Spikelets disarticulating below the glumes, 1-flowered, laterally compressed, and oblong, on the slender branches of an elongate rarely spiciform panicle; pedicels annular at the top.

Glumes scale or bristle-like or suppressed. Valve chartaceous obtuse or acute, awnless or awned, strongly 3-5-nerved, palea as long as the valve, keeled. Lodicules 2-lobed, rarely entire. Stamens 6; anthers linear. Styles short, free. Stigmas laterally exserted. Grain narrow, free or adnate to the valve and palea.

Tall, leafy grasses. Leaves long and flat. Ligule long and firmly membranous.

O. sativa, L.

Mosdene, Naboomspruit, Waterberg, in Nyl Vlei, *Galpin* M. 444.

Escape from cultivation (?).

XLIII. LEERSIA, Sw. (194).

Spikelets laterally compressed, very shortly pedicelled, panicle; rhachilla jointed above the rudimentary glumes. Floret 1, ♂. Glumes reduced to an obscure hyaline, entire, or 2-lobed rim. Valve 5-nerved, subcartilaginous, awnless, keel and margins rigidly ciliate. Pale narrow, 3-nerved, subcartilaginous, except at the hyaline margins, grooved along the outer nerves and tightly clasped by the inflexed margins of the valve, keel rigidly ciliate. Lodicules 2; fleshy, finely nerved. Stamens 6, 3, or 1. Styles distinct. Stigmas feathery. Grain ovate or oblong, compressed, embraced by the valve and the pale, free; embryo short.

Perennial; leaves narrow; panicle usually flaccid, with very slender branches.

L. hexandra, Sw.

Johannesburg, *Bryant* C. 9; Pretoria, *Burt-Davy* 1167; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 400; Warmbaths, *Burt-Davy* 1148; Shilovane, Zoutpansberg, *Junod* in *Nat. Herb.* H. 4677; Beginsel Farm, Standerton, *Burt-Davy* 1809; Vereeniging, *Burt-Davy* 7703, *Brandmüller* 5.

Chiefly found in swampy or moist places. "Water kweek."

"An excellent, nourishing, and palatable grass in its early stages; cattle and horses relish it and grow fat on it."—*Brandmüller*.

XLIV. EHRHARTA, Thunb. (201).

Spikelets laterally compressed, panicle or racemed, sometimes solitary, pedicelled; rhachilla disarticulating below the valves, more or less obscurely produced. Florets 3; lower two reduced to empty valves, uppermost ♂. Glumes persistent, membranous. Valves 3, heteromorphous; the lower two empty, usually exceeding the glumes, more or less cartilaginous, often bearded, and the upper with a callous appendage at the base, awnless or awned; the uppermost fertile, smaller, thinner, awnless, sometimes with a knob-like appendage at the base forming a hinge with the appendage of the upper empty valve. Pale narrow, keeled, finely 2-nerved, nerves very close. Lodicules 2. Stamens 6 or 3, very rarely 1. Styles distinct, short. Stigmas plumose or brush-like, exserted above the base. Grain elliptic, much compressed; hilum a fine line almost as long as the grain; embryo about one-fifth of the grain.

Perennials or annuals of very varied habit, sometimes bulbous at the base or suffrutescent; blades flat or convolute, sometimes much reduced or suppressed; ligules membranous, usually short or reduced to a narrow rim; panicle or racemes sometimes very scanty or even reduced to solitary spikelets.

E. erecta, Lam.

Houtboschberg, *Burt-Davy* 5120; Rietvlei, Belfast, in shaded donga, *Burt-Davy* 1244.

XLV. PHALARIS, Linn. (204).

Spikelets laterally compressed, in contracted, more or less spike-like, panicles; rhachilla disarticulating above the glumes, not or obscurely produced beyond the terminal floret. Florets 3; the lower two minute, rudimentary, the uppermost ♂, enclosed by the

glumes. Glumes subequal, boat-shaped, keeled, keel often winged. Empty valves very small, subulate to lanceolate, membranous, with a callous base, or one or both reduced to a minute callous scale; fertile valve thin, ultimately rigid, 5-nerved, awnless, ovate, acute. Pale almost as long as the valve, 2-nerved (sometimes obscurely). Lodicules 2, hyaline. Stamens 3. Styles long, distinct. Stigmas plumose, exerted from the top of the spikelet. Grain much compressed, ovate, free, enclosed by the valve and pale; hilum oblong, short; embryo equalling one-fourth the length of the grain.

Annuals or perennials; leaves flat; panicle terminal, stiff, spike-like, subcapitate or interrupted and lobed; pedicels very short.

P. minor, Retz.

Kolonies Plaats, Potchefstroom, *Burt-Davy* 9095.

Introduced with agricultural seeds.

P. arundinacea, Linn.

Wemmershoek, Lydenburg District, *Burt-Davy* 7692; Vlakfontein, near Amersfoort, *Burt-Davy* 4063.

Reed canary grass. "Makes good hay if cut before flowering."—*Burt-Davy*.

P. canariensis, L.

Pretoria, *Burt-Davy* 792; Brugspruit, Middelburg, *Cooper* in *Nat. Herb.* H. 4165; Zeerust, Marico, *Frost* in *Nat. Herb.* 6041; Beginsel, Standerton, *Burt-Davy* 3154.

Canary grass; introduced.

P. paradoxa, Linn. (Sp. Pl., ed. II, 1665).

Onderstepoort, Pretoria, *Parkes* in *Nat. Herb.* H. 4998; De Kroon, Brits, Rustenburg District, *Ras* in *Nat. Herb.* H. 5982.

An annual weed—not very common.

XLVI. ANTHOXANTHUM, Linn. (205).

Spikelets oblong to narrow-lanceolate, slightly laterally compressed; rachilla disarticulating above the upper glume, not produced beyond the uppermost floret. Florets 3, heteromorphous, the lower two ♂ or barren, the terminal ♀. Glumes persistent, membranous, 1–3-nerved, keeled, acuminate; upper longer. Lower two valves equal and very similar, oblong, emarginate, membranous, strongly laterally compressed, 5–7-nerved, keeled, hairy, awned, awn of the lower valve short, from a third-half way below the tip, of the upper longer, kneed, from near the base, rarely from the middle; terminal valve much shorter than the lower two, broadly elliptic, very thin, delicately 7–1-nerved. Pales of the lower two florets, if present, 2-keeled, of the terminal 1-nerved. Lodicules 0. Stamens 3 in the ♂, 2 in the ♀ florets. Styles distinct, long. Stigmas long, exerted from the top of the spikelet, plumose. Grain ovoid, slightly laterally compressed; hilum punctiform; embryo quarter the length of the grain.

Perennial or annual; blades flat, usually flaccid; panicle slender, very narrow or spike-like, sometimes reduced to scanty racemes; sweet-scented.

A. odoratum, L.

Lions Glen, Amsterdam, Ermelo District, *Buchanan* in *Nat. Herb.* H. 6152.

"Sweet vernal"; probably an escape from cultivation.

XLVII. ARISTIDA, Linn. (208).

Spikelets 1-flowered, narrow, paniced, rachilla disarticulating above the glume, not produced. Glumes usually persistent, narrow, 1–3-nerved, muticous or mucronate, awnless. Valve convolute, cylindric or oblong-cylindric, 3-nerved, awned rather rigid, tips gradually tapering or minutely bilobed, sometimes jointed at or above the middle; callus villous, shortly bearded, usually pungent; awn nearly always 3-partite from the base or above the

simple base (stipitate), very rarely simple, continuous with the valve or disarticulating from it or deciduous with a portion of the valve, foot straight or twisted bristles plumose or the lateral or all naked. Pale small, oblong, 2-nerved or nerveless. Lodicules 2, finely nerved. Stamens 3. Ovary glabrous. Styles distinct, short. Stigmas plumose, laterally exserted. Grain slender, cylindric or oblong-cylindric, terete, sometimes grooved, tightly embraced by the valve; hilum linear, almost as long as the grain; embryo short or long.

Annual or more often perennial; tufted, usually with more or less wiry culms; blades narrow, usually convolute; ligule usually a line of very short hairs; panicle varying from spike-like to effuse.

A. adscensionis, L.

Bushveld, Pretoria, *Reck* in *Nat. Herb.* H. 185; Potgietersrust, Waterberg, *Burt-Davy* 2246; Wonderboompoort, Pretoria, *Mogg* in *Nat. Herb.* H. 9940; Pyramid Estate, Potgietersrust, Waterberg, *Galpin* 8892.

A. scabrivalvis, Hack.

Skidders Court, Pretoria, *Stent* in *Nat. Herb.* H. 3379; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 427; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4429; Pyramid Estate, Potgietersrust, Waterberg, *Galpin* 8891.

A. junciformis, Trin. & Rupr.

Arcadia, Pretoria, *Burt-Davy* 7494; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 426, 423; Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21135; Billy's Vlei, Ermelo, *Burt-Davy* 9267; Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 5913, H. 994; Uitgevalen, Heidelberg, *Burt-Davy* 9185; Christiana, Bloemhof, *Burt-Davy* 13057, 14146.

Some of the species quoted above have been distributed as *A. angustata*, Stapf, but after careful examination of all our material I can find no constant characters on which to separate the two species *A. junciformis* and *A. angustata*. The slight compression of the culms is more noticable in some specimens than in others, but is not correlated with other distinctive characters. The length of the awns is variable as is also the character of the glumes. I find long mucronate and almost emucronate glumes on the same plant. The inconspicuous nodes of the sheaths, with the swollen and often protruding culm nodes, seems to be a fairly constant character in individual plants and is often correlated with much-branched culms, but not always. I have not seen Stapf's type of *A. angustata*, but *Galpin* 6901, from Maclear, Cape Province, which is identified as that species and which identification (according to the label) has been confirmed by Stapf, has these same characteristic nodes.

A. sciurus, Stapf.

Waterval Boven, Lydenburg, *Burt-Davy* 1418 and 1449; between Machadodorp and Carolina, *Mundy* in *Nat. Herb.* H. 4292A.

A. Burkei, Stapf.

Klerksdorp, Potchefstroom, *Weeber* in *Nat. Herb.* H. 5648; Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21121; Uitgevalen, Heidelberg, *Burt-Davy* 9161 and 9184; Christiana, Bloemhof, *Burt-Davy* 13030.

A. bipartita, Rupr. & Trin.

Waterval Boven, Lydenburg, *Burt-Davy* 1443B; Uitgevalen, Heidelberg, *Burt-Davy* 9125.

A. congesta, Roem. & Schult.

Johannesburg, *Bryant* D. 47; Germiston, near Johannesburg, *Fenn* in *Nat. Herb.* H. 8703; Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 5914; Crocodile River, Rustenburg, *Burt-Davy* 9302; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 421; Springbok Flats, Waterberg, *Burt-Davy* 1112; Zoutpansberg, *Junod* in *Nat. Herb.* H. 4662; Smitskraal, Bloemhof, *Burt-Davy* 10368; Christiana, Bloemhof, *Burt-Davy* 1654.

A. barbicollis, Trin. & Rupr.

Arcadia, Pretoria, *Burt-Davy* 3928; Skinners Court, Pretoria, *Junod* in *Nat Herb.* H. 4519; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 422; downs, Zoutpansberg, open veld, *Junod* 4078; Christiana, Bloemhof, *Burt-Davy* 13031.

A. sieberiana, Trin.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 425; Christiana, Bloemhof, *Burt-Davy* 13030A.

A. vestita, Thunb.

Meintjes Kop, Pretoria, *Burt-Davy* 4514; Billy's Vlei, Ermelo, *Burt-Davy* 9375.

A. spectabilis, Hack.

Kuduspoort, Pretoria District, *Rehmann* 4695*.

There is nothing in the National Herbarium that agrees with Hackel's description of this species, and I have not seen Rehmann's specimen on which he founded it.

A. sp.

Warmbaths, *Burt-Davy* in *Nat. Herb.* 2407.

This resembles *Aristida stipoides*, var. *meridionalis*, Stapf, *ex descr.*, but the spikelets are dark purple brown and the lower sheaths densely coated with fugacious wool.

A. stipoides, Lam.

Pyramid Estate, Potgietersrust, Waterberg, *Galpin* 8890.

A. uniplumis, Licht.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 424; Schweizer Rencke, Bloemhof District, *Burt-Davy* 1627; near Christiana, Bloemhof, *Burt-Davy* 12975, 10123.

XLVIII. **ORYZOPSIS**, Mich. (210).

Spikelets ovoid or lanceolate, usually awned, loosely panicked; rachilla disarticulating above the glumes, not continued beyond the floret. Floret 1, ♂, shorter than the glumes. Glumes equal or subequal, persistent, usually acute or subacuminate, thin, 3-9-nerved, side nerves often very short. Valve convolute, more or less flattened from the back, ovate to obovate or lanceolate, 5- (very rarely 3-) nerved, rather rigidly membranous to coriaceous, tips obscurely 2-lobed; callus very short and broad, obtuse; awn, if present, from the minute sinus of the tip, jointed on the valve and easily deciduous, bristle-like, naked. Pale almost as long as the valve or rather shorter, 2-nerved. Lodicules 2, rarely 3, hyaline. Stamens 3. Ovary glabrous. Styles distinct, short. Stigmas plumose, laterally exerted. Grain tightly enclosed by the hardened valve and pale, oblong, ovoid or ellipsoid, or obovoid, terete; embryo small; hilum filiform, shorter than the grain, often obscure.

Perennial; tufted; blades linear, usually long, flat, flaccid; ligules membranous; panicles effuse, or more or less contracted, always very lax, often nodding.

O. sp.

Malta, in gorge, Zoutpansberg, *Junod* 4432.

XLIX. **PSEUDOBROMUS**, K. Schum. (216).

Spikelets lanceolate, scarcely compressed, pedicelled, panicked; rachilla disarticulating above the glumes, produced beyond the base of the floret into a fine bristle. Floret 1, ♂, exerted from the glumes. Glumes persistent, slightly unequal, lanceolate to oblong, 1-3-nerved. Valve lanceolate in profile, rounded on the back, involute, passing into a straight awn or with the awn subterminal, subherbaceous, 3-nerved; callus very short, obtuse, glabrous. Pale equalling the valve, 2-keeled, entire. Lodicules 2, hyaline, unequally

2-lobed. Stamens 3. Ovary oblong, pubescent at the top. Styles distinct, very short. Stigmas slender, plumose, laterally exerted. Grain unknown.

Perennial; blades flat, broad-linear, flaccid; ligules membranous; panicle large, lax, nodding.

P. africanus, Stapf.

Houtbosch, *Rehmann*, 5732*.

L. SPOROBOLUS, R. Br. (230).

Spikelets usually very small, variously paniced, continuous on the pedicels; rhachilla more or less readily disarticulating above the glumes, not continued, or very rarely produced into a bristle. Floret 1, ♀. Glumes 2, delicately membranous, lower usually smaller, nerveless, upper 1-nerved, falling away one after the other. Valve more or less resembling the upper glume, 1-nerved or more or less distinctly 3-nerved. Pale usually almost as long as the valve, 2-nerved, folded between the nerves, often split by the maturing grain. Lodicules 2, small, broadly cuneate, glabrous, thin. Stamens 3, rarely 2. Ovary glabrous. Styles short, distinct, terminal. Stigmas plumose or subaspergilliform. Grain free, falling out or retained and dehiscing; pericarp thin, usually swelling in water, rigid, dehiscing, or the inner layers mucilaginous when wetted, and adherent, or the whole pericarp adnate and indistinct; hilum small, punctiform, basal; embryo rather large.

Annuals or perennials of various habit; ligules reduced to a ciliate or ciliolate rim; panicles contracted to spike-like, or more or less open, sometimes extremely lax; spikelets mostly $\frac{1}{2}$ –1 lin. long.

S. festivus, var. **stuppens**, Stapf.

Pretoria District, Claremont, *Kresfelder* in *Nat. Herb.* H. 20834; Wonderboom, *Burt-Davy* 3343, 7148, 2683; Irene, *Burt-Davy* 14656; *Pole Evans* £20 and in *Nat. Herb.* H. 21223; Johannesburg, *Burt-Davy* 7296; Marikana, Rustenburg District, *Burt-Davy* 161; near Nylstroom, Waterberg District, *Burt-Davy* 2023, 2068.

S. festivus, Hochst., var. **fibrosus**, Stapf [*Chevalier* in Sudania, 142, 155 (1911)].

Springbok Flats, Waterberg District, *Burt-Davy* 1111.

S. discosporus, Nees.

Pretoria District, Kaalfontein, *Mogg* 3740; Irene, *Burt-Davy* 14657; Bloemhof District, near Christiana, *Burt-Davy* 11409, 12778, 11389, 11233.

S. Ludwigii, Hochst.

Christiana, Bloemhof District, *Burt-Davy* 12809, 12950, 12498.

S. centrifugus, Nees.

Spionkop, Ermelo, *Burt-Davy* in *Nat. Herb.* H. 9236; between Machadodorp and Carolina, *Mundy* in *Nat. Herb.* H. 4286.

S. fimbriatus, Nees.

Near Machavie Station, Potchefstroom, *Burt-Davy* 1476; Schweizer Reneke, *Burt-Davy* 1684, 1636, 1691; Fourteen Streams, Bloemhof, *Burt-Davy* 1580; Kaffraria, Christiana, *Burt-Davy* 14160, 12765, 13054, 12758, 13090.

Var. **latifolia**, Stent, var. *nov.*, a more robust form, with usually broader and longer panicles and flat glaucous leaves up to 12 mm. broad = *Sporobolus Rehmanni*, Hack. (in part).

Boschveld, at Klippan, *Rehmann* 5373 (in *Herb. Alb. Mus.*); Mosdene, Naboomspruit, Waterberg, *Galpin* M. 460, 461; Hamanskraal, Pretoria District, *Burt-Davy* 1086.

The type occurs freely in the dry and sandy regions of the grass and thorn veld of the southern and south-western portions of the Transvaal. It is plentiful in association with *Eragrostis* spp. in parts of Bechuanaland and on the high veld.



Plate VII.

Sporobolus Sumatensis, Stent.

A, Lower glume; B, Upper glume; C, Valve; D, Pale (flattened out);
E, Pistil and stamens; F, Ligule.

The variety is a bushveld form that also occurs in Natal and probably extends north into the tropical regions.

Rehmann's 5373, on which Hackel founded this species (*S. Rehmannii*), has hairy leaves and a narrow panicle. I have seen no other specimen like it. Stapf (in Dyer, *Flor. Cap.*, VII, 585) has kept *S. Rehmannii* as a distinct species and has amplified Hackel's description to include all the glabrous broad-leaved forms of the bushveld and Natal. Critical examination of a number of specimens ranging from *S. fimbriatus sensu stricto* through all variations of width of leaf and panicle to the broad-leaved and robust form of the bushveld, has convinced me that this latter is only a climatic and edaphic variety of *S. fimbriatus*.

A very early grass, useful for spring grazing.

S. indicus, R. Br., var. *laxus*, Stapf.

Johannesburg, common along roadsides, *Bryant* B. 23; Carolina, *Burt-Davy* 2955; Sterk Hill, Lydenburg, *Burt-Davy* 478.

The type apparently does not occur within the limits of the Transvaal.

S. pyramidalis, Kunth. (*Flore d'Oware*, ii, p. 36).

Springbok Flats, Waterberg, *Burt-Davy* 1137; Mosdene, Naboomspruit, *Galpin* M. 458; Lydenburg, *Burt-Davy* 414.

S. robustus, Kunth. (*Rev. Gram.*, ii, p. 425).

Messina, Zoutpansberg, *Pole Evans* in *Nat. Herb.* H. 13113.

S. pectinatus, Hack. (*Oestern. Bot. Zeitschr.*, L. iii, 198).

Derdepoort, Pretoria, *Fouche* 302; near Observatory, Johannesburg, *Burt-Davy* 3097; Carolina, *Burt-Davy* 2990 and 7736; Zoutpansberg, downs, open veld, *Junod* 4091.

S. Smutsii, Stent, n. sp.

Gramen perenne; culmi decumbentes ascendentes, glabri, pluribus nodis saepe ramati; radicibus e nodis brevibus inferioribus. Innovationes steriles, saepe longae, plures atque breviter nodatae, foliatae; foliorum vaginae firmatae, carinatae, arctae, glabrae, ore barbato marginibusque ciliatis exceptis, vel setis e tuberculis inspersae, tenuiter striatae; margines crassi, cartilaginosi, pectinatique.

Panicula erecta, ovata, lanceolata, acuta (raro constricta spiciformaque), 2-4 cm. lata, 8-10 cm. longa. Rami verticillati vel superiores $\pm$ inspersi; infimi ad 3.5 cm. longi, nudi a base .5-1.25 cm., plerumque ascendentes; superiores patentem ascendentes. Ramuli breves, ad 3 mm. longi, $\pm$ secundi, appressi. Pedicelli brevissimi. *Spiculae* 1.7-2 mm. longae, ablongae, obtusae; *gluma inferior* parva .6 mm. longa, enervata, obtusa; *superior* spiculam aequans, 1-nervata, acuta vel sub-acuta, navicularis carina obtusa; *valvae* glumae superioris similis, sub-acuta, 1-nervata; *palea* extensa truncata; *antherae* 3, 1.25 mm. longae; *lodicalae* truncatae, minutae.

Rooikop, Bushveld, Pretoria District, *Pole Evans* 590, 271, 536, 556; Leeuwkraal, *Pole Evans* 668.

A very characteristic grass, with its decumbent ascending habit and short, broad, flat leaves on leafy runners. Dr. Pole Evans reports it as being very plentiful on the adjoining farms Rooikop and Leeuwkraal, but it has, so far, not been reported from any other locality.

It has little agricultural value.

S. nitens, Stent, n. sp.

Gramen perenne, rhizomata obliqua. Culmi erecti vel prostrati ascendentes saepe radicibus ramisque e nodis inferioribus, coarcti, glabri, nitentes, 1-2 nodis, 13-30 cm. longi. Foliorum vaginae arctae, strictae, laeves, glabrae, praeter margines saepe ciliatos, vaginae basis breviores compressae, pallidae, firmissimae, nitentes, persistentes; ligula corona densa pilorum formata; laminae ad 6 cm. longae, 7 mm. latae, basis rotundatae et plus minus amplectentes, apicem obtusum vel subacutum versus attenuatae, planae, firmissimae, tenuiterque striatae; glabrae, vel paucis setis e tuberculis ortis inspersae, marginibus cartilaginosis minute pectinata ciliatis. *Panicula* ovata ad lanceolata, 6-10 cm. longa,



Plate VIII.

Sporobolus nitens, Stent.

A, Lower glume ; B, Upper glume ; C, Valve ; D, Pale ; E, Pistil
and stamens ; F, Caryops.

2.5-4 cm. lata. *Rami* infimis verticillati ascendentesque, raro pauci vel omnes patentes, ceteri solitarii, vel binati, racho spirale depositi, per $\frac{1}{2}$ - $\frac{3}{4}$ longitudinis indivisi, rami supremi brevissimi, e basi divisi, excepti; *ramuli* brevissimi, plus minus secundi, appressi 1-5 spiculati; *pedicelli* brevissimi. *Spiculae* 1-5 mm. longae; *gluma inferior* ablorga, lanceolata, acuta, vel sub-acuta, enervis, 1 mm. longa; *gluma superior* 1.5 mm. longa, ovato-acuminata, valide uninerva, carina minute pectinato, ciliata; *valva* glumam superiorem equale, vel pauce longior; valide uninerva, acute acuminata, laeva; *palea* valvæ equale, lata, apicem bilobum versus attenuata, mix ad basim findata; *antherae* 3, rubescentes, circa .4 mm. longae; *lodiculæ* ovatae, parvissimae; *caryopsis* ovata, vel rotundata; pericarpium in aquam tumentum.

In the plain at the foot of the Magaliesberg Mountains, *Schlecht* 3679; Rooikop Bushveld, Pretoria District, *Pole Evans* 667; Gezina, Pretoria, *Fouche* and *Stent* in *Nat. Herb. H.* 21445; between Warmbaths and Nylstroom, *Pole Evans* 691.

Near to *S. Smutsii*, from which it differs in the hard, shining, lower sheaths, the spiral arrangement of the branches of the panicle, the densely contracted and spike-like tips of the panicle and of the branches—and the distinctly and acutely bilobed tip to the pale. Useful as a crop for sheep, but otherwise of little agricultural value.

LII. AGROSTIS, Linn. (242).

Spikelets from less than 1 lin. to $2\frac{1}{2}$ lin. long, panicked; rhachilla not continued beyond the floret or produced into a minute point or short delicate bristle, glabrous or shortly hairy. Floret 1, ♂, shorter than the glumes. Glumes equal or sub-equal, usually lanceolate and acute, rarely oblong and obtuse, awnless, membranous, usually 1-nerved, keeled. Valve broadly oblong, delicately membranous, glabrous or hairy, usually truncate, 5- (rarely 3-) nerved, awned from the back or awnless; side-nerves evanescent below, often excurrent into fine mucros or bristles; callus very small, glabrous or minutely hairy or rarely bearded. Pale delicate, hyaline, usually shorter than the valve or very short or obsolete, 2-nerved or nerveless. Lodicules 2, lanceolate, delicately hyaline. Stamens 3. Ovary glabrous. Styles distinct, very short. Stigmas plumose, laterally exserted. Grain free, enclosed in the scarcely altered floret, oblong, more or less dorsally compressed, grooved in front, rarely subterete; embryo short; hilum punctiform, basal.

Annuals and perennials, of varying habit; blades usually flat, often flaccid; ligules membranous; panicle usually much divided, often delicate, more or less effuse or contracted, rarely spike-like; branches and branchlets fine to very fine; spikelets usually very numerous, much gaping, at least, temporarily.

A. natalensis, Stapf.

Spionkop, Ermelo District, *Burt-Davy* 9232; Belfast, Lydenburg, *Burt-Davy* 1328, 1277.

A. laenantha, Nees.

Hartebeestnek, Pretoria, *Burt-Davy* 770; near Nylstroom, Waterberg, *Burt-Davy* 4165; Lake Chrissie, Ermelo, *Mundy* in *Nat. Herb. H.* 4283; Lydenburg, along water-courses, *Burt-Davy* 415; Wonderfontein, Marico, *Burt-Davy* 7591; Potchefstroom, *Burt-Davy* 1765.

A. suavis, Stapf (Kew Bull., 1908, 227).

Spionkop, Ermelo, *Burt-Davy* 9235.

LIII. CALAMAGROSTIS, Roth. (247).

Spikelets very narrow, lanceolate, acuminate, in contracted much-divided panicles; rhachilla disarticulating above the glumes, not, or very shortly, continued beyond the floret. Floret 1, ♂, much shorter than the glumes. Glumes equal or subequal, very narrow, linear-lanceolate, acuminate, membranous keeled. lower 1-nerved, upper 3-nerved. Valve narrow, lanceolate in profile, membranous, glabrous, more or less shortly bifid,

5-3-nerved with a fine short dorsal, rarely subterminal, awn; callus small, long hairy, hairs usually much exceeding the valve. Pale 2-nerved, as long as the valve or somewhat shorter. Lodicules 2, hyaline. Anthers 3. Ovary glabrous. Styles distinct, short. Stigmas plumose, laterally exserted. Grain enclosed by the hardly changed valve and pale, free, subterete; hilum basal, small; embryo small.

Perennial, usually rather robust; blades long, linear, flat; ligules scarious; panicle more or less contracted, narrow rather dense, with much-divided branches, and short branchlets and pedicels; florets surrounded by long fine hairs.

C. epigeios, Rott., var. **capensis**, Stapf.

Lichtenburg, *Burt-Davy* 62A; Doornkloof, Irene, *Pole Evans* 336.

LIII. **HOLCUS**, Linn. (257).

Spikelets in rather dense, oblong, or interrupted panicles, laterally compressed, disarticulating from the tips of the pedicels; rhachilla slightly produced beyond the upper floret, disarticulating more or less readily below the valves; joints slender, lower curved and often appendaged. Florets 2; lower ♂, upper usually ♂, sometimes ♀ or barren. Glumes 2, membranous, keeled, acute or acuminate, lower 1-nerved, upper 3-nerved, sometimes awned. Valves shorter than the glumes, characteous, very obscurely 5-3-nerved, lower awnless, upper awned. Pales narrow, 2-keeled. Lodicules 2, delicate. Stamens 3. Ovary glabrous. Styles distinct. Stigmas plumose, laterally exserted. Grain laterally compressed, enclosed by the valve and pale and often adhering to the latter, soft; hilum short; embryo small.

Annual or perennial; blades flat or convolute when dry; panicle usually more or less contracted, sometimes almost spike-like; spikelets deciduous, pallid.

H. lanatus, L.

Pretoria, *Macdonald* in *Nat. Herb. H.* 5471; Broederstroom Orchards, Haenertsberg, *Zoutpansberg*, *Reineke* in *Nat. Herb. H.* 9763.

Yorkshire Fog—probably an escape from cultivation.

LIV. **AVENAstrum**, Jess. (275).

Spikelets usually erect or sub-erect, rarely nodding, medium-sized (4-7 lin. long, rarely less or more), in nearly always erect, often stiff panicles; rhachilla more or less long-hairy, disarticulating above the glumes and between the valves, produced into a short bristle beyond the uppermost floret or ending with a rudimentary valve. Florets 3-6, ♀, or the uppermost more or less reduced. Glumes 2, persistent, scarious, more or less unequal, acute or acuminate, more or less distinctly keeled, lower 1-nerved or 3-nerved, upper 3- (very rarely 5- to sub-7-) nerved. Valves usually distinctly exserted from the glumes, more or less herbaceous, with scarious or hyaline tips, often rather firm, acute, or acuminate, bifid, with or without bristles from the lobes, 5-9-nerved, awned; awn dorsal from the middle or slightly above it, kneed and twisted below; callus short or elongate, villous. Pales shorter than the valves, 2-keeled, 2-toothed, or bifid. Lodicules 2, rather large, hyaline. Stamens 3. Ovary hairy from the middle upwards or at the top only. Styles distinct, short. Stigmas laterally exserted, plumose. Grain oblong, slightly laterally compressed, usually grooved in front, hairy at the top, pallid, rather soft, embraced by the somewhat hardened valve and the pale; hilum linear, up to half the length of the grain; embryo small.

Perennial, caespitose; blades linear, usually narrow, flat, or convolute; often setaceous ligules hyaline; panicle narrow, erect, often stiff, rarely flaccid or expanded.

A. turgidulum, Stapf.

Pretoria, *Stent* in *Nat. Herb. H.* 4363; Potchefstroom, *Burt-Davy* 5591; Vlakfontein, Wakkerstroom, *Burt-Davy* 4154.

LV. TRISTACHYA, Nees. (277).

Spikelets 5 lin. to 2 lin. long, lanceolate to linear, sessile or shortly pedicelled in clusters of three at the tips of the branches of a raceme or panicle; rachilla disarticulating between the valves and less readily or imperfectly below them, glabrous, not produced beyond the upper floret. Florets 2, heteromorphous; lower ♂, upper ♀. Glumes persistent, membranous or sub-coriaceous, more or less unequal, 3-nerved, glabrous or bristly from black glands. Lower valve membranous, 3-nerved or 5-9-nerved, with some of the nerves much shorter than the others; upper valve convolute, membranous, or ultimately cartilaginous, bifid, 7- (rarely 5-) nerved, awned from between the lobes; awn kneed, twisted below. Pales membranous, hyaline, 2-keeled. Lodicules 2, cuneate, very fleshy. Stamens 3. Ovary obovoid to oblong, glabrous or the top hairy, rudimentary in the ♂ floret. Stigmas plumose, long, laterally exerted. Grain obovoid to linear-oblong, grooved, tightly embraced by the valve and pale, free; hilum linear, long; embryo large; albumen very hard.

Perennial; generally coarse grasses, growing mostly in compact tufts; leaves rigid; ligule a line of hairs; triplets of spikelets large and few in racemes, or smaller and more numerous in scantily branched panicles, sometimes deciduous from the jointed peduncles.

T. Rehmanni, Nees.

Eloff's Plantation, Pretoria, *Macdonald* in *Nat. Herb.* H. 5442; near Observatory, Johannesburg, *Burt-Davy* 4000 and 7491; Billy's Vlei, Ermelo, *Burt-Davy* 9222; Belfast, Lydenburg, *Burt-Davy* 1341; Bloemhof, *Theunissen* in *Nat. Herb.* H. 7871.

T. pallida, Stent, n. sp.

Gramen perenne. Culmi erecti ad 64 cm. alti, fasciculati e rhizomata abliquum, graciles, 2-nodi, glabri vel parte inferiore tenuiter hirsuta, nodis pubescentibus et saepe barbatis, innovationes intravaginales.

Folia plerumque a basem nata, culmos fere basim inflorescentiae condentia; *raginae* glabrae, vel sparse hirsutae, base dense tomentosae, superiores; arctae angustaeque, inferiores latiores tandem solutae et patentes, tenuiter striatae, laevae, plerumque oribus longe hirsutae. *Laminae* angustae, arcte involutae, et saepe convolutae, glabrae vel albido-hirsutae, scaberrulae, pallido glaucae, illae innovationum ad 34 cm. longae, 1.5 n. m. latae; apice filiformo. *Panicula* contracta, ad 8 cm. longa, nodo infimo villosa; rami curvati, ascendentes, simplices, infimi verticillati, sequentes solitarii, vel raro birati; triade spicularum terminati. *Gluma inferior* lanceolata acuminata rigida, mucronata, 21 mm. longa, firmiter 3-nerva, nervi exteriores marginesque ordines duas setas e tuberculis nigris ferentes. *Gluma superior* glumam inferioram aequalis, sed tenuior, mutica, glabra firmiter 3-nervata, nervis scaberrulis. *Anthoecium inferum* ♂; *valva* 15 mm. longa, lanceolata, acuminata, glabra, 7-nervis nervi scabri, interiores brevissimi, exteriores contigui; *palea* 8 mm. longa, biloba, lobi obtusis, minute ciliatis; carinis validis, minute ciliatis; *antherae* ad 4.5 mm. longae; *lodicae* cuneatae, 5 mm. longae, carnosae. *Anthoecium superum* ♀; *valva* ovata lanceolata, biloba, 5-nerva, 7 mm. longa apice cillo ad basim aristatae, pubescente; lobi 2-3-nervis, 2 mm. longi, setis 5 mm. longis, arista 3.5 cm. longa, geniculata; callum acutum longe barbatur; *palea* angusta, obtusorotundata, vel subtruncata, carinae anguste alatae; *caryopsis* oblonga, apice hirsuta; *antherae* lodiculaeque eas in anthoecia inferes similis.

Pietersburg, *Capt. Goument* in *Nat. Herb.* H. 21318; Pyramid Estate, near Potgietersrust, upper slopes of granite mountains, *Galpin* 8893; Pretoria, *Schlechter* 3587.

Near to *T. biseriata*, Stapf (*ex descr.*), from which it differs in the 2-noded culms, the glabrous upper glume and lower valve, the 2-nerved lobes to the upper valve and the lack of a callous line at base of awn, and other minor points. I have not seen *Bicharan* 220, on which Stapf founded his *T. biseriata*, nor any other specimen that would correspond to his description of that species.




Plate IX.

Tristachya pallida, Stent.

A, Lower glume; B, Upper glume; C, Lower valve; D, Pale; E, Upper floret; F, Upper valve; G, Pale; H, Pistil, stamens, and lodicules.

T. leucothrix, Trin.

Meintjes Kop, Pretoria, *Burt-Davy* 3547; Barberton Mountains, *Burt-Davy* 338; Leeuwpoot, Carolina, *Burt-Davy* 7448; Belfast, Lydenburg, *Burt-Davy* 1324; Heidelberg, *Burt-Davy* 3146; Blesbokspruit, Ermelo, *Burt-Davy* 949; Vlaktefontein, Wakkerstroom, *Burt-Davy* 9052; Athol, Wakkerstroom, *Macdonald* in *Nat. Herb.* H. 104.

T. pedicellata, Stent (in *Bothalia*, I, iii).

Mosdene, Naboomspruit, *Galpin* M. 428; Warmbaths, *Baylis* in *Nat. Herb.* 4350; *Burt-Davy* 1154.

LVI. **TRICHOPTERYX**, Nees. (278).

Spikelets usually 2-18 lin. long, more or less lanceolate to linear, pedicelled, paniced; rachilla disarticulating between (and less readily below) the valves, glabrous, not produced beyond the upper floret. Florets 2, heteromorphic; lower ♂, rarely barren; upper ♀. Glumes persistent, membranous or subcoriaceous, unequal, 3- (very rarely 4-6-) nerved, glabrous or bristly from black glands. Lower valve membranous 3- (very rarely 7-) nerved, with the innermost side nerves much shorter; upper valve terete, membranous to cartilaginous, shortly bifid with the lobes sometimes produced into bristles, 5-9-nerved; awn from between the lobes, kneed, twisted below. Pales membranous, 2-keeled, narrower in the ♀ floret and channelled between the stout keels. Lodicules 2, cuncate, usually very fleshy. Stamens 2, rarely 3, or 0 in the lower floret. Ovary oblong, glabrous, rudimentary in the ♂ floret. Styles distinct, glabrous. Stigmas plumose, long, laterally exserted. Grain obovoid to linear-oblong, grooved or almost terete, tightly embraced by its valve and pale, free; hilum linear, long; embryo large; albumen very hard.

Perennial, rarely annual grasses of very different habit; leaves more or less rigid; ligules a line of hairs; spikelets slender, from 2 lin. to 1½ lin. long, scattered in mostly contracted, narrow panicles.

T. simplex, Hack.

The Willows, Pretoria, *Burt-Davy* 7486; Johannesburg, *Burt-Davy* 7492, *Bryant* C. 38; Waterval Boven, *Burt-Davy* 1426.

Var. **crinita**, Stapf.

Ermelo, *Burt-Davy* 5503; Spionkop, *Burt-Davy* 9230; Belfast, Lydenburg, *Burt-Davy* 1342.

Var. **sericea**, Stapf.

Waterval Boven, Carolina, *Burt-Davy* 1439, *Burt-Davy* 2983; Machadadorp, *Mundy* in *Nat. Herb.* H. 4286; Billy's Vlei, Ermelo, *Burt-Davy* 9201.

T. flavida, Stapf.

Pretoria, *Macdonald* in *Nat. Herb.* H. 5429, 5407, 5413, 54531; *Schlechter* 3606; Johannesburg, Observatory Ridge, *Burt-Davy* 3093.

T. sp. A.

Waterval Boven, Carolina, *Burt-Davy* 1426, 1408; downs, Zoutpansberg, *Junod* 4082.

A tropical species with quite glabrous spikelets, broadly truncate lower glume—a rather short, delicate, and obscurely nerved fertile valve.

“Very much spread over the veld; bad food for cattle.”—*Junod*.

LVII. **MICROCHLOA**, R. Br. (281).

Spikelets 1-2-flowered, small, sessile, crowded, unilateral on a flattened rachis, alternately 2-seriate from near the margins of the rachis or in a single row; rachilla disarticulating above the glumes, more or less produced. Floret hermaphrodite, or if 2, the lower hermaphrodite, the upper male or indicated by an empty valve. Glumes 2, persistent or (particularly the upper) deciduous, strongly 1-nerved, flattened from the

back or keeled, subequal. Valve shorter than the glumes, delicate, white, minutely or obscurely mucronulate or emarginate, 3-nerved, densely hairy along the nerves (if 2, the upper glabrous); callus small, acute, hairy. Pale slightly shorter than the valve or almost equal, 2-keeled. Lodicules 2, cuneate, glabrous, thin, faintly nerved. Stamens 3. Ovary glabrous (quite suppressed in the upper floret). Styles distinct. Stigmas plumose, laterally exerted. Grain oblong, terete, triquetrous, or compressed, embraced by the unchanged valve and pale, free; hilum punctiform; embryo equalling quarter to half the length of the grain.

Perennial, rarely annual, sometimes densely tufted; leaves narrow, often sub-setaceous; ligules reduced to a minutely ciliolate rim; spikes solitary, terminal (in the African species) or 2-4 in a terminal umbel, straight or curved.

M. setacea, R. Br.

Pretoria, *Macdonald* in *Nat. Herb.* H. 5476; Houtbosch, *Schlechter* 4697; Sabie Falls, Lydenburg, *Burt-Davy* 1550A; Belfast, Lydenburg, *Burt-Davy* 1287A; Potchefstroom, *Burt-Davy* 1225; Christiana, Bloemhof District, *Burt-Davy* 11392, 14099, 14133, 12781.

M. caffra, Nees.

The Willows, Pretoria, *Burt-Davy* 4483; Irene, near Pretoria, *Burt-Davy* 3859; near Robinsons, Carolina District, *Burt-Davy* 2720; Leeuwpoot, Carolina, *Burt-Davy* 7377; Athol, Ermelo, *Macdonald* in *Nat. Herb.* H. 94.

This species differs from the preceeding (*ex descr.* by Stapf in Dyer, *Flor. Cap.*) in the size of the spikelets and in the shape of the valve, and on these characters I have temporarily based my identifications. I have not seen R. Brown's original description of *M. setacea*, nor have I seen the type-material of either species. The differences between the two grasses are so slight that it may seem advisable, when I have had an opportunity of examining the types, to place them all under one species.

M. altera, Stapf, var. **Ne'sonii**, Stapf.

Spionkop, Ermelo District, *Burt-Davy* 9243.

LVIII. **CYNODON**, Pers. (282).

Spikelets 1-flowered, small, laterally compressed, sessile imbricate, alternately 2-seriate and unilateral on a slender keeled rhachis; rhachilla disarticulating above the glumes, produced, or not, beyond the valve. Floret hermaphrodite. Glumes narrow, keeled, acute or subulate-mucronate, the upper usually deciduous with the valve, the lower subsistent. Valve exceeding the glumes, navicular, firmly membranous, 3-nerved, awnless, keel ciliate. Pale somewhat shorter than the valve, 2-keeled. Lodicules 2, minute, obovate-cuneate, glabrous. Stamens 3. Ovary glabrous. Styles distinct, slightly shorter than the plumose styles. Grain oblong, subterete; embryo about third the length of the grain; hilum linear, two-thirds the length of the grain.

Perennial; stems creeping, rooting at the nodes and emitting from them fascicles of barren shoots and flowering culms; spikes 2-6 in terminal umbels.

C. dactylon, Pers.

Pretoria, *Mogg* in *Nat. Herb.* H. 16950; Germiston, Johannesburg, *Fenn* in *Nat. Herb.* H. 8746; Komatipoort, Barberton, *Burt-Davy* 368; Nylstroom, Waterberg, *Burt-Davy* 2086; bushveld, Pretoria, *S. Reck* in *Nat. Herb.* H. 3533; downs, open veld, Zoutpansberg, *Junod* 4101; Pilgrims Rest, Lydenburg, *Legat* in *Nat. Herb.* H. 1585; Lichtenburg, *Burt-Davy* 58; Schweizer Reneke, Bloemhof, *Burt-Davy* 1634; Christiana, Bloemhof, *Burt-Davy* 14094, etc.

"Bermuda quick." "Regte kweek," a cosmopolitan weed.

C. incompletus, Nees.

Helvetia, near Machadodorp, Lydenburg, *Burt-Davy* 7245; New Agatha, Zoutpansberg, *Burt-Davy* 1214; Spionkop, Ermelo, *Burt-Davy* 9262; near Morgenzon, Ermelo, *Burt-Davy* 1011; Carolina, *Burt-Davy* 2713; banks of the Vaal River, Bloemhof, *Burt-Davy* 1516; Vierfontein, Schweizer Reneke, Bloemhof, *Burt-Davy* 1635.

Indigenous to South Africa. An excellent lawn grass.

C. transvaalensis, *Burt-Davy* (in "Kew Bulletin," 1921, 283)

Burger's Park, Pretoria, *Stent* in *Nat. Herb.* 10798, 10799; Houghton Estate, Johannesburg, *Bryant* C. 10; Uitval, Vereeniging, *Turner* in *Nat. Herb.* 21224; Lake Chrissie, Ermelo, *Burt-Davy* 9270.

There is no doubt that all the above specimens belong to the grass described by Mr. Burt-Davy as *Cynodon transvaalensis*, but at least two of the characters on which he relies to separate it from *C. dactylon* are not constant. The distachyous inflorescence, while being certainly the most common form of inflorescence, is often replaced by a tri- and occasionally quadri-stachyous inflorescence, and the valves vary from ciliate to pronouncedly, though usually appressedly, ciliate. The delicate fine habit of the grass, with its slender stolons and culms and soft, narrow and rather short leaves, renders it quite distinct from even fine-leaved forms of *C. dactylon*. It is one of the best grasses of the country for bowling greens, etc., making a smooth, compact, and beautifully green turf. To horticulturists it is usually known as Germiston or Florida grass.

LIX. HARPECHLOA, Kunth. (285).

Spikelets of 3-4 florets, sessile, crowded, unilateral, alternately biseriate along the midrib of a flattened rhachis; rhachilla disarticulating above the glumes, continuous between the valves, the lower floret hermaphrodite, the following 1 or 2 male, the uppermost barren, rudimentary. Glumes unequal, the lower persistent, keeled, very thin, 1-nerved, the upper much longer, oblong, flattened on the back, 2-3-nerved, firm. Hermaphrodite floret about equalling the upper glume. Valve folded, obliquely oblong in profile, obtuse, white, thin, 3-nerved, densely ciliate along the nerves; callus obscure. Pale slightly shorter, 2-keeled. Lodicules cuneate, fleshy, almost 2-winged. Stamens 3. Ovary glabrous. Styles distinct. Stigmas slender, plumose, laterally exserted. Upper florets crowded in a club-shaped body, not exceeding the hermaphrodite floret, enveloped by the valve of the lower male floret; valves 2-nerved or with a trace of the middle nerve near the apex, ciliolate or glabrous; pales 2-nerved; stamens 3 or 0; ovary usually quite suppressed. Grain free, embraced by the unchanged valve and pale, oblong, obtusely triquetrous; embryo half the length of the grain; hilum punctiform, basal.

Perennial, densely caespitose; leaves firm, folded or convolute above, more or less curved; spikes terminal, solitary, rarely geminate, dark olive-grey.

H. capensis, Kunth.

Houghton Estate, Johannesburg, *Bryant* C. 12; Barberton Mountains, *Burt-Davy* 339; Zoutpansberg, *Worsdall* in *Nat. Herb.* H. 5919; Volksrust, Wakkerstroom, *Burt-Davy* 5350; Ermelo, *Burt-Davy* 3909, 948; Amersfoort, Wakkerstroom, *Burt-Davy* 9051.

LX. CTENIUM, Panz. (286).

Spikelets of 3-4 florets, sessile, compactly crowded, unilateral, alternately biseriate along the midrib of the flattened rhachis; rhachilla disarticulating above the glumes, continuous between the valves, the lower two florets barren or the second male, the third hermaphrodite, the fourth male or barren, or quite rudimentary. Glumes unequal, the lower persistent, keeled, thin, 1-nerved, the upper much longer, oblong to lanceolate, flattened or rounded on the back, firm, 2-3-nerved, with a stiff awn from the middle. Valves oblong in profile, obtuse, 3-nerved, awned just below the tips, ciliate along the

nerves or the uppermost glabrous, white, thin. Pales slightly shorter, 2-keeled or 2-nerved. Lodicules 2, quadrate-cuneate, delicate, faintly nerved. Stamens, three in the hermaphrodite, two in the male florets. Ovary glabrous. Styles distinct. Stigmas slender, long, laterally exerted. Grain free, embraced by the unchanged valve and pale, oblong; embryo up to half the length of the grain; hilum basal, punctiform.

Perennial, densely tufted, rarely annual; leaves narrow, flat, or convolute; spikes terminal, solitary or in umbels of 2-3, usually curved; spikelets prettily pectinate and awned.

C. concinnum, Nees.

Pretoria, *Macdonald* in *Nat. Herb.* H. 5480; Billy's Vlei, Ermelo, *Burt-Davy* 9218, 9226; Ermelo, *Burt-Davy* 9369, 613; Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 4332; Leeuwpoot, Carolina, *Burt-Davy* 7331, 2959; Pan-plaats, Pan Station, Middelburg, *Burt-Davy* 13320, 13289; Belfast, Lydenburg, *Burt-Davy* 1340.

LXI. **CHLORIS**, Swartz. (288).

Spikelets of 2-4 florets, sessile, crowded, unilateral, 2-seriate on a slender rachis; rachilla disarticulating above the glumes, tough between the valves, more or less produced; lowest floret hermaphrodite, the second male or barren, the following, if present, barren, often minute. Glumes 2, persistent, narrow, keeled, acute and mucronate, very thin, or broad, and the upper obtuse, more or less bilobed and rounded on the back. Hermaphrodite floret; valve narrow or broad, 3-nerved, acute or obtuse, minutely 2-toothed, usually awned from below the apex, often ciliate; pale almost equalling the valve, 2-keeled; lodicules 2, minute, delicate, glabrous; stamens 3; ovary glabrous, styles distinct, short; stigmas laterally exerted. Male floret: valve and pale as in the hermaphrodite flower, but smaller and glabrous; rudimentary florets glabrous, awned or awnless, small to very small, usually without a trace of a pale; grain oblong, triquetrous; embryo rather large; hilum punctiform, basal.

Perennial or annual; leaves flat or folded; spikes solitary or several to many in terminal umbels or short racemes; erect or stellately spreading.

C. pycnothrix, Trin.

Arcadia, Pretoria, *Burt-Davy* 8054; *Mogg* in *Nat. Herb.* H. 10086.

C. virgata, Schw.

Irene, near Pretoria, *Burt-Davy* 581; Mosdene, Naboomspruit, Waterberg, in black turf, *Galpin* M. 430; Springbox Flats, Waterberg, *Saunders* in *Nat. Herb.* H. 2138; Shilovane, Zoutpansberg, *Junod* in *Nat. Herb.* H. 4681; Standerton, *Burt-Davy* 907; Schweizer Reneke, Bloemhof, *Burt-Davy* 1650; Christiana, Bloemhof, *Burt-Davy* 12797.

"Old lands grass." "Sweet grass," a useful hay-grass, known in the Orange Free State as Kwas grass.

C. gayana, Kunth.

Skinners Court, Pretoria, *Stent* in *Nat. Herb.* H. 5084; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4454, 4451; Experiment Station, Potchefstroom, *Burt-Davy* 2704.

"Rhodes grass." One of our best indigenous hay and pasture grasses, both frost and drought resistant—very palatable.

C. petraea, Thunb.

Meintjes Kop, Pretoria, *Burt-Davy* 5381; Skinners Court, Pretoria, *Baylis* in *Nat. Herb.* H. 5376; De Kroon, Brits, Rustenburg, *Stent* in *Nat. Herb.* H. 6115; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 431; Warmbaths, Waterberg, *Burt-Davy* 2345; downs, Zoutpansberg, *Junod* 4085; Potchefstroom, *Burt-Davy* 1067; Smitskraal, Bloemhof, *Burt-Davy* 10098.

LXII. **TRIOGON**, Rott. (298).

Spikelets many flowered; flowers, perfect or the uppermost reduced, biseriate and unilateral on a slender terminal spike, disarticulating above the glumes. Glumes 1-nerved dissimilar, lower usually inequilateral, one margin being deeply notched or dilated suddenly below the middle into a lobe or wiry or segment; upper entire or notched below the mucronate or apiculate tip. Valves ovate dorsally convex, 2-fid, and awned in the cleft, or 4-fid, with the outer lobes awned and the inner membranous and rarely awned. Palea broad or narrow, complicate. Stamens 3. Styles very short. Grain very narrow, free.

Slender, densely tufted grasses; leaves usually convolute; ligule delicately hyaline and fimbriate.

T. abyssinicus, Nees (*ex* Steud. Syn. Gram., 301).

Onderstepoort, *Mogg* in *Nat. Herb.* H. 18252; Komatipoort, Barberton, *Schlechter* 11841.

LXIII. **DINEBRA**, Jacq. (302).

Spikelets 2-3-flowered, small, crowded, biseriate unilateral on slender, spreading, or deflexed spikes that are collected in narrow, pyramidal racemes, disarticulating above the glumes. Glumes slightly unequal, persistent, lanceolate, 1-nerved, keeled and awned, much exceeding the valves. Valves very small, broadly ovate, subacute, hyaline 1-nerved. Palea hyaline. Anthers 3, small. Styles free. Grain ovoid, trigonous, wrinkled, free.

An annual, leafy grass; leaves flat. Ligules membranous.

D. arabica, Jacq. (Fragment, 77, t. 121).

Pienaars River, Pretoria District, in swamplands, *Osborne* in *Nat. Herb.*; Onderstepoort, *Pole Evans* 437; Crocodile River, Rustenburg, *Burt-Davy* 9293; Mosdene Naboomspruit, Waterberg, *Galpin* M. 526.

LXIV. **ELEUSINE**, Gaertn. (304).

Spikelets 3-6-flowered, laterally compressed, densely imbricate, alternately biseriate unilateral, sessile on a flattened rachis, the uppermost terminal, perfect rhachilla disarticulating above the glumes and between the valves, or tough, produced, sometimes terminating with a rudimentary valve. Florets ♂. Glumes 2, subequal, persistent, obtuse or obscurely mucronate, membranous, strongly keeled, 3-5-nerved, the lateral nerves close to the keel, the lower shorter, with the keel crested. Valves very similar, 3-nerved near the base; lateral nerves submarginal above, with 1-2 short additional nerves close to the keel. Palea slightly shorter than the valves, 2-keeled, keels winged. Lodicules 2, minute, cuneate. Stamens 3. Ovary glabrous. Styles slender from a broadened base, distinct. Stigmas plumose, laterally exserted. Grain broadly oblong to globose, broadly grooved; pericarp loose, delicate, breaking up irregularly or almost circumscissile; seed finely striate; embryo suborbicular, basal; hilum punctiform, basal.

E. indica, Gaertn.

Koster River, Rustenburg, *Burt-Davy* 146; Haenertsberg, Zoutpansberg, *Legat* in *Nat. Herb.* H. 140; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4432; Waterval Boven, Carolina, *Burt-Davy* 1432; Belfast, Lydenburg, *Burt-Davy* 1264; Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 5910.

"Goose grass." A common annual weed throughout the Transvaal.

E. coracana, Gaertn.

Devon, Bethal, *Moodie* in *Nat. Herb.* H. 3429.

A species cultivated as a forage grass and also as a cereal, from which the natives brew a kind of beer; known to the natives as M'Poko.

LXV. **DACTYLOCTENIUM**, Willd. (305).

Spikelets 3-5-flowered, laterally compressed, densely imbricate, biseriate, sessile, unilateral on a flattened rhachis, the uppermost reduced; rhachilla tardily disarticulating above the glumes, tough between the valves. Florets ♂, the uppermost rudimentary. Glumes 2, unequal, strongly keeled, the lower ovate, acute, thin, persistent, the upper elliptic-oblong in profile, obtuse, mucronate or awned, firm, deciduous. Valves ovate, subacuminate, 3-nerved, mucronate or awned, deciduous with the grains. Pales about as long as the valves, 2-keeled, subsistent. Lodicules 2, cuneate, minute. Stamens 3. Ovary glabrous. Styles distinct, very long, subterminally exerted. Grain subglobose, slightly laterally compressed, not grooved or hollowed, rugose or punctate; pericarp very delicate, irregularly breaking away; embryo scarcely equalling half the length of the grain; hilum basal, punctiform.

Annual or perennial; leaves flat, subflaccid; spikes in umbels of 2-6, erect or stelately spreading; tips of the rhachis barren, mucroniform, usually curved.

D. aegyptiacum, Willd.

Pretoria, *Reck* in *Nat. Herb.* H. 3968; *Burt-Davy* 39; Warmbaths, Waterberg, *Burt-Davy* 1758; Mosdene, Naboomspruit, *Galpin* M. 432.

"Crowfoot grass," occasionally cultivated as a hay crop.

LXVI. **LOPHACME**, Stapf (309A).

Spikelets laterally compressed, subsessile, somewhat distant on the long, slender, simple subdigitate branches of a panicle; rhachilla slender, glabrous, disarticulating above the glumes and between the two lowest valves, tough above. Florets, about 6, the lowest 2 ♂, shorter than the contiguous glumes, the following gradually reduced, barren, embracing each other and forming a tuft of awns. Glumes unequal, narrow, membranous, 1-nerved, keeled, persistent. Fertile valves linear-lanceolate in profile, 2-toothed, membranous, 3-nerved, with the side nerves evanescent above, finely awned from between the teeth; callus minutely hairy; barren valves entire, glabrous, passing into fine awns, without a callus. Pales very narrow, 2-keeled, slightly shorter than the valves. Lodicules 2, very minute, cuneate, hyaline. Stamens 3. Ovary glabrous. Styles distinct, short. Stigmas loosely plumose, laterally exerted. Grain unknown.

Perennial (?); blades flat; ligule a ciliate rim; panicle subdigitate, of very slender somewhat flexuous spike-like racemes.

L. digitata, Stapf.

Pretoria, *Macdonald* in *Nat. Herb.* H. 5468; Johannesburg, *Bryant* D. 11; Ermelo, *Burt-Davy* 5468; Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 997, H. 10644.

LXVII. **CROSSOTROPIS**, Stapf (309B).

Spikelets laterally compressed, subsessile, more or less distinctly 2-ranked on the rigid simple branches of a panicle; rhachilla slender, disarticulating above the glumes and between the valves. Florets 3-9, ♂, or the uppermost more or less reduced, equalling the glumes or slightly exerted, or over-topped by the awn-like tips of the glumes. Glumes subequal or equal, narrow, membranous, strongly 1-nerved, keeled, persistent. Valves somewhat distant, linear-oblong in profile, shortly 2-lobed, mucronate or shortly awned from the sinus, membranous, 3-nerved, side nerves submarginal, subpercurrent, rigidly ciliate, margins inflexed; callus small, hairy. Pales narrow, 2-keeled, slightly shorter than the valves. Lodicules 2, cuneate, small. Stamens 3. Ovary glabrous. Styles distinct, very slender. Stigmas plumose, laterally exerted. Grain oblong, strongly compressed from the back, concave or flat, enclosed by the slightly altered valve and pale; embryo about half the length of the grain; hilum basal, punctiform.

Annual or perennial; blades usually flat; ligules hyaline; panicle contracted and narrow, or open with the branches spreading at right angles; spikelets rather close to very distant, the uppermost terminal.

C. grandiglumis, Rendl.

The Willows, Pretoria, *Stent* in *Nat. Herb.* H. 9952; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 434; Lydenburg *Schlechter* 3948; Natalspruit, *Wentworth Sykes* in *Nat. Herb.* H. 8887; Christiana, Bloemhof, *Burt-Davy* 14093, 13041.

LXVIII. ENNEAPOGON, Desv. (310).

Spikelets 3-flowered, paniculate; rhachilla disarticulating above the glumes, minutely scaberulous or almost smooth. Lowest floret ♂, the intermediate male or barren, the uppermost rudimentary, minute. Glumes 2, persistent, membranous, acute or obtuse or minutely truncate, 3-5- or sub-7-nerved. Hermaphrodite floret; valve very broad, rounded on the back, rather firm, more or less villous, 9-nerved, 9-awned; awns subulate, equal or subequal, plumose, ciliate or scaberulous; callus minute, short. Pale oblong, 2-keeled, exceeding the valve. Lodicules 2, minute, cuneate, fleshy. Stamens 3. Ovary glabrous. Styles distinct, short. Stigmas laterally exerted, loosely plumose. Second floret like the lower, but the valve about half as long, glabrous, the ovary rudimentary or suppressed. Uppermost floret reduced to a tuft of minute awns. Grain oblong, dorsally more or less compressed; hilum punctiform, subbasal; embryo large, occupying three-quarters, or more, of the front.

Perennial, rarely subannual; blades usually narrow, often convolute; ligules reduced to a line of hairs; panicle contracted, more or less spike-like, elegantly bristly plumose from the numerous awns.

E. pretoriensis, *Stent* (*Bothalia*, I, 174).

Wonderboom, Pretoria, *Burt-Davy* 8905; northern slopes of Magaliesberg, Pretoria, *Fouche* 1.

E. scoparius, Stapf.

Bushveld, Pretoria, *Reck* in *Nat. Herb.* H. 186; Bloemhof District, Proposal, *Burt-Davy* 9473; Smitskraal, *Burt-Davy* 12915; Christiana, *Burt-Davy* 14128; Schweizer Reneke, *Burt-Davy* 1681.

LXIX. SCHMIDTIA, Steud. (312).

Spikelets closely 4-6-flowered; rhachilla disarticulating above the glumes and between the valves, joints extremely short, glabrous. Florets hermaphrodite, except the rudimentary uppermost one. Glumes 2, persistent, membranous, acute or subobtuse, 9-11-nerved. Valves broad, rounded on the back, rather firm and villous below, 9-nerved, five of the nerves excurrent into straight subulate scabrid awns, the four alternate ones into very thin lanceolate muticous lobes; callus slender, minute, bearded. Pales oblong, 2-keeled, rather longer than the body of the valves. Lodicules 2, minute, cuneate, fleshy. Stamens 3. Ovary glabrous. Styles distinct, slender. Stigmas laterally exerted; loosely plumose. Grain oblong, dorsally compressed; hilum punctiform, subbasal; embryo large, occupying three-quarters or more of the back of the grain.

Perennial or annual (?); more or less glandular-pubescent; blades rather rigid, flat, or convolute; ligule a line of hairs; panicle contracted, narrow, oblong, or spike-like; spikelets turgid, many-bristled.

S. bulbosa, Stapf.

Zandbult, near Warmbaths, Waterberg, *Bateman* in *Nat. Herb.* H. 10304; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 448; Christiana, Bloemhof, *Burt-Davy* 14098; Schweizer Reneke, *MacLaitchie* in *Nat. Herb.* H. 8963; Kaffraria, Bloemhof District, *Burt-Davy* 12945.

Bateman's specimens from Zandbult represents a form with very short awns.

This is known as "Sand quick," and is a useful pasture grass of the dry sandy regions of the Western Transvaal and Bechuanaland.

LXX. **TRIRAPHIS**, R. Br. (317).

Spikelets 5-15-flowered, laterally compressed, pedicelled, panicle; rhachilla disarticulating above the glumes and between the valves. Florets ♂, the uppermost gradually reduced. Glumes subequal, lanceolate to linear-lanceolate, acuminate or truncate, or minutely 2-toothed and aristulate, 1-nerved, keeled, thin. Valves oblong, 3-lobed, thin, 3-nerved, 3-awned, the middle lobe more or less bifid, awned from the sinus, the side lobes shorter, entire, asymmetric, awned from the inner side, margins inflexed, nerves ciliate, particularly the lateral; awns fine, scabrid, often longer than the valves; callus slender, acute, bearded. Pales linear or linear-oblong, somewhat shorter than the valves. Lodicules 2, cuneate, delicate, minute. Stamens 3. Ovary glabrous. Styles distinct, slender. Stigmas laterally exserted, very slender, plumose. Grain tightly embraced by the scarcely changed valve and pale, linear, terete, or obtusely triquetrous; embryo short; hilum basal, punctiform.

Annual or perennial; blades narrow, linear; ligule a ciliate membranous rim; panicle contracted, spike-like, or open, much-branched; spikelets distinctly pedicelled.

T. Rehmanni, Hack.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 438; Henley-on-Klip, Heidelberg, *Stent* in *Nat. Herb.* H. 21141; Welverdiend, Potchefstroom, *Burtt-Davy* 14581; Bloemhof District, Christiana, *Burtt-Davy* 1297, 13130; Schweizer Reneke, *Burtt-Davy* 1683.

T. Schlechteri, Pilg.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 439.

Perennial; about 120 cm. high, culms simple, 5-noded internodes included or shortly exserted, except the uppermost, which is very long and long exserted. Sheaths terete, tight, and finely striate, the lower splitting into rigid fibres; ligule long ciliate; blades up to about 50 cm. long and 6-7 mm. wide, long, tapering to a fine setaceous point, primary lateral nerves about five on each side of and scarcely differing from the midrib, glabrous and smooth all over or slightly scaberulous beneath. Panicle about 37 cm. long, contracted; branches in fascicles, longest about 1.5 cm., branchlets and spikelets appressed; spikelets about 14 mm. long, including the awns—glumes reddish, bifid, mucronate lanceolate, scabrid, 1-nerved, lower 5 mm., upper 6 mm. long. Valves linear-lanceolate, 5 mm. long, middle awn about 5 mm. long; scabrid slightly longer than the scabrid side awns.

This agrees with *Schlechter* 11664, from Lourenco Marques in the Kew Herbarium, and which was distributed under Pilger's name *T. Schlechteri*. I can, however, find no reference in any literature to this species, and it appears that no description has been published.

LXXI. **FINGERHUTHIA**, Nees. (326).

Spikelets strongly laterally compressed, in compact spike-like panicles, jointed on and deciduous from the pedicels; rhachilla tough. Florets 1, ♂, or if more, then the uppermost ♂ or rudimentary. Glumes 2, subequal, narrow, thin, complicate, 1-nerved, keeled, shortly awned or mucronate. Valves oblong to lanceolate, mucronate, rather firm, 7-5- (rarely 3-) nerved, the upper smaller. Pales slightly shorter than the valves, ovate-oblong, 2-keeled, flaps broad. Lodicules 2, cuneate. Stamens 3. Ovary glabrous, slightly constricted below the apex (at least, after fecundation). Styles distinct, rather long. Stigmas very slender, finely plumose, subterminally exserted. Grain unknown.

Perennial, caespitose; innovation shoots intravaginal; blades narrowly linear; ligule a dense line of silky hairs; panicle compact, spike-like; the lowest spikelets barren, consisting of a few empty glumes.

F. africana, Lehm.

Skinnners Court, Pretoria, *Stent* in *Nat. Herb.* H. 5168; Springkok Flats, Waterberg, *Sampson* in *Nat. Herb.* H. 4452, *Burt-Davy* 1129; Christiana, Bloemhof, *Burt-Davy* 13078, 13099, 14123, 1497.

LXXIIA. **ARUNDO**, Linn. (331).

Spikelets 2-7-flowered, laterally compressed, in very compound panicles, rhachilla disarticulating above the glumes and between the valves, joints short, glabrous; florets hermaphrodite, the uppermost reduced. Glumes equal, broadly lanceolate, shortly acuminate, keeled, membranous, 3-5-nerved. Valves more or less equalling the glumes, ovate to lanceolate-ovate, acuminate, finely bifid or entire, long hairy below, 5-9-nerved, three nerves more or less percurrent or excurrent, the rest short, the middle nerve often produced into a short, fine bristle; callus short, shortly bearded. Pales slightly exceeding half the length of the valve, 2-keeled. Lodicules 2, obovate, nerved, glabrous. Stamens 3. Ovary glabrous. Styles distinct, almost as long as the laterally exerted plumose stigmas. Grain obovoid-oblong, broad, loosely enclosed by the valve and pale; hilum basal, punctiform; embryo occupying almost wholly one side of the grain.

Perennial, with creeping rhizomes, extravaginal shoots and very tall and stout culms; leaves rather evenly distributed over the culms; sheaths slightly exceeding the internodes; blades long, broad, flat; ligules very short, membranous; panicles large, much compound; spikelets hairy.

A. donax, Linn.

Although this grass is not represented in the National Herbarium, it is known to occur fairly plentifully on river banks in the subtropical parts of the Transvaal.

LXXII. **PHRAGMITES**, Trin. (333).

Spikelets loosely 3-10-flowered, awnless, in large panicles; rhachilla disarticulating above the first and between the following valves, slender, glabrous, joints very short; lowest flower male or abortive, the following hermaphrodite, the uppermost florets reduced. Glumes thin, unequal to subequal, lanceolate, acute, more or less rounded on the back, 3-nerved, or the lowest sometimes sub-5-nerved. Valves heteromorphous, the lowest linear-lanceolate, much longer than the subtending glume, otherwise of a similar structure, quite glabrous, persistent, the following valves very thin, linear, long and more or less caudate-acuminate, 3-nerved, middle nerve percurrent, side nerves fine, short, callus long, slender, with very long silky hairs. Pales linear-oblong, about half as long as the valves, 2-keeled. Lodicules 2 (or sometimes 3 in the lowest floret), obovate, 2-3-nerved, glabrous, Stamens 3, or 2 in the lowest floret. Ovary glabrous, in the lowest flower rudimentary or quite suppressed. Styles distinct, rather short. Stigmas laterally exerted, densely plumose. Grain loosely enclosed by the valve and pale, free, oblong, semiterete; hilum oblong, short basal; embryo about half as long as the grain.

Perennials with a creeping rhizome; extravaginal innovation shoots, and tall sheathed annual or perennial culms; blades flat; ligule a narrow, ciliate, membranous rim; panicle lax, usually very large and much compound; spikelets conspicuously silky from the long callus hairs.

P. communis, Trin.

Skinnners Court, Pretoria, *Burt-Davy* in *Nat. Herb.* H. 610; Selous River, Rustenburg, *Burt-Davy* 158; Komatipoort, Barberton, *Burt-Davy* 358; Lichtenburg, *Burt-Davy* 62; Zeerust, Marico, *Burt-Davy* 120; Vlakfontein, near Amersfoort, Wakkerstroom, *Burt-Davy* 4039.

LXXIII. **POGONARTHRIA**, Stapf (337A).

Spikelets laterally compressed, subsessile, more or less imbricate, secund on the irregularly spirally arranged branches of a panicle; rhachilla disarticulating above the glumes and between the valves, tips of the joints ciliate. Florets 2-8, ♂. Glumes rigidly

membranous, 1-nerved. Valves oblong, rigidly membranous, acuminate, quite glabrous, 3-nerved; side nerves evanescent above the middle. Pales 2-keeled, slightly shorter than the valves. Lodicules 2, minute, delicate. Stamens 3. Ovary glabrous. Styles distinct. Stigmas plumose. Grain tightly embraced by the scarcely altered valve and pale, linear-oblong, obtusely triquetrous or oval in cross-section; embryo less than half the length of the grain; hilum basal, punctiform.

Perennial, stiff; blades rigid, usually convolute; ligule a fringe of cilia; panicles straight, with spreading, more or less curved, branches in irregular spirals; spikelets secund, crowded, livid, purplish or dark grey.

P. falcata, Rendl.

The Pyramids, Pretoria, *Burt-Davy* 4478; Henley-on-Klip, Heidelberg, *Stent* in *Nat. Herb.* H. 21145; Barberton, *Burt-Davy* 8011; Pruizen, Pietpotgietersrust, Waterberg, *Burt-Davy* 2220; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 449; Springbok Flats, Waterberg, *Burt-Davy* 1110; Warmbaths, *Burt-Davy* 2403; Slurry, Marico, *O'Connor* in *Nat. Herb.* H. 18962; Shilavane, Zoutpansberg, *Junod* in *Nat. Herb.* 4685; Waterval Boven, Carolina, *Burt-Davy* 1425; Schweizer Reneke, Bloemhof, *Burt-Davy* 1653, 1682; Christiana, Bloemhof, *Burt-Davy* 14129, 14152.

A coarse wiry grass of little value, except when quite young.

LXXIV. DIPLACHNE, Beauv. (337).

Spikelets shortly pedicelled or subsessile, somewhat distant or remote on the simple slender branches of a panicle; rhachilla disarticulating above the glumes and between the valves, glabrous. Florets 2-10, ♂, or the uppermost reduced. Glumes unequal or subequal, membranous, 1-nerved, keeled, persistent. Valves oblong to linear-oblong, 2-toothed or minutely notched, rarely quite entire, muticous or mucronulate from the sinus, very rarely shortly awned from below the apex, membranous, 3-nerved, usually finely ciliate in the lower part of the nerves or sometimes quite glabrous; side nerves percurrent or almost (or sometimes very shortly) excurrent. Pales 2-keeled, shorter than the valves. Lodicules 2, cuneate, fleshy, nerved. Stamens 3. Ovary glabrous. Styles distinct, slender. Stigmas plumose, laterally exerted. Grain enclosed by the slightly altered valve and pale, oblong to obovoid-oblong, dorsally compressed, sometimes quite flat, rarely terete; embryo equalling third to half the length of the grain; hilum punctiform, basal.

Mostly perennial; tufted, somewhat coarse grasses; blades long, narrow, flat or involute; ligules membranous, sometimes reduced to a rim; panicles consisting of slender usually long, simple, loosely spike-like and more or less distant branches; spikelets light, or olive-green, often tinged with purple and dark.

D. fusca, Beauv.

Skinnners Court, Pretoria, *Stent* in *Nat. Herb.* H. 3367; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 540; Christiana, Bloemhof, *Burt-Davy* 11410, 12788.

D. biflora, Hack.

Eloff's Plantation, Pretoria, *Macdonald* in *Nat. Herb.* H. 5446; Meintjes Kop, Pretoria, *Burt-Davy* in *Nat. Herb.* H. 3546; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4427; downs, Zoutpansberg, on open veld, *Junod* 4106; Waterval Boven, Carolina, *Burt-Davy* 1407.

LXXV. ERAGROSTIS, Beauv. (341).

Spikelets usually strongly laterally compressed, pedicelled in open or contracted panicles, rarely sessile in simple or compound spikes, very rarely articulate on the pedicels; rhachilla disarticulating above the glumes and between the valves or tough and persistent, glabrous, sometimes more or less scaberulous, very rarely minutely hairy. Florets 2 to many, ♂, or the uppermost reduced. Glumes unequal or equal, usually membranous,

1-nerved, or the upper sometimes 3-nerved, keeled, persistent or deciduous. Valves more or less imbricate, ovate to lanceolate, acute to obtuse, entire, mucous, membranous to chartaceous, 3-nerved, glabrous, very rarely minutely pubescent; side nerves short or almost percurrent. Pales equal to the valves or slightly shorter, membranous, 2-keeled, deciduous or persistent on the rhachilla. Lodicules 2, small, cuneate, more or less fleshy. Stamens 3, rarely 2. Ovary glabrous. Styles distinct. Stigmas plumose, laterally exerted. Grain enclosed by the scarcely altered valve and pale and deciduous with them, or more commonly falling with the deciduous valve, leaving the more or less persistent pale behind, oblong to obovoid or globose, round or very obtusely triquetrous or quadrangular in cross-section; pericarp thin, sometimes slightly swelling or separating; embryo often half as long as the grain (or sometimes longer); hilum punctiform, basal.

Perennial or annual, of very varying habit; blades narrow; ligule reduced to a fringe of usually minute hairs; panicles lax to effuse or contracted to spike-like, or transformed into simple or compound spikes; spikelets usually more or less olive-green or olive-grey breaking up variously, very rarely deciduous as a whole.

E. caesia, Stapf.

Belfast, Lydenburg, *Burt-Davy* 1344.

E. curvula, Nees.

Skinnners Court, Pretoria, *Burt-Davy* 4375, 4365; *Stent* in *Nat. Herb.* H. 5162; Irene, near Pretoria, *Burt-Davy* 32; Germiston, *Fenn* 8742; Elsbury, Witwatersrand, *Rogers* 12147; near Nylstroom, Waterberg, *Burt-Davy* 2065; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 451; Zoutpansberg, *Methuen* in *Nat. Herb.* H. 8802; Belfast, Lydenburg, *Burt-Davy* 1327; Helvetia, near Machadodorp, Lydenburg, *Burt-Davy* 7246; Carolina, *Burt-Davy* 2986, 7389, 2961; Athol, Ermelo, *Macdonald* in *Nat. Herb.* H. 98; Potchefstroom, *Burt-Davy* 2166; Welverdiend, Potchefstroom, *Burt-Davy* 14563; Vlaktefontein, Amersfoort, Wakkerstroom, *Burt-Davy* 4146.

A useful pasture grass.

E. lehmanniana, Nees.

Pyramid Estate, near Potgieters, Waterberg, *Galpin* 8904; Crocodile River, Rustenburg, *Burt-Davy* 9301; Schweizer Reneke, Bloemhof, *Burt-Davy* 1651, 1637; Christiana, Bloemhof, *Burt-Davy* 14139, 14149, 13447.

E. chloromelas, Steud.

Skinnners Court, Pretoria, *Stent* in *Nat. Herb.* H. 4362; Irene, near Pretoria, *Burt-Davy* 3847; Meintjes Kop, Pretoria, *Burt-Davy* 3945; Lydenburg, *Schlechter* 3969; Panplaats, Pan Station, Middelburg, *Burt-Davy* 13331; Ermelo, *Tennant* in *Nat. Herb.* H. 6954; Davel, Ermelo, *Burt-Davy* 1656; Standerton, *Burt-Davy* 3319; Potchefstroom, *Burt-Davy* 1478; Christiana, Bloemhof, *Burt-Davy* 11701; Schweizer Reneke, *Burt-Davy* 1640.

E. chloromelas would appear to be a very variable species—some of the specimens quoted above are very dissimilar in appearance and yet agree, more or less, with Stapf's description of the species. Many of the specimens previously placed under *E. chloromelas*, *E. nebulosa*, or *E. curvula* in the National Herbarium I have separated out into distinct groups which appear to me to represent distinct species. These I am recording below simply as sp. A, sp. B, etc., pending a revision of the whole genus.

E. sp. A.

Potchefstroom, *Burt-Davy* 844; Schweizer Reneke, *Burt-Davy* 1643.

A species with a rather stout, leafy culm, erect or geniculate, basal sheaths up to 9 cm. long, rigidly convolute leaves, long, rather contracted panicle and narrow linear, few-flowered spikelets.

E. sp. B.

Skinner's Court, Pretoria, *Burt-Davy* 3557.

Densely caespitose, with rigid basal sheaths, slender culms, rigidly convolute leaves, spreading ovoid panicle, and broader and rather obtuse few-flowered spikelets.

E. sp. C.

Uitgevalle, Heidelberg, *Burt-Davy* 13659; Beginsel Farm, Standerton, *Burt-Davy* 927, 1018; Christiana, Bloemhof, *Burt-Davy* 13062.

Densely caespitose, with short, whitish, rather thin basal sheaths, finer leaves, panicle ovate spreading or slightly contracted, and rather short contracted branchlets.

E. sp. D.

Henley-on-Klip, *Stent* in *Nat. Herb.* H. 21128; Pyramid Estate, Potgietersrust, *Galpin* 8901.

In habit very like sp. B, and may be the same species with narrower spikelets.

E. sp. E.

Welverdiend Station, Potchefstroom, *Burt-Davy* 14598; Christiana, Bloemhof, *Burt-Davy* 13062; Machavie Station, Potchefstroom, *Burt-Davy* 1481.

Near to *E. Burchellii* (ex descr.), but spikelets 4-7-flowered, keels of the valves scabrid.

E. sp. F.

Meintjes Kop, Pretoria, *Pole Evans* 438; Henley-on-Klip, Heidelberg, *Stent* in *Nat. Herb.* H. 21146; Pyramid Estate, Potgietersrust, *Galpin* 8901.

Culms slender, $1\frac{1}{2}$ to 2 ft. high—leaves very narrow and setaceous convolute and flexuous, panicle delicate with numerous very fine capillary divisions and scattered, 2-4-flowered, spikelets, conspicuously white bearded in the axils of the branches and branchlets

A very distinct species.

E. sp. G.

Gezina, Pretoria, *Stent* in *Nat. Herb.* H. 21470; Pyramid Estate, near Potgietersrust, *Galpin* 8903; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 453.

E. nebulosa, Stapf.

Vlaktfontein, Ermelo, *Burt-Davy*, 4164; Uitgevalle, Heidelberg, *Burt-Davy* 13672.

E. porosa, Nees.

Christiana, Bloemhof, *Burt-Davy* 12795, 12803.

E. bicolor, Nees.

Pyramid Estate, Potgietersrust, *Galpin* 8899; Springbok Flats, Waterberg, *Burt-Davy* 1132.

E. Wilmsii, Stapf.

In moist places near Pretoria, *Schlechter* 4155.

E. Burttdavii, *Stent* (in *Bothalia*, I, 176).

Bloemhof, *Burt-Davy* 10506.

E. sporoboloides, Stapf.

Pretoria, *Macdonald* in *Nat. Herb.* H. 5478; near Bamboesspruit, Bloemhof, *Burt-Davy* 1519.

MacDonald's specimen has a panicle 15 cm. long and often 5-flowered spikelets and may, when the revision of the genus is undertaken, have to be placed under a separate species.

E. atherstonei, Stapf.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 443; Warmbaths, Waterberg, *Burt-Davy* 859; Pyramid Estate, Potgietersrust, Waterberg, *Galpin* 8902; near Christiana, Bloemhof, *Burt-Davy* 1608.

One of the dominant grasses round Vryburg, Cape Province, and found mostly in association with *E. lehmanniana*. Atherston's specimen from Lydenburg in the Kew Herbarium, on which Stapf founded the species, is very poor and not typical of the grass—it has a narrow panicle, erect unbranched culms, and 3-4-flowered spikelets. Round Vryburg, where the grass is abundant, the culms are usually geniculate and much-branched and the panicles often 8 cm. broad and 17 cm. long, while glands are not always present below the nodes. There is no doubt, however, that the grass collected by Atherston at Lydenburg is the same as those quoted above and as the Vryburg grass.

***E. plana*, Nees.**

Pretoria, *Burt-Davy* 827; near Nylstroom, Waterberg, *Burt-Davy* 1998, 2082; Pyramid Estate, Potgietersrust, *Galpin* 8897; Mosdene, Naboomspruit, *Galpin* M. 538; Houtboschberg, *Burt-Davy* 1256; Vlaktefontein, near Amersfoort, Wakkerstroom, *Burt-Davy* 4011; Standerton, *Burt-Davy* 889; Carolina, *Burt-Davy* 2964; Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 5909.

"Os-gras." "A hard, wiry, tufted grass, much grazed, when young, by horses, mules, and oxen, but soon becoming too hard and wiry."—*Burt-Davy*.

"Only grass I have found that can hold its own with Kikuyu."—*MacCallum*.

***E. biflora*, Hack.**

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 445; Mailas Kop, *Schlechter* 4559.

E. sp. H.

Turffontein, Johannesburg, *Bryant* B. 68; Uitgevalen, Heidelberg, *Burt-Davy* 9139; Vierfontein, Schweizer Reneke, Bloemhof, *Burt-Davy* 1647.

Annual; near to *E. bicolor*, but taller and more robust with larger spikelets of usually 3 florets. Rhachilla continued beyond the terminal floret.

***E. Galpinii*, Stent (in *Bothalia*, I, 176).**

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 456; Pyramid Estate, near Potgietersrust, Waterberg, *Galpin* 8900; near Nylstroom, Waterberg, *Burt-Davy* 2034; Warmbaths, Waterberg, *Burt-Davy* 1149; Letaba River, Zoutpansberg, *Junod* 2350.

***E. patentissima*, Hack.**

Experiment Farm, Ermelo, *Burt-Davy* 9364.

***E. denudata*, Hack.**

Groenkloof, Pretoria, *Phillips* H. 21682; Warmbaths, Waterberg, *Burt-Davy* 1144; Vierfontein, Schweizer Reneke, Bloemhof, *Burt-Davy* 1624; Christiana, Bloemhof, *Burt-Davy* 12992, 14130, 11412, 11397.

***E. sclerantha*, Nees.**

Spionkop, Ermelo, *Burt-Davy* 9240; Booth and Bell in *Nat. Herb.* H. 3400; Grasdale, Ermelo, *Burt-Davy* 9263.

***E. chalcantha*, Trin.**

Eloff's Plantation, Pretoria, *Macdonald* in *Nat. Herb.* H. 5447; Sunnyside Kopjes, Pretoria, *Macdonald* in *Nat. Herb.* H. 5457; Skinners Court, Pretoria, *Burt-Davy* 3805; Ermelo, *Burt-Davy* 1668; Derby Station, Marico, *Burt-Davy* 7167; Beginsel Farm, Standerton, *Burt-Davy* 930, 1017; Blesbokspruit, Ermelo, *Burt-Davy* 951; Henley-on-Klip, Heidelberg, *Stent* in *Nat. Herb.* H. 21136; Uitgevalen, Heidelberg, *Burt-Davy* 9159; Pan Station, Middelburg, *Burt-Davy* 13308; near Zeerust, Marico, *Burt-Davy* 7193; downs, Zoutpansberg, *Junod* 4097, 4102; Potchefstroom, *Burt-Davy* 1066; Vaal River, Bloemhof, *Burt-Davy* 1506.

"A short leafy grass, common on Rooi grasveld and giving a nice leafy bite for sheep."—*Burt-Davy*.

E. pallens, Hack.

Mosdene, Naboomspruit, Waterberg, *Galpin* M. 459; Vaal River, Bloemhof, *Burtt-Davy* 1058; near Christiana, Bloemhof, *Burtt-Davy* 9623.

Burchell's 2190, on which Stapf has founded his species *E. dura*, is an undeveloped specimen of *E. pallens*.

E. major, Hack.

Hamanskraal, Pretoria, *Burtt-Davy* 1092; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 462; Warmbaths, Waterberg, *Burtt-Davy* 1143; Standerton, *Burtt-Davy* 1782; Vlakfontein, Amersfoort, Wakkerstroom, *Burtt-Davy* 4056; Potchefstroom, *Burtt-Davy* 1216.

"Stink gras," an introduced weed.

E. barbinodis, Hack.

Wonderboom, Pretoria, *Pole Evans* in *Nat. Herb.* H. 21472; Hamanskraal, Pretoria, *Burtt-Davy* 1087; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 450, 455; Naboomspruit, *Mogg* in *Nat. Herb.* H. 11665; near Nylstroom, *Burtt-Davy* 2066, 1997; Krabbefontein, Tzaneen, Zoutpansberg, *Burtt-Davy* 2368.

Perennial; often with long surface stolons that root at the nodes and send up fascicles of barren shoots and flowering culms. Panicle often more or less contracted and up to 20 cm. long.

E. superba, Nees.

Skidders Court, Pretoria, *Burtt-Davy* 5104; Hamanskraal, Pretoria, *Burtt-Davy* 1093; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 463; Springbok Flats, Waterberg, *Burtt-Davy* 9078; Warmbaths, *Burtt-Davy* 2227; Komatipoort, Barberton, *Burtt-Davy* 371; near Magatas Nek, Zoutpansberg, *Burtt-Davy* 152; Christiana, Bloemhof, *Burtt-Davy* 14140, 14095.

E. brizoides, Nees.

Eloff's Plantation, Pretoria, *Macdonald* in *Nat. Herb.* H. 5475; Skidders Court, Pretoria, *Burtt-Davy* 1181; The Willows, Pretoria, *Burtt-Davy* 4490; Houghton Estate, Johannesburg, *Bryant* C. 34; Pyramid Estate, Potgietersrust, *Galpin* 8906; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* H. 4463; crags and downs, Zoutpansberg, *Junod* 4095; Belfast, Lydenburg, *Burtt-Davy* 1374; Amersfoort, Wakkerstroom, *Burtt-Davy* 9047; Uitgevalen, Heidelberg, *Burtt-Davy* 9163; Beginsel Farm, Standerton, *Burtt-Davy* 3079; Amsterdam, Ermelo, *MacDonald* in *Nat. Herb.* H. 100; Ermelo, *Burtt-Davy* 3877, *Tennant* in *Nat. Herb.* H. 6970, *Burtt-Davy* 3917.

E. obtusa, Munro.

Skidders Court, Pretoria (cult. ?), *Aaron* in *Nat. Herb.* H. 7902; Potchefstroom, *Burtt-Davy* 1064; Christiana, Bloemhof, *Burtt-Davy* 12497, 1512, 11411.

E. lappula, Nees, var. *divaricata*, Stapf.

Skidders Court, Pretoria, *Burtt-Davy* 1021; Potgietersrust, Waterberg, *Burtt-Davy* 2400.

E. aspera, Nees.

Potgietersrust, Waterberg, *Burtt-Davy* 2418; Magoebas Kloof, Zoutpansberg, *Mogg* in *Nat. Herb.* H. 10668; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 446; Tzaneen, Zoutpansberg, *Sampson* in *Nat. Herb.* 4443.

E. gummiflua, Nees.

Skidders Court, Pretoria, *Burtt-Davy* 3806; Turffontein, Johannesburg, *Bryant* D. 5; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 457; Pyramid Estate, Potgietersrust, Waterberg, *Galpin* 8905; Springbok Flats, Waterberg, *Burtt-Davy* 1127; Warmbaths, Waterberg, *Burtt-Davy* 1146; Potgietersrust, *Burtt-Davy* 2340; downs, Zoutpansberg, *Junod* 4105; Waterval Boven, Carolina, *Burtt-Davy* 1440; Ermelo, *Burtt-Davy* 1658; Potchefstroom, *Burtt-Davy* 1070; Christiana, Bloemhof, *Burtt-Davy* 12785, 14153.

E. ciliaris, Link.Barberton, *Burtt-Davy* 8012.**E. namaquensis**, Nees.Pyramid Estate, Potgietersrust, *Galpin* 8896 ; Mosdene, Naboomspruit, *Galpin* M. 539.**E. viscosa**, Trin. (Mem. Acad. Petersb., Ser. VI, i, 1831, 397).Pyramid Estate, Potgietersrust, *Galpin* 8908.**E. abyssinica**, Schrad. [Linnaea XII (1838), 450].Skinners Court, Pretoria, *Burtt-Davy* 2126 ; Groenkloof, Pretoria, *Goodwin* in *Nat. Herb.* H. 8782 ; Mosdene, Naboomspruit, Waterberg, *Galpin* M. 440 ; Wel-verdiend Station, Potchefstroom, *Burtt-Davy* 14596.

Escape from cultivation—"Teff."

LXXVI. **KOELERIA**, Pers. (346).

Spikelets laterally compressed in spike-like panicles ; rhachilla glabrous or finely hairy, disarticulating above the glumes and between the valves, produced with or without a rudimentary valve. Florets 1-5, ♂, or the uppermost more or less reduced. Glumes 2, persistent, subequal or unequal, subacute to acuminate, keeled, the lower usually 1-nerved or like the upper 3-nerved, margins hyaline. Valves exceeding the glumes, acute or obtuse with the margins and tips broadly hyaline, 3-5-nerved ; side nerves usually faint, conniving above, middle nerve percurrent or excurrent into a mucro or a short subterminal awn ; callus very minute, glabrous. Pales shorter than the valves or almost as long, 2-keeled, 2-toothed, conspicuously hyaline and white. Lodicules 2, hyaline. Stamens 3. Ovary glabrous. Styles distinct, very short. Stigmas laterally exerted, plumose. Grain oblong, laterally compressed, whitish, soft, tightly embraced by the hardened back of the valve ; hilum basal, short, obscure ; embryo small.

Perennial or annual ; blades usually very narrow ; ligules hyaline ; panicle usually cylindric, often interrupted, glabrous and glistening from the hyaline white margins of the valves and pales, or more or less hairy.

K. cristata, Pers.Skinners Court, Pretoria, *Stent* in *Nat. Herb.* H. 4364 ; Leeuwpoort, Carolina, *Burtt-Davy* 7438 ; Ermelo, *Burtt-Davy* 5467 ; near Amersfoort, Wakkerstroom, *Burtt-Davy* 4130 ; Volksrust, Wakkerstroom, *Burtt-Davy* 5354 ; Vlaktefontein, Wakkerstroom, *Burtt-Davy* 9038.LXXVII. **STIBURUS**, Stapf (371A).

Spikelets laterally compressed, subsessile or shortly pedicelled, in spike-like cylindric panicles ; rhachilla disarticulating above the glumes and between the valves. Florets 4-5, ♂, the uppermost reduced, shortly exerted from the glumes. Glumes equal or subequal, membranous, lanceolate, caudate-acuminate, 1-nerved. Valves very similar to the glumes, but 3-nerved ; callus very minute. Pales shorter than the valves, 2-keeled. Lodicules 2, minute, hyaline, cuneate. Stamens 3. Ovary glabrous. Styles short, distinct. Stigmas plumose, laterally exerted. Grain enclosed by the hardly changed valve and pale, free, oblong, terete ; embryo short ; hilum basal, punctiform.

Perennial ; tufted ; blades very narrow, usually subsetaceous, long ; ligule a ciliate rim ; panicle cylindric, dense, usually dark purple, greyish-villous.

S. alopecurioides, Stapf.Houtbosch, *Schlechter* 4747.

LXXVIII. **POA**, Linn. (378).

Spikelets mostly 2-6-flowered, in loose or close (rarely in spike-like) panicles; rhachilla disarticulating above the glumes and between the valves, glabrous or scantily and minutely hairy; flowers hermaphrodite or the upper imperfect. Glumes thin, membranous, keeled, acute or obtuse, 1-3-nerved. Valves membranous, sometimes rather firm, obtuse or acute, 5-7-nerved; callus small, obtuse, often with a tuft of long wool. Pales shorter than the valves, 2-keeled. Lodicules 2, more or less 2-lobed. Stamens 3. Ovary glabrous. Styles short, free. Stigmas plumose, laterally exserted. Grain ovoid, oblong or linear, often grooved, free or adherent to the pale; hilum punctiform, basal; embryo small.

Annual or perennial; blades flat and flaccid or convolute, and more or less rigid; ligules hyaline; panicles open, often effuse, rarely contracted, spike-like; spikelets rather small, awnless.

P. annua, L.

Irene, near Pretoria, *Van der Byl* in *Nat. Herb.* H. 5258; Lydenburg, *Burt-Davy* 401; Belfast, *Burt-Davy* 1390; Caledon, Lake Chrissie, Ermelo, *Hamilton* in *Nat. Herb.* H. 6740; Volksrust, Wakkerstroom, *Meyer* in *Nat. Herb.* H. 7783; Potchefstroom, *Burt-Davy* 1034.

LXXIX. **BROMUS**, Linn. (389).

Spikelets laterally compressed, at least, after flowering, variously panicle; rhachilla disarticulating above the glumes and between the valves. Florets usually numerous, ♂, the uppermost reduced. Glumes more or less unequal, acute to acuminate, persistent, lower 1-7-, upper 3-9-, nerved, and sometimes mucronate or aristulate. Valves lanceolate or broadly oblong, rounded on the back or keeled, 5-13-nerved, usually awned; awn terminal (rarely 3 or 0) or somewhat distant from the often 2-toothed tip, straight or recurved and then often loosely twisted below, not kneed. Pales entire or bifid; keels usually rigidly ciliate or ciliate. Lodicules 2, oblong or lanceolate, entire or lobed. Stamens 3, rarely 2. Ovary obovoid with an often large, villous, 3- or 2-lobed terminal appendage. Styles short, lateral on the appendage. Stigmas plumose, laterally exserted, or in the cleistogamic species permanently enclosed. Grain linear to linear-oblong, convex-concave, and usually adherent to the valve and pale, or at least to the latter; hilum filiform long; embryo small.

Annual or perennial, or very varying habit; blades linear, flat, often flaccid; ligules membranous, hyaline; panicle contracted, often very dense, or open, and even effuse, or reduced to a raceme; spikelets rather large, erect or pendulous, from ovoid to linear-cuneate; awns very long or short, sometimes reduced to a mucro or 0.

B. leptocladus, Nees.

Groenkloof, Pretoria, *Burt-Davy* 10958; Ermelo, *Burt-Davy* 4166; Volksrust, Wakkerstroom, *Nourse* in *Nat. Herb.* H. 6196.

B. unioloides, H. B. K.

Irene, near Pretoria, *Burt-Davy* 3841; Lydenburg, *Burt-Davy* 413; Vlakfontein, Amersfoort, Wakkerstroom, *Burt-Davy* 9035; Kolonies Plaats, Potchefstroom, *Burt-Davy* 9080.

"Rescue grass" (naturalized). One of the best winter grasses, relished by all stock, susceptible to smut in the Karroo.

B. tectorum, Linn.

Skinnars Court, Pretoria, *Burt-Davy* 1909.

B. inermis, L.

Groenkloof, Pretoria, *Melle* in *Nat. Herb.* H. 11555.

With the exception of *B. unioloides*, which is known as Rescue or Prairie grass, the *Bromus* spp. are introduced weeds.

LXXX. **LOLIUM**, Linn. (395).

Spikelets usually more or less compressed, 2-ranked, alternate, sessile in the hollows of the rhachis of a simple spike; rhachilla glabrous, disarticulating above the glumes and between the valves. Florets 3-11, or sometimes more, ♂ or the uppermost reduced. Glumes of terminal spikelets equal and similar, lower suppressed in the lateral spikelets, upper linear to oblong, obtuse to acute, flat or slightly rounded, coriaceous, prominently 7-9-nerved. Valves oblong, rounded on the back, subobtusely, minutely 2-toothed (or acute in profile), more or less chartaceous except at the short hyaline tips, glabrous, 5-nerved, muticous or awned; awn a straight bristle from close to the tip. Pales equalling the valves or nearly so, 2-keeled, keels more or less crested. Stamens 3. Lodicules 2, lanceolate with a lateral tooth. Ovary glabrous, truncate. Styles distinct, very short. Stigmas laterally exerted, plumose. Grain elliptic-oblong to linear-oblong, tightly enclosed by the valve and pale, adhering to both; embryo short; hilum linear, almost as long as the grain.

Annual or perennial; blades linear, flat; ligules hyaline; spikes terminal; spikelets more or less erect, 2-ranked, with the (upper) glume opposite the hollow of the rhachis.

L. temulentum, L.

Onderstepoort, Pretoria, *Burtt-Davy* in *Nat. Herb.* H. 5105; Zwartkoppies, Pretoria, *Burtt-Davy* 7095; Vogelfontein, near Germiston, Witwatersrand, *Macdonald* in *Nat. Herb.* H. 6675; Ermelo, *Burtt-Davy* 3889; Belfast, Lydenburg, *Lewis* in *Nat. Herb.* H. 5007; Amersfoort, Wakkerstroom, *Robertson* in *Nat. Herb.* H. 16576; Potchefstroom, *Burtt-Davy* 852, 1762.

"Darnel" (drabok). A weed in grain fields; injurious to stock and to humans.

L. multiflorum, Lam.

Koloneis Plaats, Potchefstroom, *Burtt-Davy* 9086.

Perennial rye-grass—escape from cultivation.

LXXXI. **OROPETIUM**, Trin. (398).

Spikelets sessile, solitary, more or less immersed in the hollows of the rhachis of a simple distichous terminal spike; rhachilla very slender, glabrous, disarticulating above the glumes and between the valves. Florets 1-4, ♂, or uppermost rudimentary, shorter than the upper glume. Glumes of terminal spikelet equal and similar, lower glume of the lateral spikelets suppressed or (if present) more or less reduced, sometimes split in two, upper very narrow, rigid, except at the hyaline margins, subacute to acuminate, 1-3-nerved, nerves often more or less confluent into a broad rigid midrib. Valves oblong or lanceolate in profile, elliptic to broadly oblong when expanded, distinctly or obscurely keeled, at least in the upper part, truncate or minutely 3-toothed, hyaline, glabrous, hairy along the nerves, finely 3-nerved; callus bearded. Pales equalling the valves, faintly 2-nerved. Lodicules 2, minute, cuneate. Stamens 3. Ovary glabrous. Styles slender, distinct. Stigmas laterally exerted. Grain oblong, subterete or slightly compressed from the sides; embryo about one-third the length of the grain; hilum punctiform, basal.

Dwarf, densely tufted perennials, rarely annuals; blades setaceous; ligule hyaline, ciliate, short; spikes very slender, straight or gradually curved; spikelets closely or loosely 2-ranked on the opposite or subopposite sides of the rhachis.

O. capense, Stapf.

Christiana, Bloemhof, *Burtt-Davy* 12782.

INDEX TO VOLUME I.

	PAGE		PAGE
Acacia Girraffae.....	196	Aloe, spicata, Linn.....	142
" spirocarpoides.....	195	" spicata, Thunb.....	142
Achneria.....	231	" spiralis, Thunb.....	143
Acokanthera spectabilis.....	59	" succotrina, Weston.....	142
Acroceras.....	227	" variegata, Linn.....	143
Actinopeltella, Doidge.....	216	" variegata, Thunb.....	142
" nitida, Doidge.....	216	" vera.....	141
Actinopeltis.....	216	" viscosa, Thunb.....	143
Agave.....	142	Amazonia Goniomae, Doidge.....	204
Agropyrum.....	234	Amphilophis.....	225, 244
Agrostis.....	228, 229, 283	" glabra, Stapf.....	244
" lacnantha, Nees.....	283	" insculpta, Stapf.....	245
" natalensis, Stapf.....	283	Anastrabe integerrima.....	207
" suavis, Stapf.....	283	Andianda.....	34
Aira.....	233	Andradea arborea.....	30
Aizoon hispanicum, Linn.....	146, 159	Andropogon.....	226, 246
" microphyllum, Bartl.....	155	" amplectens, Nees.....	247
Alambo.....	43	" annulatus, Forsk.....	245
Albizzia fastigiata.....	27	" appendiculatus, Nees.....	247
Allophylus, Linn.....	57, 59	" auctus, Stapf.....	249
" africana.....	59	" Buchanani, Stapf.....	249
" decipiens, Radkl.....	59, 61	" ceresiaeformis, Nees.....	220, 252
" erosus, Radkl.....	59, 61	" contortus, L.....	253
" melanocarpus, Radkl.....	60, 61	" cymbarius, Linn.....	251
" monophyllus, Radkl.....	59	" dichroos, Steud.....	249
Alloterpis.....	226, 257	" dregemus, Nees.....	249
" semialata, Hitch. var. Ecklonii, Stapf.....	257	" eucomis, Nees.....	247
Aloe, Linn.....	139, 141, 143	" filifolius, Steud.....	247
" and Mesembryanthemum, Genera in Thunbergs Herbarium.....	139	" flikpendulus, Hochst.....	251
" sp.....	12	" halepensis var. effusus, Stapf.....	243
" abyssinica.....	12	" hirtiflorus var. semiberbis, Stapf.....	246
" arachnoidea, Miller.....	140	" hirtus, Linn.....	249
" arachnoides, Thunb.....	139	" huillensis, Rendl.....	247
" arborescens.....	12	" intermedius var. punctatus, Stapf.....	244
" barbadensis.....	142	" monticola var. trinii, Stapf.....	244
" dichotoma, Thunb.....	140, 143	" nardus.....	66
" disticha, Thunb.....	140	" nardus var. marginatus, Stapf.....	248
" ferox.....	20	" nardus var. validus, Stapf.....	248
" fruticosa, Lam.....	142	" pertusus var. capensis, Stapf.....	245
" humilis, Miller.....	140	" plurinodis, Stapf.....	248
" humilis, Thunb.....	140	" rufus, Kunth.....	249
" latifolia, Haw.....	141	" Ruprechtii, Hack.....	251
" lingua, Thunb.....	140, 141	" Schinzii, Hack.....	247
" linguaeformis, Linn.....	140, 141	" schirensis, Hochst., var angustifolia, Stapf.....	247
" maculata, Thunb.....	141	" schoenanthus var. versicolor.....	248
" mitriformis.....	12	Anthephora.....	224, 227, 255
" natalensis.....	12	" pubescens, Nees.....	222, 255
" perfoliata, Thunb.....	141, 142	Anthistiria imberbis, Retz.....	219
" picta, Thunb.....	141	Anthoxanthum.....	233, 276
" platylepis.....	20	" odoratum, L.....	276
" plicatilis, Miller.....	140, 141	Anthracnose.....	179
" pumala, Thunb.....	141	Aphysa Rhynchosiae (Kalch. and Cke.) Theiss. and Syd.....	65
" pumila, Linn.....	141	Apicra spiralis, Baker.....	143
" retusa, Linn.....	142	" spiralis, Willd.....	143
" retusa, Thunb.....	142		
" sinuata, Thunb.....	142, 143		

	PAGE		PAGE
Apodytes dimidiata.....	210	Babala grass.....	257, 274
Apple.....	179, 182	Baker, Edmund G.—	
Areynhllum, Torrey and Gray.....	115	Revision of South African Species of	
Aristida.....	222, 229, 276	Rhynchosia.....	113
" sp.....	278	Bambusa.....	235
" adscensionis, L.....	277	Bark, bitter.....	34
<i>Aristida angustata</i> , Stapf.....	277	Bastard Sneezewood.....	34
" barbicollis, Trin. and Rupr.....	277	Beech.....	98
" bipartita, Trin. and Rupr.....	277	Behnia reticulata.....	72, 79
" Burkei, Stapf.....	277	Bermuda quick.....	288
" congesta, Roem. and Schult.....	277	Bersama, Fresen.....	53, 57, 63
" junciformis, Trin. and Rupr.....	277	" The Genus.....	33
" scabrivalvis, Hack.....	277	" abyssinica, Fresen.....	36, 38, 63, 64
" Sciurus, Stapf.....	277	Boontjes elands.....	187
" sieberiana, Trin.....	278	Boscia caffra.....	196
" spectabilis, Hack.....	278	Blauwzaad.....	264
" stipoides, Lam.....	278	Brachiaria.....	226, 261
" stipoides var. meridionalis, Stapf.....	278	" arrecta, Stent.....	263
" uniplumis, Licht.....	278	" brizantha, Stapf.....	222, 262
" vestita, Thun.....	278	" Isachne, Stapf.....	262
Arundinaria.....	235	" Marlothii, Stent.....	263
" tessellata.....	69	" mutica, Forsk.....	262
Arundinella.....	231, 256	" nigropedata, Stapf.....	222, 262
" Ecklonii, Nees.....	257	" serrata, Stapf.....	262
Arundo.....	233, 295	Brachylaena sp.....	6
" donax, Linn.....	295	" dentata.....	6
Ascomycetes, South African; in the		" discolor.....	6
National Herbarium—		" elliptica.....	6
Part I.....	5	" neriifolia.....	6
Part II.....	65	Brachypodium.....	235
Part III.....	195	Brachystegia sp.....	221
Ascostratum insigne, Syd.....	206	Briza.....	234
Asparagus sp.....	9, 16	Brizopyrum.....	230, 232
" striatus.....	9	Bromus.....	234, 235
<i>Asterella rehmi</i> , P. Henn.....	12	Bromus spp.....	302
Asterina.....	210	" inermis, L.....	302
? Asterina sp.....	210, 213, 214	" leptocladus, Nees.....	302
<i>Asterina capensis</i>	71	" lectorum, Linn.....	302
" celtidicola, P. Henn, var.....		" unioloides, H.B.K.....	302
" microspora, Doidge.....	204	Brown, N. E.—	
Asterina crotonicola, Doidge.....	76, 82	The Genera Aloe and Mesembryan-	
" dilabens, Syd.....	204	themum in Thunberg's Herbarium	139
" gerbericola, Doidge.....	202, 204, 210	Buffels gras.....	264
" macowaniana, Kalch. and Cke.....	212	Burchellia capensis.....	22
" Scolopiae, Doidge.....	77, 82	Bush tea.....	105
" solaris, K. and Cke.....	10	Buxus MacOwani.....	79
" Streptocarpi, Doidge.....	203, 216	Calamagrostis.....	229, 283
" undulata.....	202, 204	" epigeios, Rott., var. capensis,	
Asterinella.....	199	Stapf.....	284
" nimusopsidis, Doidge.....	80	Calonectria capensis, Doidge.....	218
" Pterocelastri, Doidge.....	198	Calothyrium psychotriae, Doidge.....	76
Asterodonthis.....	15	Canary grass.....	276
" solaris (K. and Cke.), Theiss.....	10, 11	Cape Plane.....	89, 93
Asteromyxa, Theiss. and Syd.....	199	Capnodium.....	213
" hirtula.....	199	" citricolum, McAlp.....	213
? " inconspicua, Doidge.....	199	Capparis citrifolia.....	204
Atropis.....	234	Cardiospermum, Linn.....	57, 58
<i>Auerswaldia disciformis</i> , Wint.....	21	" halicacabum.....	58
" examinans, (M. and B.).....	21	Carissa.....	79
Avena.....	234	" sp.....	10
Avenastrum.....	234, 284	" acuminata.....	10
" turgidulum, Stapf.....	284	" arduina.....	10, 72, 78
Avocado.....	179, 180	" grandiflora.....	10
<i>Azopopus semialatus</i> , Hook., var. <i>ecklonii</i> ,		Carrot seed grass.....	255
Stapf.....	257	Cassine capensis.....	212

	PAGE		PAGE
Cassipourea verticillata.....	79	<i>Copisma pilosum</i>	131
Catacauma dalbergiicola (P. Henn), Th. and Syd.....	25	<i>rotundifolium</i> , E. Mey.....	119
grammicum (P. Henn), Theiss. and Syd.....	26	<i>tenue</i>	126
Peglerae, Doidge.....	25	<i>totum</i>	130
Pterocarp, Syd.....	25	<i>trichodes</i> , E. Mey.....	138
Schotiae, Doidge.....	65	<i>viscidulum</i> , E. Mey.....	137
Catha edulis.....	6	Cordia cafræ.....	27
Celastrineae.....	208	<i>Corynelia clavata</i> (L.), Sacc.....	218
Celastrus.....	212	<i>carpopbila</i> , Syd.....	217
acuminatus.....	197, 208	<i>fructicola</i> (Pat.), v. Hohn.....	217
albatus.....	212	<i>tripos</i> , Cke.....	216
buxifolius.....	200, 212	<i>uberata</i> , Fr.....	218
Harveyanus.....	200	Coryneliaceae.....	195
nemorosus.....	200	Crossotropis.....	233, 292
polyacanthus.....	212	<i>grandiglumis</i> , Rendl.....	293
pyracanthus.....	200	Croton silvaticum.....	67
ruber.....	200	Crotonis rivularis.....	77
senegalensis.....	200	Crow foot grass.....	292
Cenchrus.....	227	Ctenium.....	230, 289
Chaetacme aristata.....	79	<i>coninum</i> , Nees.....	290
Chaetobromus.....	232	Cunonia capensis.....	213
Chaetostigmella.....	210	Cyanospermum, Benth.....	114
asterinicola, Doidge.....	210	Cyathea dregei.....	28
capensis, Doidge.....	210	Cyclopia, Vent. The Genus.....	105
Chaetothyrium.....	212	<i>Ashtonii</i>	106, 107, 109
Chilanthus arboreus.....	199, 206	<i>Bolusii</i>	107, 109
Chloris.....	230, 290	<i>brachypoda</i>	106, 108
gayana, Kunth.....	290	<i>brachypoda</i> , var. <i>intermedia</i>	108
petraea, Thunb.....	231, 290	<i>galiodes</i> , D. C.....	107, 109
pycnothrix, Trin.....	290	<i>Burtonii</i>	106, 107
virgata, Schw.....	290	<i>genistoides</i> , Br.....	105, 106, 108
Chrysopogon.....	225, 244	<i>latifolia</i> , D. C.....	106, 107
montanus, Trin. var. <i>tremulus</i> , Stapf.....	244	<i>montana</i>	107, 109
monticola.....	219	<i>montana</i> var. <i>glabra</i>	109
Chrysosciæ, Benth.....	114, 115	<i>pubescens</i> , E. and Z.....	107, 108
<i>Chrysosciæ calycina</i> , E. Mey.....	116	<i>sessiliflora</i> , E. Mey.....	107, 109
<i>grandiflora</i> , E. Mey.....	115	<i>subternata</i> , Vog.....	106, 107
<i>media</i> , E. Mey.....	116	<i>tenuifolia</i> , Lehn.....	105, 106, 108
<i>parviflora</i> , E. Mey.....	116	<i>Vogelii</i> , Harv., var. <i>brachypoda</i> , Harv.....	108
Citrus.....	179	<i>Vogelii</i> , Harv., var. <i>intermedia</i>	108
sp.....	21, 213	<i>Vogelii</i> , Harv., var. <i>subternata</i> , Harv.....	107
Claviceps Paspali (Ergot).....	258	Cycloschizon.....	6
Cleistachne.....	224, 253	<i>alyxiae</i> (Mass).....	6
sorghoides, Benth.....	253	<i>brachylaenae</i> (Rehm.), P. Henn.....	6
? Clypeolella, sp.....	210, 214	<i>fimbriatum</i> , Doidge.....	6
Clypeosphaeria natalensis, Doidge.....	81	<i>pritzeli</i> , P. Henn.....	6
Cocconia capensis, Doidge.....	8	Cyclothea Bosciæ, Doidge.....	196
concentrica, Syd.....	7	<i>Cylista angustifolia</i> , Ecklon and Zeyher.....	116
<i>parinarii</i> , P. Henn.....	26	<i>angustifolia</i> , E. Mey.....	116
porrigo (Cke.), Sacc.....	7	<i>argentea</i> , Ecklon. & Zeyher.....	116
Coffee Pear (Pleurostyliæ).....	34	<i>lancifolia</i>	115
Common bush tea.....	106	Cymbopogon.....	226, 247
Conophytum truncatum, N. E. Br.....	165	<i>excavatus</i> , Stapf.....	248
Copisma.....	114, 117	<i>marginatus</i> , Stapf.....	248
<i>Copisma diversifolium</i> , E. Mey.....	126	<i>plurinodes</i> , Stapf.....	248
<i>effusum</i> , E. Mey.....	128	<i>validus</i> , Stapf.....	248
<i>falcatum</i> , E. Mey.....	127	Cynodon.....	230, 288
<i>gibbum</i> , E. Mey.....	125	<i>dactylon</i> , Pers.....	67, 222, 288, 289
<i>glabrum</i> , E. Mey.....	132	<i>incompletus</i> , Nees.....	67, 289
<i>glandulosum</i> , E. Mey.....	132	<i>transvaalensis</i>	289
<i>grandifolium</i> , E. Mey.....	120	Cynosurus.....	233, 235
<i>nitidum</i> , E. Mey.....	138	Dactylis.....	234
<i>paniculatum</i> , E. Mey.....	130	Dactyloctenium.....	230, 292
<i>pictum</i> , E. Mey.....	125		

	PAGE		PAGE
<i>Eragrostis</i> Burtt-davii, Stent.....	176, 223	Flacourtiaceae.....	83
" caesia, Stapf.....	297	Florida Grass.....	289
" chalcantha, Trin.....	222, 299		
" chloromelas, Steud.....	297	<i>Galega pinnata</i> , Thunberg.....	117
" ciliaris, Link.....	301	<i>Galenia</i>	155
" curvula, Nees.....	172, 222	" filiformis, N. E. Br.....	155
" curvula, var. valida.....	172, 297	<i>Galenia herniariaefolia</i> , Fenzl.....	155
" denudata, Hack.....	299	<i>Gasteria</i>	140, 141, 143
" dura, Stapf.....	300	" disticha.....	141
" echinochloidea.....	222	" pulchra, Haw.....	141
" Galpinii, Stent.....	176, 299	" Thunbergii, N. E. Br.....	140
" gangetica, Steud.....	178	" verrucosa.....	140
" gummiflua, Nees.....	300	<i>Gerbera cordata</i>	202, 211
" lappula, Nees, var. divaricata		<i>Germiston grass</i>	289
Stapf.....	300	<i>Gloeosporium</i>	179
" lehmanniana, Nees.....	222, 297, 299	<i>Glottidium</i>	40
" major, Hack.....	300	<i>Glottiphyllum longum</i> , N. E. Br.....	157
" namaquensis, Nees.....	300	<i>Glycine angustifolia</i> , Jacq.....	113, 116
" nebulosa, Stapf.....	297	" argentea, Thunb.....	126
" obtusa, Munro.....	222, 300	" densiflora, Roth.....	115
" pallens, Hack.....	300	" erecta, Thunbg.....	115
" patentissima, Hack.....	299	" glabra, Spreng.....	132
" plana, Nees.....	222, 299	" glandulosa, Thunberg.....	132
" porosa, Nees.....	298	" heterophylla, Thunbg.....	132
" sclerantha, Nees.....	299	" secunda, Thunberg.....	121
" sporoboloides, Stapf.....	298	" Totta, Thunbg.....	130
" superba, Nees.....	300	<i>Gonioma Kamassi</i>	204
" valida, Stent.....	172	<i>Goose grass</i>	291
" viscosa, Trin.....	301	<i>Gramineae</i>	170
" Wilmsii, Stapf.....	298	<i>Graminae, South African—</i>	
<i>Ergot</i> (Claviceps Paspali).....	258	A New Genus and Seven New Species	170
<i>Erianthus junceus</i> , Stapf.....	236	Graminae of the Transvaal.....	222
<i>Eriochloa</i>	226, 258	Key to the South African Genera....	224
" acrotricha, Hack.....	260	<i>Grass—</i>	
" Fouchei, Stent.....	259, 260	babala.....	257, 274
" ramosa, Kuntz.....	260	bermuda quick.....	288
<i>Eriochrysis</i>	225	Blauwzaad.....	264
<i>Eriosema puberulum</i> , Ecklon. and Zeyher...	130	buffels.....	264
" sericeum, Ecklon. and Zeyher....	133	canary.....	276
<i>Euclea macrophylla</i>	16	carrot seed.....	255
" natalensis.....	15	crow foot.....	292
<i>Eugenia</i> sp.....	79	dek.....	222
" capensis.....	25	ditch.....	258
" zeyheri.....	75	dobo.....	89
<i>Eulalia</i>	224, 236	esparte.....	251
" geniculata, Stapf.....	236	elephant.....	48
" villosa, Nees.....	236	finger.....	268
<i>Eu-Montagnelleae</i>	219	Florida.....	289
<i>Euphorbia</i> sp.....	206	Germiston.....	289
" schintzii.....	23	goose.....	291
<i>Eurhynchosis</i>	114, 117	guinea.....	264
<i>Eusesbania</i>	40, 41, 42	Indian buffels.....	265
<i>Eustachys</i>	231	kits.....	273
" a capensis.....	209	kodo millet.....	258
Davyi.....	209	kopje.....	272
" bituminosa, D. C.....	114, 138	krul.....	262
" McNaughtonii.....	197	kwas.....	290
" a.....	234, 235	millet large seeded.....	262
" vulpioides, Steud.....	234	Molasses.....	257
" sp.....	24, 25, 29	Natal buffels.....	272
" capensis.....	26, 29	Natal red top.....	271
" ochstetteri, var. glabrior.....	24	Old land.....	290
" howardiana.....	24	Os.....	299
<i>Finger grass</i>	268	para.....	262
<i>Fingerhuthia</i>	231, 294	pearl millet.....	274
<i>Fingerhuthia africana</i> , Lehm.....	222, 232, 294	penhalonga.....	262
		perennial rye.....	303

	PAGE		PAGE
Grass—		<i>Hyparrhenia variabilis</i> , Stapf.....	251
prairie or rescue.....	302	<i>Hysteroglyphium</i> Acaciae, Doidge.....	196
reed canary.....	276	" Fraxini (Pers), De Not.,	
regte kweek.....	288	var. <i>Oleastri</i> , Desm ..	195
Rhodes.....	290	<i>spenicolum</i> , Doidge.....	195
rooi.....	254	<i>Hysterostoma</i>	15
South African millet.....	273	" Faureae, Doidge.....	197
spear.....	258	" <i>orbiculata</i> , Syd.....	13
steek.....	253	<i>Hysterostomella circularis</i> , Har. and Pat....	7
stink.....	300	" <i>concentrica</i> , Syd.....	7
Sweet vernal.....	276	" <i>tenella</i> , Syd.....	9
tambookie.....	222, 251	<i>Hysterostomina</i>	9, 10
turf.....	238	" <i>tenella</i> , Syd.....	9
turpentine.....	248	<i>Ibbetsonia</i>	105
Warmbaths.....	262	<i>Imperata</i>	224, 235
wildebeest.....	242	" <i>arundinacea</i> Cyr., var. <i>Thunbergii</i> ,	
Zuurpol.....	242	Hack.....	235
Greyia, Hook. and Harv.....	57, 59	Indian Buffels grass.....	265
" Sutherlandi, Hook. and Harv.....	59	? <i>Ipomoea</i> sp.....	208
<i>Grumilea capensis</i>	201	<i>Irene</i>	81, 209
Guinea grass.....	264	" <i>implicata</i> , Doidge.....	206
<i>Halleria elliptica</i>	17	" <i>Peglerae</i> , Doidge.....	207, 209
" <i>lucida</i>	17, 207, 209	" <i>Podocarp</i>	218
<i>Harpechloa</i>	230, 289	" <i>Rinorea</i> , Doidge.....	80
" <i>capensis</i> , Kunth.....	289	" <i>Zeyheri</i> , Doidge.....	75
<i>Haworthia</i>	141	<i>Ischaemum</i>	225, 236
" <i>arachnoidea</i> , Duval.....	140	" <i>arcuatum</i> , Stapf.....	238
" <i>arachnoidea</i> , Haw.....	140	" <i>faciculatum</i> , var. <i>arcuatum</i> , Hook.....	238
" <i>fasciata</i> , Haw.....	141	" <i>glaucoctachyum</i> , Stapf.....	238
" <i>granata</i> , Haw.....	141	" <i>Mellei</i> , Stent.....	237, 238
" <i>margaritifera</i> , Haw.....	141	" <i>purpurascens</i> , Stapf.....	238
" <i>retusa</i> , Duval.....	142	<i>Isipinga</i> , Doidge.....	15
<i>Haworthia viscosa</i> , Haw.....	143	" <i>areolata</i> , Doidge.....	15
<i>Heidelberg tea</i>	106	" <i>contorta</i> , Doidge.....	15
<i>Helichrysum</i> sp.....	70	<i>Jasminum angulare</i>	80
<i>Helminthosporium</i> sp.....	209	<i>Keur</i>	98
<i>Hemarthria</i>	224, 241	<i>Kiggelaria africana</i>	204
<i>Hemarthria fasciculata</i> , Stapf.....	241	<i>Klits</i> gras.....	273
<i>Heteropogon</i>	225, 252	<i>Knowltonia vesicatoria</i>	208
<i>Heteropogon contortus</i> , Roem. and Schult.....	253	<i>Kodo Millet</i>	258
<i>Heuning tea</i>	106	<i>Koelaria</i>	233, 234, 301
<i>Hidrosia bullata</i> , E. Mey.....	136	<i>Koelaria cristata</i> , Pers.....	222, 301
<i>Hippobromus</i> , E. and Z.....	57, 61	<i>Koker Boom</i>	140
" <i>alatus</i> , E. and Z.....	61, 71	<i>Kopje gras</i>	272
Hofmeyer, Joan, and E. P. Phillips—		<i>Krul gras</i>	262
The Genus <i>Olinia</i>	97	<i>Kwas gras</i>	290
The Genus <i>Cyclopia</i> , Vent.....	105	Kweek —	
<i>Holcus</i>	233, 284	regte.....	288
" <i>lanatus</i> , L.....	284	water.....	275
<i>Homostegia albizziae</i> , (Cke.).....	27	<i>Lagurus</i>	229
" <i>amphimelaena</i> ,.....	18	<i>Lamarkia</i>	232
Honey tea.....	105, 106	<i>Lasiochloa</i>	231, 232
Hordeum.....	229	<i>Leersia</i>	228, 275
Hottentots tea.....	106	" <i>hescandra</i> , Sw.....	275
<i>Hyparrhenia</i>	226, 248	<i>Leguminosae</i>	5, 41, 113, 187
" <i>auctus</i> , Stapf.....	249	<i>Lembosia</i>	5
" <i>Buchanani</i> , Stapf.....	249	" <i>albersii</i>	10
" <i>cymbaria</i> , Stapf.....	251	" <i>congesta</i> , Wint.....	9
" <i>dichroa</i> , Stapf.....	249	" <i>radiata</i> , Doidge.....	5
" <i>dregeana</i> , Stapf.....	249	" <i>piriensis</i> , Doidge.....	78
" <i>filipendula</i> , Stapf.....	251	<i>Lembosiodothis</i>	9
" <i>glauca</i> , Stent.....	250, 251	" <i>Leptocarydon</i>	233
" <i>hirta</i> , Stapf.....	249	" <i>Vulpiastrum</i> , Stapf.....	220
" <i>rufa</i> , Stapf.....	249	<i>Lepturus</i>	229
" <i>Ruprechtii</i> , Fourn.....	251		
" <i>Tamba</i> , Anderss.....	251		

	PAGE		PAGE
<i>Lessertia tenuifolia</i>	30	<i>Mesembryanthemum</i> bicolorum, Thunb.....	149
<i>Leucophrys</i>	227	" brachiatum, Ait.....	152
<i>Lolium</i>	234, 303	" bracteatum, Ait.....	149, 162
" multiflorum, Lam.....	303	" brevifolium, Ait. 149, 150, 156	156
" temulentum, L.....	303	" brevifolium, Haw.....	149
<i>Lophacme</i>	232, 292	" brevifolium, Salm Dyck	149
" digitata, Stapf.....	292	" brevifolium, Sond.....	156
<i>Machailo</i>	93	" calamiforme, L.....	149
<i>MacOwaniella</i> , Doidge.....	9, 10	" calcareum, Marle.....	163
" congesta (Wint.), Doidge... 9, 10		" calycinum, Haw.....	156
" myrsinicola, Doidge.....	196	" candens, Haw.....	156
<i>Maerua pedunculosa</i>	204	" caninum, Lam.....	167
<i>Marattia fraxinea</i>	219	" capense, Houtt.....	168
<i>Mbomvane neinci</i>	89	" capillare, Linn. f... 149, 150	150
<i>Melanose</i>	179	" capillare, Thunb... 149, 156	156
<i>Melanthus</i> , Linn.....	57, 58	" ciliatum, Ait. Thunb... 151	151
" insignis.....	58	" clavatum, Haw.....	157
" major.....	28	" clavatum, Jacq.....	164
" villosus, Bolus.....	58	" coccineum, Haw.....	149
<i>Melica</i>	233	" collinum, Sond.....	158
<i>Melinis</i>	224, 228, 257	" copticum, Jacq.....	146
" minutiflora.....	257	" copticum, Linn.....	159
" minutiflora, var. pilosa.....	257	" corallinum, Thunb....	151
" tenuinervis, Stapf.....	257	" cordifolium, Linn. f.	
<i>Meliola</i>	209, 215	Thunb.....	152
" amphitricha.....	73, 76	" crassifolium, Linn... 154, 158	158
" capensis (K. and Cke.), Th.....	71, 82	" crassifolium, Thunb... 152	152
" carissae, Doidge.....	72, 82	" criniflorum, Linn. f.	
" comata, Doidge.....	208, 215	Thunb.....	152, 157
" dracoenicola, Pat. and Har.....	72, 82	" crystallinum, Linn.	
" Evansii.....	208	Thunb.....	152
" gemellipoda, Doidge.....	80	" crystallophanes, Eckl.	
" hyalophragmia.....	212	and Zey.....	145
" Knowltoniae, Doidge.....	208	" cymbiforme, Haw.....	162
" macowaniana, Thum.....	212	" decussatum, Thunb. 152, 153	153
" oleicola, Doidge.....	73, 82	" debile, Haw.....	152
" Panici, Earle.....	74, 82	" defoliatum, Haw.....	164
" Toddaliae, Doidge.....	80, 209	" deltoides, Linn. Thunb.	153
<i>Meliolina</i>	209, 210	" denticulatum, Haw... 153	153
<i>Mesembryanthemum</i>	139, 143, 151, 155	" difforme, Linn.....	153
" Genera Aloe and, in		" difforme, Thunb....	153
Thunbergs Herbarium	139	" digitatum, Ait.....	153
" acinaciforme, Linn 145, 156		" digitiforme, Thunb... 153	153
" aculeatum, N. E. Br... 163		" dissimile, N. E. Br... 157	157
" Aitonis, Jacq. 145, 146, 160		" dolabriforme, Linn... 153	153
" albidum, Linn. Thunb. 145		" echinatum, Ait.....	161
" anatomicum, Haw..... 154		" Ecklonis, Salm Dyck.	167
" angulatum, Thunb... 145, 160		" edule, Linn. Thunb... 154	154
" angulatum, var. gracile, Sond.....	146	" elongatum, Eckl. and Zey.....	160
" angulatum, var. ovatum, Sond.....	160	" emarcidum, Thunb... 154	154
" apetalum, Linn. f... 145, 159		" emarginatum, Linn.	
" apetalum, Thunb.... 146		Thunb.....	154
" arboriforme, Burch... 166		" erectum, Haw.....	155
" articulatum, Thunb... 144, 146, 147, 162		" erigeriflorum, Jacq. 149, 150	150
" auratum, Sond..... 148		" exiguum, N. E. Br... 153	153
" aureum, Linn..... 148		" expansum, D. C..... 157	157
" aureum, Thunb.... 148		" expansum, Linn... 154, 157	157
" barbatum, Berger.... 148		" expansum, Thunb.... 154	154
" barbatum, Haw..... 148		" expersum, N. E. Br... 158	158
" barbatum, Linn..... 148		" falcatum, Linn..... 164	164
" barbatum, Thunb.... 148		" falcatum, Thunb.... 154	154
" bellidiflorum, Thunb... 148		" fasciculatum, Thunb. 154, 165	165
" bicolor..... 149		" fastigiatum, Haw..... 155	155
		" fastigiatum, Thunb... 155	155
		" felinum, Hill..... 161	161
		" filicaule, Haw..... 152	152

	PAGE		PAGE
<i>Mesembryanthemum filiforme</i> , Thunb.	155	<i>Mesembryanthemum spathulatum</i> , Thunb. .	162
" <i>floribundum</i> , Haw.	156	" <i>spathulatum</i> , Willd. 152,	162
" <i>forficatum</i> , Thunb. 145,	156	" <i>spathulifolium</i> , Berger	162
" <i>geniculiflorum</i> , Linn.		" <i>spectabile</i> , Haw.	167
Thunb.	153, 156	" <i>spinosum</i> , Linn.	163, 164
" <i>glaucum</i> , Linn.		" <i>spinosum</i> , Thunb.	163
Thunb.	156, 167	" <i>splendens</i> , Linn.	
" <i>glomeratum</i> , Salm Dyck	164	Thunb.	163
" <i>gracile</i> , Haw.	164	" <i>stelligerum</i> , Haw.	148
" <i>gramineum</i> , Haw.	157	" <i>stipulaceum</i> , Linn.	164
" <i>granulicaule</i> , Haw.	147	" <i>stipulaceum</i> , Thunb. .	164
" <i>Hackelianum</i> , Berger.	160	" <i>stratum</i> , N. E. Br.	166
" <i>hispidum</i> , Linn.	157	" <i>subglobosum</i> , Haw.	149
" <i>hispidum</i> , Thunb.	156	" <i>subincanum</i> , Haw.	164
" <i>itonsum</i> , Haw.	148	" <i>tenuifolium</i> , Linn. .	162, 164
" <i>juncum</i> , Haw.	148	" <i>tenuifolium</i> , Thunb. .	164
" <i>lacerum</i> , Haw.	154	" <i>testiculate</i> , Ait.,	
" <i>laeve</i> , Thunb.	157	Thunb.	164
" <i>lanceum</i> , Thunb. .	157, 159	" <i>tetragonum</i> , Moench. .	165
" <i>lateriflorum</i> , D. C.	149	" <i>tetragonum</i> , Thunb. 154,	164
" <i>limpidum</i> , Ait.	152	" <i>thunbergii</i> , Haw.	157
" <i>lineare</i> , Thunb.	157	" <i>tortuosum</i> , Linn.	
" <i>linguaeforme</i> , Thunb. .	157	Thunb.	165
" <i>linguiforme</i> , Linn.	157	" <i>trichotomum</i> , Thunb. .	165
" <i>longum</i> , Haw.	157	" <i>tricolor</i> , Sims.	157
" <i>loreum</i> , Linn.	158	" <i>tricolor</i> , Willd.	152, 157
" <i>loreum</i> , Thunb.	158	" <i>tripolium</i> , Linn.	154, 165
" <i>micans</i> , Linn. Thunb. 156,	158	" <i>tripolium</i> , Thunb. .	157, 165
" <i>moniliforme</i> , Thunb. .	158	" <i>truncatum</i> , Thunb. 165,	166
" <i>murinum</i> , Haw.	161	" <i>tuberosum</i> , Linn.	166
" <i>mustellinum</i> , Haw.	161	" <i>tuberosum</i> , Thunb. .	166
" <i>mutabile</i> , Haw.	149	" <i>umbellatum</i> , Linn.	167
" <i>namaquense</i> , Sond.	153	" <i>uncinatum</i> , Linn.,	
" <i>noctiflorum</i> , Linn.		Thunb.	167
Thunb.	158, 167	" <i>uncinellum</i> , Haw.	167
" <i>nodiflorum</i> , Linn.		" <i>varians</i> , Haw.	157
Thunb.	159	" <i>verruculatum</i> , Linn. .	167
" <i>ovatum</i> , Thunb. 157, 159,	160	" <i>verruculatum</i> , Thunb. .	167
" <i>papuliferum</i> , D. C.	155	" <i>veruculoides</i> , Sond.	167
" <i>papulosum</i> , Linn. f. 146,	160	" <i>villosum</i> , Linn.	167
" <i>parviflorum</i> , Jacq.	166	" <i>violaceum</i> , D. C.	164
" <i>pinnatifidum</i> , Lin. f.		<i>Michrochloa</i>	229, 230, 287
Thunb.	160	" <i>altera</i> , Stapf, var., Nelsonii,	
" <i>pinnatum</i> , Thunb.	161	Stapf.	288
" <i>plenifolium</i> , N. E. Br.	155	" <i>caffra</i> , Nees.	288
" <i>polyanthon</i> , Haw.	164	" <i>setacea</i> , R. Br.	288
" <i>polyanthon</i> , Salm		<i>Microcycylus osyridis</i> , Sacc.	18
Dyck.	164	<i>Microcycylus tassianus</i> , Syd.	18
" <i>pomeridianum</i> , Linn.		<i>Micropeltis Marathiae</i> , P. Henn.	219
Thunb.	152, 161	<i>Mikania capensis</i>	210
" <i>praecultum</i> , N. E. Br.	156	<i>Millet</i> —	
" <i>productum</i> , Haw.	164	<i>kodo</i>	258
" <i>pruinosum</i> , Thunb.	161	<i>large seeded</i>	262
" <i>pugioniforme</i> , Houtt 152,	161	<i>pearl</i>	274
" <i>pugioniforme</i> , Linn.	161	<i>South African</i>	273
" <i>pugioniforme</i> , Thunb.	161	<i>Mimusops obovata</i>	80
" <i>pyropaeum</i> , Haw. .	152, 157	<i>Miscanthidium</i>	224, 235
" <i>reptans</i> , Ait.	152	" <i>juncum</i> , Stapf.	236
" <i>ringens</i> , Thunb.	161	<i>Monocymbium</i>	225, 252
" <i>roseum</i> , Willd.	156	" <i>cerisiiforme</i> , Stapf.	252
" <i>rostratum</i> , Linn. .	153, 161	<i>Montaganelia maxima</i> , Mass.	20
" <i>sabulosum</i> , Thunb.	161	<i>peglerae</i> , Pole Evans.	20
" <i>scabrum</i> , Linn.	162	<i>Moopa-Koma</i>	86
" <i>secundum</i> , Thunb. 147,	162	<i>Morelle</i>	93
" <i>serratum</i> , Linn.	162	<i>Morenoella</i>	206
" <i>serratum</i> , Thunb.	162	<i>Phillipsii</i> , Doidge.	205
" <i>sessile</i> , Thunb.	150, 162	<i>Mosdenia</i>	170, 223, 229, 256

	PAGE		PAGE
Mosdenia waterbergensis, Stent.....	170, 256	Oryza sativa, L.....	275
M'Poko.....	291	Oryzopsis.....	229, 278
Myrica sp.....	21	" sp.....	278
Myrsine melanophlebos.....	11, 20, 196, 217	Os-gras.....	299
Mystroxydon.....	208	Osyris compressa.....	18
Natal Buffels gras.....	272	Oxyanthus Gerrardi.....	201
Natal Red Top.....	271	Palawaniella Dovyvalidis, Doidge.....	198
Natal Species of the Sapindaceae.....	57	" eucleae, Doidge.....	16
Natalia lucens, Harv. and Sond.....	64	Panicum.....	228, 263
Ochna, The Genus.....	87	" sp.....	68
" arborea.....	8, 88, 89, 90, 92	" arcuratum.....	265
" atropurpurea, D. C....	87, 88, 89, 90, 94	" arrectum, Nees.....	263
" atropurpurea, var. angustifolia,		" brizanthum, Hochst.....	262
Phillips.....	95	" coloratum, Linn.....	222, 265
" atropurpurea, var. natalita....	87, 88, 94	" deustum, Thunb.....	264
" Chilversii, Phillips.....	88, 90	" dregeanum, Nees.....	265
" Holstii, Engl.....	88, 89, 90, 93	" Ecklonii, Nees.....	265
" leptoclada, Aliv.....	88, 90, 94	" helopus var. glabrescens, K. Schum	261
" natalita, Engl. and Gilg....	88, 89, 90, 93	" Holubii, Stapf.....	263
" O'Connorii, Phillips.....	88, 89, 90, 92	" Isachne, Rott.....	262
" pretoriensis, Phillips.....	90, 95	" laevifolium, Hack.....	222, 264
" pulchra, Hook.....	87, 90, 91	" longijubatum, Stapf.....	265
Ocotea bullata.....	17, 205	" Marlothii, Hack.....	263
Old land grass.....	290	" maximum, Jacq....	62, 66, 68, 222, 264
Olea capensis.....	7, 11, 20, 22, 211	" miliaceum, L.....	265
" exasperata.....	11	" minus, Stapf.....	221, 265
" ? foveolata.....	27	" minus Stapf. var. planifolium,	
" laurifolia.....	11, 22, 73	Stapf.....	265
" peglerae.....	74	" natalensis, Hochst.....	222, 265
" Pegleri.....	211	" nigropedatum, Munro.....	262
" verrucosa.....	10, 13, 195	" novemnerve, Stapf.....	265
" woodiana.....	11, 20	" obscurans, Stapf.....	265
Oleaceae ?.....	74	" proliferum var. longijubatum, Stapf	265
Oligostroma maculiformis (Wint), Doidge..	31	" proliferum var. paludosum, Stapf.	265
" proteae, Syd.....	31	" serratum, Spreng.....	262
Olinia.....	97	Panicum stagninum, Retz.....	263
" The Genus.....	97	" trichopus, Hochst.....	261
" acuminata, Klotz... 97, 98, 100, 102,	104	Papilionaceae.....	40
" cymosa.....	97, 98, 99, 100, 104	Para grass.....	262
" cymosa var. acuminata.....	97, 100, 102	Parasterina brachystoma (Rehm.), Th., var.	
" cymosa var. intermedia.....	97	" laxa.....	201
" cymosa var. latifolia.....	97	Parasterina laxa, Doidge.....	201
" micrantha.....	97	" reticulata, Doidge.....	200
" radiata, Phill. and Hofmeyr 97, 98,		Parastigmataea, Doidge.....	22
100, 102, 103, 104		" nerviseta, Doidge.....	22
" ternata, Gilg.....	97	Parenglerula macowaniana (Thum.), v. Hohn	212
" usambarensis.....	97	Parinarium capense.....	26, 27
Olyra.....	235	Parinarium mobola.....	26
Oncoba Kraussiana.....	204	Parodiella schimperi, P. Henn.....	65
Ophiodothis edax, Sacc.....	32	Paspalum.....	226, 258
Ophiodothella edax (B. and Br.), V. Hohn	32	" digitaria, Poir.....	258
Oplismenus.....	227, 271	" dilatatum, Poir.....	258
" africanus.....	272	" distichum, Linn.....	258
" africanus, Beauv., var. simplex,		" scrobiculatum, L.....	220
Stapf.....	272	" scrobiculatum, L., var. Commer-	
" hirtellus, P. Beauv.....	272	sonii Stapf.....	258
Oropetium.....	229, 303	Pear—	
Oropetium capense, Stapf.....	303	" avocado.....	179, 180
Orthodanum.....	114, 117	" coffee (Pleurostylia).....	34
" argenteum, E. Mey.....	133	" real red.....	88, 93
" glabratum, Meisner.....	133	Pearl Millet.....	274
" latifolium, E. Mey.....	133	Peltophorum africanum.....	29
" Muhlenbeckii.....	133	Penhalonga grass.....	262
" sordidum, E. Mey.....	133	Pennisetum.....	227, 273
Oryza.....	228, 274	" benthami.....	68
		" cenchroides, Rich.....	222, 274

	PAGE		PAGE
Penisetum macrourum, Trin.....	274	Phyllachora caffra, Syd.....	27
" sphacelatum.....	68	" Chrysopogonis, Syd.....	219
" sphacelatum var. tenuifolium, Stapf.....	274	" crotonicola, Rehm.....	67
" spicatum, R. and S.....	274	" crotonis, (Cke.), Sacc.....	67
" Thunbergii, Kunth.....	274	" cynodontis (Sacc.), Niessl.....	67
" Thunbergii, var. Galpinii, Stapf	274	" dalbergiicola, P. Henn.....	25
" typhoideum.....	274	" Digitariae, Syd.....	220
Pentameris.....	231	" Doidgeae, Syd.....	220
Pentaschistis.....	231	" dombeyae, Syd.....	28
Perennial rye grass.....	303	" Evansii, Syd.....	67
Perischizon oleifolium (Kalch. and Cke.), Syd.....	22	" ficuum, Niessl.....	24
Perisporiales.....	212	" graminis, var. Cynodontis Sacc.....	67
Perisporium.....	209, 210	" grammica, P. Henn.....	26
Perisporium irenicolum, Doidge.....	209	" heterospora, P. Henn.....	68
" Meliolae.....	209, 210	" hieronymi, P. Henn.....	28
Perotis.....	229, 256	" Julocrotonis, Bres.....	67
Perotis latifolia, Ait.....	256	" Leptocarydii, Syd.....	220
Persea Americana.....	179, 180	" Lessertiae, Doidge.....	30
Phaeochorella parinari (P. Henn.), Theiss. and Syd.....	26	" lucens, (Cke.), Sacc.....	68
Phaeodimeriella.....	210	" melianthi (Thuem.), Sacc.....	28
Phaeodimeriella asterinicola, Doidge.....	210	" myrsinicola, Doidge.....	81
" capensis, Doidge.....	210	" osyridis, Cke.....	18
" plumbea, Doidge.....	211	" Peltophori, Syd.....	29
Phaeodothis stenostoma (Ell. and Tr.), Theiss. and Syd.....	66	" Penniseti, Syd.....	68
Phaeodothis Tristachyae, Syd.....	67	" Pterocarpi, Syd.....	25
Phaeostigma circumsedens, Doidge.....	210	" sanguinolenta, Th. and Syd., var. microspora.....	221
Phalaris.....	233, 275	" schweinfurthii, P. Henn. var. nervisequens, Lingelsh.....	24
" arundinacea, Linn.....	276	" stenostoma, Ell. and Tracy.....	66
" canariensis, L.....	276	" strelitziae, Sacc.....	31
" minor, Retz.....	276	" Teleae, Doidge.....	68
" paradoxa, Linn.....	276	" Tricholaenae, P. Henn.....	69
Phillips, E. P.—		" vivensis (Cke.), Sacc.....	27
The Genus Bersama.....	33	" Winkleri, Syd.....	220
A Revision of the South African Species of Sesbania.....	40	Phyllachorella rikatiensis, Doidge.....	30
The Natal Species of the Sapindaceae.....	57	Phyllanthus amapodensis.....	98
The Thorn Pears (Scolopia spp.).....	83	Physalospora caffra, Syd.....	27
The Genus Ochra.....	87	" cydoniae.....	179, 182
Species of Elephantorrhiza in the South African Herbaria.....	187	" cynodontis.....	67
Phillips, E. P., and Joan Hofmeyr—		" dombeyae, Syd.....	28
The Genus Olinia.....	97	" Perseae.....	182
The Genus Cyclopia, Vent.....	105	Placoasterella rehmi (P. Henn.), Theiss. and Syd.....	12
Phoberos.....	83	Plane, Cape.....	89, 93
" Ecklonii, Arn.....	86	Electronia obovata.....	98
" Mundtii, Arn.....	85	Pleistomella Halleriae, Doidge.....	17
" zeyheri, Arn.....	85	Pleurostelia (Coffee pear).....	34
Phragmites.....	233, 295	" capensis.....	200
" communis, Trin.....	295	Poa.....	234, 302
Phragmocaulis vivensis (Cke.), Theiss. and Syd.....	27	" annua, L.....	302
Phragmodothella nervisequens, Doidge.....	22	Poaagrostis.....	228
Phragmodothis asperata, Syd.....	23	Podocarpus elongata, Hk.....	217, 218
Phragmosperma.....	219	" falcata.....	218
" marattiae, (P. Henn.), Theiss and Syd.....	219	" latifolia.....	218
Phragmothyriella parenchymatica, Doidge.....	206	Pogonarthria.....	231, 295
Phyllachora.....	195	" falcata, Rendl.....	296
" aberiae.....	29	Pollinia villosa, Spreng.....	236
" albizziae, Cke.....	27	Polypogon.....	229
" amaniensis, P. Henn.....	29	Polyrhizon bewsii, Doidge.....	15
" anthistiricola, Syd.....	219	" Celastri, Doidge.....	16
" Brachystegiae, Doidge.....	221	Polystomella caulicola, Doidge.....	
		Polystomellaceae.....	
		Polystomellacearum.....	9, 15, 16
		Polytropia, Harvey and Sonder.....	114, 116
		" ferulaefolia.....	117
		" pinnata, Ecklon. and Zeyher.....	117

	PAGE		PAGE
Polytropia, <i>umbellata</i> , Ecklon. and Zeyher.	117	Rhynchosia [confusa, Burt-Davy 115, 118,	122, 125
Potamophila.....	233, 274	" congestiflora.....	113
" <i>prehensilis</i> , Benth.....	274	" connata.....	115
Prairie Grass.....	302	" <i>Cooperi</i> , Harvey.....	129
Prionachne.....	230	" <i>crassifolia</i> , Benth.....	113, 118, 127
Protea sp.....	32	" <i>cyanosperma</i> , Benth.....	114
" <i>abyssinica</i>	32	" <i>cyanosperma</i> , Benth.....	114, 115
" <i>flanagani</i>	31	" <i>densiflora</i> , D. C.....	115
" <i>grandiflora</i>	31, 71	" <i>Dieterlenae</i> , Bak. fil.....	118, 125
" <i>neriifolia</i>	31	" <i>elegans</i>	65
Pseudobromus.....	229, 278	" <i>elegantissima</i> , Schinz.....	113, 119, 132
" <i>africanus</i> , Stapf.....	279	" <i>ferulaefolia</i> , Benth.....	117
<i>Psoralea prostata</i>	117	" <i>ficifolia</i> , Benth.....	114, 138
<i>Psychotria capensis</i>	76	" <i>flavissima</i> , Hochst.....	124
<i>Pterocarpus angolensis</i>	25	" <i>Fleckii</i> , Schinz.....	113, 118, 122
" <i>ceriseus</i>	70	" <i>Galpini</i> , Bak. fil.....	119, 135
<i>Pterocelastrus variabilis</i>	198	" <i>gibba</i> , E. Mey.....	65, 118, 125
" <i>var. tricuspidatus</i>	198	" <i>gibba</i> var. <i>pictum</i> , Benth.....	125
<i>Ptaeroxylon</i> , E. and Z.....	57, 62	" <i>glabra</i> , D. C.....	132
" <i>utile</i> , E. and Z.....	62, 98	" <i>glabra</i> , Ecklon. and Zeyher.....	132
<i>Pueraria</i>	114	" <i>glandulosa</i> , D. C.....	119, 132
<i>Putterlickia pyracantha</i>	79, 200	" <i>glandulosa</i> , Ecklon and Zeyher	132
" <i>verrucosa</i>	200	" <i>graciliflora</i> , Harms.....	114, 131
<i>Pyrenacantha scandens</i>	208, 215	" <i>grandifolia</i> , Steud.....	120
<i>Quar</i>	98	" <i>grandifolia</i> , Walp.....	120
<i>Rapanea melanophleos</i>	81	" <i>Harmsiana</i> , Schlechter 113, 118, 123	
Real red pear.....	88, 93	" <i>Harmsiana</i> , Sch., var. <i>Burchellii</i> , Burt-Davy.....	114, 124
Reed canary grass.....	276	" <i>Harmsiana</i> , Sch., var. <i>grandifolia</i> , Bak. fil.....	118
Regte Kweek.....	288	" <i>Harmsiana</i> , var. <i>grandiflora</i>	124
Rescue or prairie grass.....	302	" <i>Harveyi</i>	118, 121
Revision of South African Species of <i>Sesbania</i>	40	" <i>hirsuta</i> , Ecklon. and Zeyher	113, 118, 125, 126
<i>Rhipidodendrum distichum</i> , Willd.....	141	" <i>hirsuta</i> var. <i>angustifolia</i> , Harvey.....	126
<i>Rhodes Grass</i>	290	" <i>Jacottetii</i> , Schinz.....	118, 123
<i>Rhoicissus thomboidea</i>	79	" <i>komatiensis</i> , Harms.....	113, 119, 134
<i>Rhynchosia</i>	113, 138	" <i>leucoscias</i> , Benth.....	116
<i>Rhynchosia</i> , Revision of South African Species.....	113	" <i>leucoscias</i> , var. <i>angustifolia</i> , Harvey.....	116
" <i>spp.</i>	65	" <i>longiflora</i> , Schinz.....	113, 119, 131
" <i>acuminata</i> , Ecklon. and Zeyher	125	" <i>longipes</i> , Harms.....	113, 118, 127
" <i>adenodes</i> , Ecklon. and Zeyher	114, 118, 121, 128, 137	" <i>memnonia</i> , D. C.....	137
" <i>adenodes</i> , Ecklon. and Zeyher	var. <i>Cooperi</i> , Bak. fil 118, 128, 129	" <i>memnonia</i> , Harvey.....	114, 127, 137
" <i>adenodes</i> , var. <i>laxiflora</i>	129	" <i>memnonia</i> , D. C., var. <i>prostrata</i> , Harvey.....	118, 127
" <i>adenodes</i> , Ecklon. and Zeyher,	var. <i>robusta</i>	" <i>microscias</i> , Benth.....	116
" <i>albissima</i> , Gandoger 114, 119, 127, 137	121	" <i>minima</i> , D. C.....	114, 126
" <i>amatymbica</i> , E. and Z.....	128	" <i>minima</i> , L.....	118
" <i>ambacensis</i> , K. Schum.....	134	" <i>monophylla</i> , Schlechter 113,	114, 118, 120
" <i>amboense</i> , Schinz.....	132	" <i>monophylla</i> , var. <i>Eylesii</i> , Bak. fil.....	120
" <i>angulosa</i> , Schinz.....	119, 136	" <i>namaensis</i> , Schinz.....	113, 114
" <i>angustifolia</i>	113	" <i>nervosa</i> , Benth.....	118, 124
" <i>angustiflora</i> , D. C.....	116	" <i>nervosa</i> , Benth, var. <i>pauciflora</i> , Harvey.....	122, 125
" <i>argentea</i> , Harvey.....	118, 126	" <i>nervosa</i> , forma <i>unifoliata</i> , Bak. fil.....	124
" <i>Bakeri</i> , Schinz.....	119, 135	" <i>nitens</i> , Benth.....	119, 135
" <i>bullata</i> , Benth.....	114, 119, 136	" <i>nitida</i> , Harvey.....	138
" <i>Burkei</i> , Burt-Davy.....	114, 118, 121	" <i>Orthodanum</i> , Benth.....	113, 119, 133
" <i>caribaea</i> , D. C.....	118, 125	" <i>Orthodanum</i> var. <i>densiflora</i>	133
" <i>var. pictam</i>	125	" <i>Orthodanum</i> var. <i>Muhlenbeckii</i> , Harvey.....	133
" <i>var. Thornecroftii</i>	125		
" <i>chrysantha</i> , Schlechter 113, 118, 129			
" <i>chrysoscias</i> , Benth.....	115, 116		
" <i>cinnamomea</i> , Schinz 113, 114,	119, 132		
" <i>clivorum</i> , S. Moore.....	113, 119, 134		
" <i>clivorum</i> , var. <i>longicalycina</i>	135		

	PAGE		PAGE
Rhynchosia ovata, Wood and Evans.	113,	<i>Schmidelia decipiens</i> , Arn.	61
" pauciflora, Bolus.	114, 115,	" <i>dregeana</i> , Sond.	59
" Pegleri, Bak. fil.	119, 137	" <i>erosa</i> , Arn. Sond.	60
" Pentheri, Schlechter	113, 118, 127, 128	" <i>melanocarpa</i> , Arn.	60
" Pentheri var. Hutchinsoniana,		" <i>monophylla</i> , Presl.	59, 79
" Burt-Davy and Bak. fil.	128	" <i>natalensis</i> , Sond.	57, 60
" pinnata, Harvey.	117	" <i>rehmanniana</i> , Szyszy.	61
" puberula, Harvey.	118, 130	Schmidtia.	222, 232, 293
" pycnantha, Harms.	135	" <i>bulbosa</i> , Stapf.	293
" quadrata, Harvey.	118, 122	Schneeppia.	6
" <i>Rehmanni</i> , Schinz.	138	" <i>brachylaenae</i> , Rehm.	6
" <i>remota</i> , Conrath.	132	" <i>radiata</i> , Doidge.	5
" reptabunda, N. E. Br.	118, 123	Schotia speciosa.	65
" rigidula, D. C.	113, 119, 133	Scleropoa.	234
" <i>riparia</i> , Ecklon and Zeyher.	132	Scolecodothis capensis, Doidge.	27
" rotundifolia, Walp.	117, 119	Scolecopeltis Cassipoureae, Doidge.	79
" Rudolfi, Harms.	119, 134	Scolopia.	83, 89, 93
" Schlechteri, Bak. fil.	119, 138	" spp.	77, 83
" <i>secunda</i> , Harvey.	118, 121	" Ecklonii, Szyszy.	83, 84, 86
" <i>secunda</i> (Thunberg).	118	" Ecklonii, var. Gerrardii, Phillips.	86
" <i>sericea</i> , Conrath.	132	" Ecklonii, var. Engleri, Phillips.	84, 86
" sigmodes, Benth.	114, 118, 129	" <i>Engleri</i> , Gilg.	83, 86
" simplicifolia, E. Mey.	117, 120	" Flanaganii, Sim.	83, 84
" spectabilis, Schinz.	114, 119, 134	" <i>Gerrardii</i> , Harv.	83, 86
" stenodon, Bak. Fil.	127	" Mundii, Warb.	83, 84, 85, 212
" Stuhlmanni, Harms.	115	" Thuneroftii, Phillips.	84
" Totta, D. C.	113, 114, 119,	" Zeyheri, Szyszy.	77, 83, 84
" 130, 131, 132,	138	Secale.	233
" Totta var. brevipetiolata, Bak fil.	131	Sehima.	225, 238
" Totta, var. Fenchelii, Schinz.	130	" <i>galpinii</i> , Stent.	239, 240
" Totta, var. graciliflora.	131	" <i>notata</i> .	239
" Totta, D. C., var. namaensis,		Sesbania.	40, 47
" Schinz.	130, 132	" Revision of South African Species	40
" Totta, var. pilosa, Harvey.	131	" <i>aculeata</i> , Pers.	42, 48, 50, 56
" Totta, var. unifoliata, Burt-		" <i>aegyptiaca</i> , Poir.	41, 44, 45
" Davy.	131	" <i>aegyptiaca</i> , var. bicolor, Wight	
" trichodes, Harvey.	138	" and Arn.	41, 45
" <i>uncinata</i> .	114, 119, 136	" <i>aegyptiacus</i> , Hiern.	50
" <i>uniflora</i> , Harvey.	113, 116	" <i>affinis</i> .	49
" <i>unifoliata</i> .	114	" <i>arabica</i> , Hochst.	42, 51
" <i>venulosus</i> , K. Schum.	114, 119,	" <i>atropurpurea</i> , Taub.	45
" 125, 132		" <i>caerulescens</i> .	41, 46
" viscidula, Steud.	114, 119, 137, 138	" <i>cinerascens</i> , Welw.	41, 47
" Woodii, Schinz.	113, 119, 136	" Dalzielii, Phillips and Hutch.	41, 49
<i>Rhytisma melianthi</i> (Thuem.), Sacc.	28	" Dummeri, Phill. and Hutch.	41, 45
" <i>porrigo</i> , Cke.	7	" Goetzei, Harms.	42
Rinorea ardesiaciflora.	80	" grandiflora, Poir.	41, 46
Rizalia confusa, Doidge.	211	" hamata, Phill. and Hutch.	42, 54, 55
Roodebosje.	98	" <i>Hildebrandtii</i> , Taub.	43
Rooi gras.	254	" <i>Hockii</i> , De Wild.	46
Rooihout.	89, 93	" Kirkii, Phill. and Hutch.	42, 54, 55
Rooi Yzerhout.	89, 93	" <i>leptocarpa</i> , D. C.	41, 50
Rosenscheldia horridula, Doidge.	20	" <i>macowaniana</i> , Schinz.	52
<i>Rottboellia compressa</i> , var. <i>fasciculata</i> Hack	241	" <i>macrantha</i> , Welw.	41, 47
Rye grass, Perennial.	303	" <i>microphylla</i> , Harms.	42, 52
Sacciolepis.	227, 228	" <i>mossaubicensis</i> , Klotzsch.	42, 52
Sand quick.	294	" <i>pachycarpa</i> , D. C.	42, 50
Sapindaceae, The Natal Species of.	57	" <i>phaerocarpus</i> , Hiern.	50
Sapindus, Linn.	57, 62	" <i>pubescens</i> , D. C.	41, 48, 49
" oblongifolius, Sond.	62	" <i>punctata</i> , D. C.	41, 43, 44, 45
Schismus.	232	" <i>punctata</i> , Hiern.	47, 50
Schizachrium.	225, 245	" Rogersii, Phill. and Hutch.	42, 55
" semiberbe, Nees.	246	" <i>sericea</i> .	49
<i>Schmidelia africana</i> , D. C. Harv. and Sond.	60	" <i>speciosa</i> , Taub.	41, 43
		" <i>sphaerocarpa</i> .	42, 51
		" <i>sphaerosperma</i> , Welw.	51
		" <i>tetraptera</i> , Hochst.	42, 53, 54, 55

	PAGE		PAGE
Sesbania Wildemannii, Phill. and Hutch..	41, 49	Sweet vernal grass.....	276
Setaria.....	227, 272	Tambookie grass.....	222, 251
" aurea, A. Brown.....	67, 273	Tawri.....	51
" flabellata, Stapf.....	273	Tea—	
" Gerrardii, Stapf.....	273	Bush.....	105
" Holstii, Stapf.....	273	common bush.....	106
" imberbis, R. and S.....	273	Heidelberg.....	106
" italica, Beauv.....	273	heuning.....	106
" lindenbergiana, Stapf.....	272	honey.....	105, 106
" nigrirostris, Dur. and Schinz 67, 222,	272	Hottentots.....	106
" perennis, Hack.....	273	sugar.....	106
" sulcata, Raddi.....	67, 222, 272	vici.....	105
" verticillata, Beauv.....	273	Teclea natalensis.....	68
Seynesia elegantula, Syd.....	10	Teff.....	301
Sibomvu.....	88	Telimena Arundinariae, Doidge.....	69
Signodostyles villosa, Benth.....	129	" corticicola, Doidge.....	70
Sinande.....	261	Tephrosia elongata.....	32
Sneeze-wood.....	98	" suberosa.....	32
" bastard.....	34	Teratospaeria fibrillosa, Syd.....	71
Sorghum.....	225, 243	Tetrachne.....	231
" ethiopicum, Rupr. ?.....	243	Tetrapterae.....	55
" halepense, Nees.....	243	Theissenula.....	212
" micratherum, Stapf.....	243	" Woodiana, Doidge.....	212
" versicolor, Stapf.....	243	Themeda.....	225, 254
South African Millet Grass.....	273	" triandra, Forsk.....	222, 254
Spartina.....	230	" triandra, Forsk, var. Burchellii..	254
Spear grass.....	253	" triandra, Forsk, var. glauca.....	254
Sphaeria examinans, Mont. and Berk.....	21	" triandra, Forsk, var. hispida.....	254
Sporobolus.....	228, 279	" triandra, var. punctata.....	254
" centrifugus, Nees.....	279	Thorn Pears, The (Scolopia spp.).....	83
" discosporus, Nees.....	279	Toddalia lanceolata.....	209
" festivus, Hochst., var. fibrous,		" natalensis.....	79
Stapf.....	279	Trabutia evansii, Theiss. and Syd.....	24
" festivus, var. stuppeus, Stapf.....	279	" ficuum (Neissl.), Theiss. and Syd	24
" fimbriatus, Nees.....	279, 281	" nervisequens (Lingelsh), Theiss.	
" fimbriatus, Nees., var. latifolia..	279	and Syd.....	24
" indicus, R. Br., var. laxus, Stapf 222,	281	" nervisequens, var. robusta.....	25
" Ludwigii, Hochst.....	222, 279	Trachypogon.....	224, 241
" nitens, Stent.....	281, 282	" plumosus, Nees.....	242
" pectinatus, Hack.....	281	" polymorphus, var. capensis, Hack.	242
" pyramidalis, Beauv.....	281	Tragus.....	255
" Rehmanni, Hack.....	281	" Koeleriodes, Aschers.....	255
" robustus, Kunth.....	281	" racemosus, All.....	255
" Smutsii, Stent.....	223, 280, 281,	Trichelia emetica.....	7
" tenellus.....	222	Trichocladus ellipticus.....	15, 78, 79
Steek gras.....	253	Trichocycelus, N. E. Br.....	151
Stenotaphrum.....	226	" ciliatus, N. E. Br.....	151
Stent, Sydney M—		Tricholaena.....	228, 269
South African Graminae—		" glabra, Stapf.....	271
A New Genus and Seven New		" rosea, Nees.....	271
Species.....	170	" setifolii, Stapf.....	271
Graminae of the Transvaal.....	222	Trichopeltaceae.....	79
Key to the South African Genera	224	Trichopeltula carissae, Doidge.....	78
Stephania hernandifolia.....	22	" kentaniensis, Doidge.....	79
Stiburus.....	232, 301	Trichothyrium dubiosum.....	216
" alopecurioides, Stapf.....	301	" elegans, Doidge.....	213
Stigmathea.....	22	" robustum, Doidge.....	215
" Rhynchosiae, Kalch. and Cke... 65		Tricopteryx.....	231, 287
Stigmatacearum.....	22	" sp. A.....	287
Stink gras.....	300	" flavida, Stapf.....	287
Stipa.....	229	" simplex, Hack.....	287
" dregeana, var. elongata.....	74	" simplex, Hack., var. crinita,	
Strelitzia angusta.....	31	Stapf.....	287
Streptocarpus Rexii.....	203, 216	" simplex, Hack., var. Sericea,	
Striga lutea.....	264	Stapf.....	287
Sugar Tea.....	106	Tripogon.....	230, 291
Systemma Pterocarpi, Doidge.....	70		

	PAGE		PAGE
<i>Tripogon abyssinicus</i> , Nees.....	291	<i>Urochloa helopus</i> , Stapf.....	261
<i>Tripopora Cookei</i> , (Cke.), Sacc.....	216	" <i>pullulans</i> , Stapf., var. <i>mossambi-</i>	
" <i>tripos</i> (Cke.), Lind.....	216, 217	<i>censis</i> , Stapf.....	261
<i>Triraphis</i>	233, 294	" <i>trichopus</i> , Stapf.....	261
" <i>Rehmanni</i> , Hack.....	294	<i>Vigna</i> sp.....	65
" <i>Schlechteri</i> , Pilg.....	294	<i>Virgilia capensis</i>	98
<i>Trisetum</i>	234	<i>Viscum nervosum</i>	8
<i>Tristachya</i>	231, 285	<i>Vlei tea</i>	105
" <i>biseriata</i> , Stapf.....	285	<i>Vulpia</i>	234
" <i>leucothrix</i> , Trin.....	67, 287	<i>Water kweek</i>	275
" <i>pallida</i> , Stent.....	285, 286	<i>Warmbaths grass</i>	262
" <i>pedicellata</i> , Stent.....	178, 287	<i>Wildebeeste gras</i>	242
" <i>Rehmanni</i> , Nees.....	178, 285	<i>Xylosma Flanaganii</i> , Bolus.....	84
<i>Turf-grass</i>	238	<i>Xymalos</i>	10
<i>Turpentine grass</i>	248	" <i>monospora</i>	79, 206
<i>Umbomvane umncince</i>	88	<i>Xyochlaena</i> sp.....	228, 271
<i>Umpanzi</i>	98	" <i>monachne</i> , Stapf.....	271
<i>Umpanzite</i>	98	<i>Yorkshire Fog</i>	284
<i>Umtisa Listeriana</i>	79	<i>Zamarke</i>	50
<i>'Un Tshaba</i>	251	<i>Zukalia</i>	212
<i>Urelytrum</i>	224, 225, 239	" <i>parenchymatica</i> , Doidge.....	206, 212
" <i>squarrosus</i> , Hack.....	241	" <i>Woodiana</i> , Doidge.....	212
<i>Urochlaena</i>	232	<i>Zuur pol</i>	242
<i>Urochloa</i>	227, 260		
" <i>bulbodes</i> , Stapf.....	261		

